

Q T
104
B886c
1886

QUIZ-COMPENDS

PHYSIOLOGY

DR. BRUBAKER.



NLM 00052909 3

BERTS' of Medicine.

THE FIFTH ENLARGED EDITION. JUST READY.

Recommended as a Text-book at University of Pennsylvania, Long Island College Hospital, Yale and Harvard Colleges, Bishop's College, Montreal, University of Michigan, and over twenty other Medical Schools.

A HANDBOOK OF THE THEORY AND PRACTICE OF MEDICINE.
By FREDERICK T. ROBERTS, M.D., M.R.C.P., Professor of Materia Medica and Therapeutics and of Clinical Medicine in University College Hospital; Assistant Physician in Brompton Consumptive Hospital. The Fifth Edition, partially rewritten, and carefully revised throughout.

Price, in Cloth, \$3.00; Leather, \$6.00.

SURGEON GENERAL'S OFFICE	
LIBRARY	
ANNEX	
Section	Physiology
Form 113c W.D.,S.G.O.	No. 291336

U. S. GOVERNMENT PRINTING OFFICE: 1928

sy:
501tio
co
A:co
Y:tal
in
M

ad

be known. — Professor J. A. Tucker, Cincinnati.

"I sha' include it among the text-books of the College of Physicians and Surgeons, and strongly recommend it to my classes."—*Professor John S. Lynch, Baltimore.*

"It is unsurpassed by any work that has fallen into our hands, as a compendium for students preparing for examination. It is thoroughly practical, and fully up to the times."—*The Clinic.*

"Our opinion of it is one of almost unqualified praise. The style is clear, and the amount of useful and, indeed, indispensable information which it contains is marvelous. We heartily recommend it to students, teachers, and practitioners."—*Boston Medical and Surgical Journal.*

"That Dr. Roberts' book is admirably fitted to supply the want of a good handbook of medicine, so much felt by every medical student, does not admit of a question."—*Students' Journal and Hospital Gazette.*

P. BLAKISTON, SON & CO., Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

YEO'S Manual of Physiology.

300 ILLUSTRATIONS.

FULL GLOSSARY AND INDEX.

By GERALD F. YEO, M.D., F.R.C.S., Professor of Physiology in King's College, London. Demi-octavo, 750 pages. Over 300 carefully printed engravings on wood.

Bound in Cloth, \$4.00; Leather, \$5.00.

RECOMMENDATIONS.

"By his excellent manual Prof. Yeo has supplied a want which must have been felt by every teacher of physiology. In the noble text-book of Prof. Foster, English readers have a work which is unsurpassed, but its great size and comprehensiveness, and its somewhat minute discussion of many doubtful points, make it a formidable object to the eyes of the first and second years' students. Dr. Yeo has written a book which is intended for junior students, but which, although written in simple, and as far as possible untechnical language, is accurate and complete. * * * Moreover, being intended chiefly for medical students, and written by one who is not only an able physiologist, but an accomplished physician, the needs of the physician and surgeon are never lost sight of. * * * The text is profusely illustrated with excellent wood engravings. * * * In conclusion, we heartily congratulate Prof. Yeo on his work, which we can recommend to all those who wish to find, within a moderate compass, a reliable and pleasantly written exposition of all the essential facts of physiology as the science now stands."—*The Dublin Journal of Medical Science*, May, 1884.

"For students' use it is one of the very best text-books in Physiology."—*Prof. L. B. How, Dartmouth Medical College, Hanover, N. H.*

"The work will take a high rank among the smaller text-books of Physiology."—*Prof. H. P. Bowditch, Harvard Medical School.*

"The brief examination I have given it was so favorable that I placed it in the list of text-books recommended in the circular of the University Medical College."—*Prof. Lewis A. Stimpson, M.D., 37 East 37th Street, New York.*

"There are many points in physiology that are either not comprehended or are misunderstood by the great majority of students. In this work these points are made especially clear, and in a particular manner those that are of most importance to the medical or dental practitioner. We have had long experience in teaching this branch of medical science, and unreservedly commend this work to the student of physiology."—*Archives of Dentistry.*

"It is an excellent book and well adapted for the uses for which it is intended. It is a decidedly modern book, being carefully pruned of all ancient redundancies and containing all that is new and proven. The arrangement is very good, indeed, the best, and corresponds closely with that of Dalton's. It is written in simple, pure English. * * * It will be valuable for students."—*D. Tod Gilliam, M.D., Professor of Physiology, Starling Medical College, Columbus, O.*

"After a careful examination of this Manual of Physiology, I can truthfully say that it is a most valuable addition to the list of text-books upon the subject. That it should and will receive a welcome from both students and teachers there can be no doubt; for in addition to the familiar but well presented facts of most text-books, it contains all the most important facts of physiological science which have been established in the last few years. The author presents his subject in a manner that is clear, concise and logical. Each section has had a careful revision, and reveals the author's familiarity with the scope and tendencies of modern physiology. It will prove an interesting and instructive book to those commencing the study of this subject."—*A. P. Brubaker, M.D., Demonstrator of Physiology, Jefferson Medical College, Philadelphia.*

P. BLAKISTON, SON & CO., Publishers and Booksellers.

1012 WALNUT STREET, PHILADELPHIA.

THE FIFTH EDITION.

Holden's Anatomy.

208 WOOD ENGRAVINGS.

A MANUAL OF THE DISSECTION OF THE HUMAN BODY. By LUTHER HOLDEN, M.D., Late President of the Royal College of Surgeons of England; Consulting Surgeon to St. Bartholomew's Hospital. Fifth Edition. Edited by JOHN LANGTON, M.D., F.R.C.S., Surgeon to, and Lecturer on Anatomy at, St. Bartholomew's Hospital; Member of the Board of Examiners, Royal College of Surgeons of England; With 208 fine Wood Engravings. Octavo. About 880 pages.

Bound in Cloth, \$5.00; Leather, \$6.00.

* * * This new edition has been revised with the object of making it a more thorough text-book. Over one hundred pages new matter have been added, and many new illustrations, some of which are very finely engraved and printed. Notwithstanding these new features, the price of the book has been lowered.

GILLIAM'S ESSENTIALS OF PATHOLOGY.

JUST PUBLISHED.

The object of this book is to explain to the student, in a plain, practical way, the fundamentals of Pathology, as an introduction to larger books.

THE ESSENTIALS OF PATHOLOGY. By D. TOD GILLIAM, M.D., Professor of Physiology, Starling Medical College, formerly Professor of General Pathology, Columbus Medical College, Columbus, Ohio. 12mo. 296 pages. 47 Illustrations.

Price, Cloth, \$2.00.

P. BLAKISTON, SON & CO., Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

HUMAN PHYSIOLOGY.

THIRD EDITION. ILLUSTRATED.

BRUBAKER.

? QUIZ COMPENDS ?

A NEW SERIES OF MANUALS FOR THE USE OF
STUDENTS AND PHYSICIANS.

Price of each, Cloth, \$1.00. Interleaved, for taking Notes, \$1.25.

These Compends are based on the most popular text-books, and the lectures of prominent professors, and are kept constantly revised, so that they may thoroughly represent the present state of the subjects upon which they treat.

The Authors have had large experience as Quiz Masters and attachés of colleges, and are well acquainted with the wants of students.

They are arranged in the most approved form, thorough and concise, with illustrations whenever they can be used to advantage.

Can be used by students of *any* college.

They contain information nowhere else collected in such a condensed, practical shape.

Size is such that they may be easily carried in the pocket, and the price is low.

They will be found very serviceable to physicians as remembrancers.

LIST OF VOLUMES.

No. 1. **ANATOMY.** Third Edition. 63 Illustrations. By SAMUEL O. L. POTTER, M.D., late A. A. Surgeon U. S. Army.

No. 2. **PRACTICE OF MEDICINE.** Part I. Second Edition. Revised and Enlarged. By DAN'L E. HUGHES, M.D., Demonstrator of Clinical Medicine, Jefferson College, Philadelphia.

No. 3. **PRACTICE OF MEDICINE.** Part II. Second Edition. Revised and Enlarged. Same author as No. 2.

No. 4. **PHYSIOLOGY.** Third Edition, with Illustrations. Enlarged and Revised. By A. P. BRUBAKER, M.D., Demonstrator of Physiology, Jefferson Medical College, Philadelphia.

No. 5. **OBSTETRICS.** Second Edition. Enlarged. By HENRY G. LANDIS, M.D., Professor of Obstetrics and Diseases of Women and Children, Starling Medical College, Columbus, Ohio. Illustrated.

No. 6. **MATERIA MEDICA.** A New Revised Edition. By SAMUEL O. L. POTTER, M.D., late A. A. Surgeon U. S. Army.

No. 7. **INORGANIC CHEMISTRY.** Revised Edition. By G. MASON WARD, M.D., Demonstrator of Chemistry, Jefferson College, Philadelphia.

No. 8. **VISCERAL ANATOMY.** Second Edition. Revised. By SAMUEL O. L. POTTER, M.D., late A. A. Surgeon, U. S. Army. With Illustrations.

No. 9. **SURGERY.** Second Edition. Revised and Enlarged. By ORVILLE HORWITZ, B.S., M.D., Resident Physician at Pennsylvania Hospital, Philadelphia. With 62 Illustrations.

No. 10. **ORGANIC CHEMISTRY.** Including Medical Chemistry, Urine Analysis and the Analysis of Water and Food. By HENRY LEFFMANN, M.D., Demonstrator of Chemistry in Jefferson College, Philadelphia.

No. 11. **PHARMACY.** By F. E. STEWART, M.D., PH.G., Quiz Master in Pharmacy and Chemistry, Philadelphia College of Pharmacy, Lecturer at the Medico Chirurgical College, etc.

Others in preparation.

Price, each, Cloth, \$1.00. Interleaved, for taking Notes, \$1.25.

P. BLAKISTON, SON & CO.,

MEDICAL PUBLISHERS AND BOOKSELLERS,

1012 WALNUT ST., PHILADELPHIA.

A

COMPEND

OF

HUMAN PHYSIOLOGY.

ESPECIALLY ADAPTED FOR THE USE OF
MEDICAL STUDENTS.

BY

ALBERT P. BRUBAKER, A.M., M.D.,

DEMONSTRATOR OF PHYSIOLOGY IN THE JEFFERSON MEDICAL COLLEGE; PROFESSOR
OF PHYSIOLOGY, PENNSYLVANIA COLLEGE OF DENTAL SURGERY;
MEMBER OF THE PATHOLOGICAL SOCIETY
OF PHILADELPHIA.

THIRD EDITION, REVISED AND ENLARGED.

WITH ILLUSTRATIONS,

AND


A TABLE OF PHYSIOLOGICAL CONSTANTS.

PHILADELPHIA:

P. BLAKISTON, SON & CO.,

1017 WALNUT STREET.

1886.



Annex

QT

104

B886c


1886

File # 3680, no. 2

Entered according to Act of Congress, in the year 1886, by

P. BLAKISTON, SON & CO.,

In the office of the Librarian of Congress, at Washington, D. C.



PRESS OF WM. F. FELL & CO.,
1220-24 Sansom street.

PREFACE TO SECOND EDITION.

This Compend of Physiology is the outgrowth of the author's system of examinations in the Quiz room during a number of years, and was written at the request of medical students who desired a compact and convenient arrangement of the fundamental facts of human physiology. As most medical students enter upon the study of physiology before they have acquired a thorough knowledge of anatomy, it was thought desirable that such anatomical details should also be inserted as would be essential to a clear conception of the functions about to be studied. It was believed that it would be practically useful to students during their attendance upon lectures and in reviewing the subject prior to examinations. The fact that during the first year after its publication the first edition has been exhausted, proves that it has met the needs of students.

In preparing a second edition the author has carefully revised the entire work, and inserted some fifteen pages of additional matter, which it is hoped will still further increase the usefulness of the book.

To those teachers of physiology who have kindly noticed and recommended the Compend to their students I tender my thanks, and trust that in its improved condition it will continue to merit their approval.

ALBERT P. BRUBAKER.

PREFACE TO THIRD EDITION.

A third edition of the Compend having been called for, the author has taken the opportunity to make some alteration to the text, to add some new material, and to insert a number of illustrations, which it is hoped will elucidate the text.

1210 Race Street.

A. P. B.

February, 1886.

TO MY FATHER,
HENRY BRUBAKER, A.M., M.D.,
THIS LITTLE VOLUME
IS AFFECTIONATELY INSCRIBED.

TABLE OF CONTENTS.

	PAGE
INTRODUCTION.....	9
CHEMICAL COMPOSITION OF THE BODY.....	10
STRUCTURAL COMPOSITION OF THE BODY.....	15
FOOD.....	18
DIGESTION.....	23
ABSORPTION.....	33
BLOOD.....	39
CIRCULATION OF BLOOD.....	45
RESPIRATION.....	52
ANIMAL HEAT.....	58
SECRETION.....	60
Mammary Glands.....	63
Vascular or Ductless Glands.....	65
EXCRETION.....	66
Kidneys.....	66
Liver.....	73
Skin.....	76
NERVOUS SYSTEM.....	80
Spinal Nerves.....	82
Properties and Functions of Nerves.....	84
Cranial Nerves.....	87
Spinal Cord.....	101
Medulla Oblongata.....	109
Pons Varolii.....	112
Crura Cerebri.....	113
Corpora Quadrigemina.....	114
Corpora Striata and Optic Thalami.....	114

	PAGE
Cerebellum	116
Cerebrum.....	118
Sympathetic Nervous System.....	123
SENSE OF TOUCH	126
SENSE OF TASTE.....	127
SENSE OF SMELL.....	129
SENSE OF SIGHT.....	130
SENSE OF HEARING.....	136
VOICE AND SPEECH	141
REPRODUCTION.....	144
Generative Organs of the Female.....	144
Generative Organs of the Male.....	147
Development of Accessory Structures.....	148
Development of the Embryo.....	153
TABLE OF PHYSIOLOGICAL CONSTANTS.....	159
TABLE SHOWING RELATION OF WEIGHTS AND MEASURES OF THE METRIC SYSTEM TO APPROXIMATE WEIGHTS AND MEASURES OF THE U. S.....	162
INDEX	163

COMPEND

OF

HUMAN PHYSIOLOGY.

Physiology, from φύσις, nature, and λόγος, a *discourse*, in its original application embraced the study of all natural objects, inorganic as well as organic. In its modern application physiology signifies *the study of life*; the investigation of the vital phenomena exhibited by all organic bodies, vegetable and animal.

It may be divided into—

1. *Vegetable physiology*, which treats of the phenomena manifested by the several structures of which the plant is composed.
2. *Animal physiology*, which treats of the phenomena manifested by the organs and tissues of which the animal body is composed.

Human Physiology is the study of the functions exhibited by the human body in a state of health.

A *Function* is the action of an organ or tissue.

The Functions of the Human Body may be classified into three groups, viz.:—

1. *Nutritive functions*, which have for their object the preservation of the individual; *e.g.*, digestion, absorption, circulation of the blood, respiration, assimilation, animal heat, secretion and excretion.
2. *Animal functions*, which bring the individual into conscious relationship with external nature; *e.g.*, sensation, motion, language, mental and moral manifestations.
3. *Reproductive function*, which has for its object the preservation of the species.

The facts of human physiology have been determined by means of anatomy, chemistry, pathology, comparative anatomy, vivisection, the application of physics, etc.

The Body may be studied from a *chemical* and *structural* point of view.

CHEMICAL COMPOSITION OF THE HUMAN BODY.

Of the **Sixty-four Chemical Elements**, about *sixteen* enter into the composition of the body, in the following proportions:—

Oxygen.....72.00	}	O. H. and C. are found in all the tissues and fluids of the body, without exception.
Hydrogen..... 9.10		
Nitrogen..... 2.50		O. H. C. and N. found in most of the fluids and all tissues except fat.
Carbon13.50		
Sulphur..... .147	In fibrin, casein, albumen, gelatin; as potassium sulpho-cyanide in saliva; as alkaline sulphate in urine and sweat.
Phosphorus.... 1.15	In fibrin and albumen; in brain; as tri-sodium phosphate in blood and saliva, etc.
Calcium..... 1.30	As calcium phosphate in lymph, chyle, blood, saliva, bones and teeth.
Sodium..... .10	As sodium chloride in all fluids and solids of the body, except enamel; as sodium sulphate and phosphate in blood and muscles.
Potassium..... .026	As potassium chloride in muscles; generally found with sodium as sulphates and phosphates.
Magnesium.... .001	Generally in association with calcium, as phosphate, in bones.
Chlorine..... .085	In combination with sodium, potassium and other bases, in all the fluids and solids.
Fluorine..... .08	As calcium fluoride in bones, teeth and urine.
Iron..... .01	In blood globules; as peroxide in muscles.
Silicon.....a trace	In blood, bones and hair.
Manganese, a trace	Probably in hair, bones and nails.

Of the four chief elements which together make up *97 per cent.* of the body, O. H. N. are eminently *mobile, elastic*, and possess great *atomic heat*. C. H. N. are distinguished for the narrow range and feebleness of their affinities and chemical inertia. C. has the greatest atomic cohesion. O. is noted for the number and intensity of its combinations, and its remarkable display of chemical activity.

Chemical Elements do not exist alone in the body, but are combined in characteristic proportions to form compounds, the *proximate principles*, which are the ultimate compounds to which the fluids and solids can be reduced.

Proximate Principles exist in the body under their own form, and can be extracted without losing their distinctive properties.

There are about *one hundred* proximate principles, which are divided into four classes, viz.: *inorganic, organic non-nitrogenized, organic nitrogenized, and principles of waste.*

I. INORGANIC PROXIMATE PRINCIPLES.

SUBSTANCE.	WHERE FOUND.
Oxygen.....	Lungs and blood.
Hydrogen.....	Stomach and intestines.
Nitrogen.....	Blood and intestines.
Carbonic anhydride.....	Expired air of lungs.
Carburetted hydrogen	} ... Lungs and intestines.
Sulphuretted hydrogen	
Water.....	Found in all solids and fluids.
Sodium chloride.....	In all fluids and solids except enamel.
Potassium chloride.....	In muscles, liver, saliva, gastric juice, etc.
Ammonium chloride.....	Gastric juice, saliva, tears, urine.
Calcium chloride.....	Bones, teeth, urine.
Calcium carbonate.....	Bones, teeth, cartilage, internal ear, blood.
Calcium phosphate	} ... In all fluids and solids of the body.
Magnesium phosphate	
Sodium phosphate	
Potassium phosphate	
Sodium sulphate	} ... Universal, except milk, bile and gastric juice.
Potassium sulphate	
Sodium carbonate	} ... Blood, bones, lymph, urine, etc.
Potassium carbonate	
Magnesium carbonate.....	Blood and sebaceous matter.

The inorganic principles enter and leave the body under their own form. *Water* is an essential constituent of all the tissues of the body, constituting about 70 per cent. of the entire body weight. It is introduced into the body in the form of drink and as a constituent of all kinds of food. The average quantity consumed daily is about four pints. While in the body, water acts as a general solvent, gives pliability to various tissues, and promotes the passage of inorganic and organic matters through animal membranes. It also promotes chemical changes which are essential to absorption and assimilation of food and the elimination of products of waste. It is probable that water is also formed within the body by the

union of oxygen with the surplus hydrogen of the food. It is eliminated by the skin, lungs and kidneys.

Sodium chloride is present in all the solids and fluids of the body, with the exception of enamel. It regulates osmotic action, holds the albuminous principles of the blood in solution, and preserves the form and consistence of blood corpuscles and the cellular elements of the tissues, by regulating the amount of water entering into their composition.

Calcium phosphate is the most abundant of all the inorganic principles with the exception of water, and is present to a great extent in bone, teeth, muscles and milk. It gives the requisite consistency and solidity to the different tissues and organs. In the blood, it is held in solution by the albuminous constituents.

The *Sodium* and *Potassium phosphates* are present in most of the solids and fluids, and give to them their alkaline reaction. They are chiefly derived from the food.

II. ORGANIC NON-NITROGENIZED PRINCIPLES.

The organic non-nitrogenized principles are derived mainly from the vegetable world, but are also produced within the animal body. They are divided into: 1st the *carbo-hydrates*, comprising starch and sugar, bodies in which the oxygen and hydrogen exist in the proportion to form water, the amount of carbon being variable; 2d, the *hydro-carbons*, comprising fats, bodies having the same elements entering into their composition, but with the carbon and hydrogen increased and the oxygen diminished in amount.

SUGARS. C. O. H.

Glycogen, or Liver sugar.

Lactose, or Milk sugar.

Glucose, or Grape sugar.

Inosite, or Muscle sugar.

Sugar is found in many of the tissues and fluids of the body; *e.g.*, liver, milk, placenta, blood, muscles, etc. The varieties of sugar are soluble in water, assume the crystalline form upon evaporation, and are converted into alcohol and carbonic acid by fermentation. Sugar is derived from the food, absorbed into the blood, where it largely disappears. After playing its part in the nutritive processes of the body, it is oxidized, and thus contributes to the formation of heat. It is finally eliminated under the form of carbonic acid and water. There is no experimental proof that sugar contributes directly to the formation of fat in the animal body.

FATS. C. O. H.

Palmitin,	} Neutral Fats.	Palmitic acid,	} Fatty Acids.
Stearin,		Stearic acid,	
Olein,		Oleic acid,	

The *Neutral fats*, when combined in proper proportions, constitute a large part of the fatty tissue of the body; they are soluble in ether, chloroform and hot alcohol; insoluble in cold alcohol and water, and liquefy at a high temperature; when a neutral fat is subjected to a high temperature in the presence of water and an alkali, it is decomposed, with the assimilation of water, into a fatty acid and glycerine. The fatty acid combines with the alkali and forms an oleate, palmitate or stearate, according to the fat used. A similar decomposition of the neutral fats is said to take place in the small intestine during digestion. When thoroughly mixed with pancreatic juice, the fats are reduced to a condition of *emulsion*, a state in which the fat is minutely subdivided and the small globules held in suspension.

The *Fatty acids* combined with sodium, potassium and calcium, are found as salts in various fluids of the body, such as blood, chyle, fæces, etc. Phosphorized fats in nervous tissue, butyric acid in milk, propionic acid in sweat, are also constituents of the body.

The Fats are derived from the food, both animal and vegetable. They are deposited in the form of small globules in the cells of the different tissues, are suspended in various fluids, are deposited in masses in and around various anatomical structures and beneath the skin. Independent of the fat consumed as food, there is good experimental evidence that fat is also produced within the animal body from a partial decomposition of the albuminous compounds. Fat serves as a non-conductor of heat, gives roundness and form to the body, and protects various structures from injury. The fats are ultimately oxidized, thus giving rise to heat and force, and are finally eliminated as carbonic acid and water.

III. ORGANIC NITROGENIZED PRINCIPLES.

ALBUMENS. C. O. H. N. S. P.

Albumen.	Myosin.	Mucin.
Albuminose.	Protagon.	Chondrin.
Fibrin.	Pepsin.	Elastin.
Casein.	Pancreatin.	Keratin.
Ostein.	Salivin.	Globulin.

The *Albuminous compounds* are organic in their origin, being derived from the animal and vegetable world; they are taken into the body as

food, appropriated by the tissues, and constitute their organic basis; they differ from the non-nitrogenized substances in not being crystalline, but amorphous, in having a more complex but just as definite composition, and containing in addition to C. O. H., nitrogen, with, at times, sulphur and phosphorus. The albumens possess characteristics which distinguish them from all other substances: viz., a *molecular mobility*, which permits *isomeric* modifications to take place with great facility. Under favorable conditions they promote chemical changes, by their presence (*catalysis*), in other substances: *e. g.*, during digestion, salivin and pepsin cause starch and albumen to be transformed into sugar and albuminose respectively. Different albumens possess varying proportions of water, which they lose when subjected to desiccation, becoming solid; but upon exposure to moisture they again absorb water, regaining their original condition—they are *hygroscopic*. Another property is that of *coagulation*, which takes place under certain conditions: *e. g.*, the presence of mineral acids, heat, alcohol, etc.

After death the albuminous compounds undergo putrefactive changes, giving rise to carburetted and sulphuretted hydrogen and other gases.

Albumen exists in the blood, lymph, chyle, constituting the pabulum of the tissues; it is coagulated by heat, mineral acids and alcohol.

Peptones are formed in the stomach from the digestion of albuminous principles of the food; they are coagulated by tannic acid, chlorine, acetate of lead, and characterized by great diffusibility, which permits them to pass through animal membranes with facility.

Fibrin can be obtained from freshly drawn blood by whipping; it also coagulates spontaneously, and when examined microscopically exhibits a filamentous structure.

Casein is the albuminous principle of milk.

Ostein constitutes the organic basis of bone, with which are mingled the salts of lime.

Myosin is found in muscles, *protagon* in brain, *pepsin*, *pancreatin* and *salivin*, in the digestive fluids.

Mucin, *chondrin*, *elastin*, *keratin* and *globulin*, are found in mucus, cartilage, elastic tissues, hair, nails, and red corpuscles, respectively.

As the properties of the compounds formed by the union of elements are the resultants of the properties of the elements themselves, it follows that the ternary substances, sugars, starches and fats, possess a great inertia and a notable instability; while in the more complex albuminous compounds, in which sulphur and phosphorus are united to the four chief elements, molecular mobility, resulting in isomerism, exists in a high degree. As

these compounds are unstable, of a great molecular mobility, they are well fitted to take part in the composition of organic bodies, in which there is a continual movement of composition and decomposition.

IV. PRINCIPLES OF WASTE.

Urea,	Xanthin,	Sodium,	} Urates.
Creatin,	Tyrosin,	Potassium,	
Creatinin,	Hippuric Acid,	Ammonium,	
Cholesterin,	Calcium Oxalate,	Calcium,	

These principles, which represent waste, are of organic origin, arising within the body as products of disassimilation or retrograde metamorphosis of the tissues; they are absorbed by the blood, carried to the various excretory organs, and by them eliminated from the body.

The excrementitious substances will be fully considered under excretion.

Proximate Quantity of the Chemical Elements and Proximate Principles of Body Weighing 154 lbs.

	lbs.	oz.		lbs.	oz.
Oxygen.....	111	..	Water	111	..
Hydrogen.....	14	..	Albuminoids.....	23	7
Nitrogen.....	3	8	Fats.....	12	..
Carbon.....	21	..	Calcium phosphate.....	5	13
Calcium... ..	2	..	Calcium carbonate.....	1	..
Phosphorus.....	1	12	Calcium fluoride.....	..	3
Sodium, etc.....	..	12	Sodium sulphate, etc.....	..	9
	<hr/>			<hr/>	
	154	..		154	..

STRUCTURAL COMPOSITION of THE BODY.

The Study of the Structure of the body reveals that it is composed of dissimilar parts, *e. g.*, bones, muscles, nerves, lungs, etc.; while these, again, by closer examination, can be resolved into elementary structures, the *tissues*, *e. g.*, connective tissue, muscular, nervous, epithelial tissue, etc.

Microscopical examination of the tissues shows that they are composed of fundamental structural elements, termed *cells*.

Cells are living physiological units; the simplest structural forms capable of manifesting the phenomena of life.

Cells vary in their anatomical constitution in the different structures of the body, and may be classed in three groups, viz.: 1. Cells possessing a distinct *cell wall*, *cell substance* and a *nucleus*. 2. Cells possessing a *cell*

substance and a *nucleus*. 3. Cells possessing the *cell substance* only. They vary in size, from the $\frac{1}{3000}$ to the $\frac{1}{300}$ of an inch in diameter; when young and free to move in a fluid medium they assume the spherical form; but when subjected to pressure, may become flattened, cylindrical, fusiform or stellate.

Structure of Cells. The cell wall is not an essential structure, as many cells are entirely devoid of it. It is a thin, structureless, transparent membrane, permeable to fluids.

The Cell Substance in young cells is a soft, viscid, albuminous matter, unstable, insoluble in water, and known as *protoplasm*, *bioplasm*, *sarcode*, etc.; in older cells the original cell substance undergoes various transformations, and is partly replaced by fat globules, pigment and crystals.

The Nucleus is a small vesicular body in the interior of the cell substance, and frequently contains smaller bodies, the *nucleoli*.

MANIFESTATIONS OF CELL LIFE.

Growth. Cells when newly formed are exceedingly small, but as they approach maturity they increase in size, by the capability which the cells possess of selecting and appropriating new material as food, vitalizing and organizing it. The extent of cell growth varies in different tissues; in some the cells remain exceedingly small, in others they attain considerable size. In many instances the cell substance undergoes transformation into new compounds destined for some ulterior purpose.

Reproduction. Like all organic structures cells have a limited period of life; their continual decay and death necessitates a capability of reproduction. Cells reproduce themselves in the higher animals mainly by *fission*. This is seen in the white blood corpuscles of the young embryos of animals; the corpuscle here consists of a cell substance and nucleus. When division of the cell is about to take place, the nucleus elongates, the cell substance assumes the oval form, a constriction occurs, which gradually deepens, until the original cell is completely divided and two new cells are formed, each of which soon grows to the size of the parent cell.

In cells provided with a cell membrane the process is somewhat different. In the ova of the inferior animals, after fertilization has taken place, a furrow appears on the opposite sides of the cell substance, which deepens until the cell is divided into two equal halves, each containing a nucleus; this process is again repeated until there are four cells, then eight, and so on until the entire cell substance is divided into a mulberry mass of cells,

completely occupying the interior of the cell membrane. The whole process of segmentation takes place with great rapidity, occupying not more than a few minutes, in all probability.

Motion. Spontaneous movement has been observed in many of the cells of the body. It may be studied, for example, in the movements of the spermatozooids, the waving of the cilia covering the cells of the bronchial mucous membrane, the white corpuscles of the blood, etc.

By a combination and transformation of these original structural elements, and material derived from them, all the tissues are formed which enter into the structure of the human body.

CLASSIFICATION OF TISSUES.

I. **Homogeneous Substance**, a more or less solid, albuminous structure, filling the spaces between the cells and fibres of various tissues, *e. g.*, cartilage, bone, dentine, etc.

II. **Limiting Membrane**, a thin, homogeneous membrane, structureless, composed of coagulated albumen, and often not more than the $\frac{1}{1000}$ of an inch in thickness, found lining the blood vessels and lymphatics, forming the basement membrane of the skin and mucous membranes, the posterior layer of the cornea, the capsule of the crystalline lens, etc.

III. **Simple fibrous or filamentous tissue**—the elements of which are real or apparent filaments.

(a) *Connective or areolar; white fibrous tissue*; constituting tendons, ligaments, aponeuroses, periosteum, dura mater, synovial membranes, vascular tunics, etc.

(b) *Yellow elastic tissue*, found in the middle coats of arteries, veins, lymphatics, ligamentum nuchæ, vocal cords, ligamenta subflava, etc.

IV. **Compound membranes** (membrano-cellular or fibro-cellular tissues), cells aggregated into laminæ.

(a) *Epidermic tissue*; (b) *epithelial tissue*; (c) *glandular tissue*; (d) *cornea*.

V. **Cells containing coloring matter**, or pigment cells, *e. g.*, skin, choroid membrane, etc.

VI. **Cells coalesced or consolidated by internal deposits**, *e. g.*, hair, nails, bone, teeth, etc.

VII. **Cells imbedded in an intercellular substance**, *e. g.*, cartilage, crystalline lens, etc.

VIII. **Cells aggregated in clusters**, forming tissues more or less solid, *e. g.*, adipose tissue, lymphatic glands.

IX. **Cells** imbedded in a matrix of capillaries, *e. g.*, gray or vesicular nervous matter.

X. **Cells** whose coalesced cavities form tubes containing liquids or secondary solid deposits, *e. g.*, vascular tissue, dentine.

XI. **Cells** free, isolated, or floating—fluid tissue—*e. g.*, red and white blood corpuscles, lymph and chyle corpuscles.

FOOD.

A Food may be defined to be any substance capable of playing a part in the nutrition of the body.

Food is required for the repair of the waste of the tissues consequent on their functional activity, for the generation of heat and the evolution of force.

Hunger and Thirst are sensations which indicate the necessity for taking food; they arise in the tissues at large, and are referred to the stomach and fauces, respectively, through the sympathetic nervous system.

Inanition or Starvation results from an insufficiency or absence of food, the physiological effects of which are hunger, intense thirst, intestinal uneasiness, weakness and emaciation; the quantity of carbonic acid exhaled diminishes and the urine is lessened in amount; the volume of the blood diminishes; a fetid odor is exhaled from the body; vertigo, stupor followed by delirium, and at times convulsions, result from a disturbance of the nerve centres; a marked fall of the bodily temperature occurs, from a diminished activity of the nutritive process. Death usually takes place, from exhaustion.

During starvation the loss of different tissues, before death occurs, averages $\frac{4}{10}$, or 40 per cent. of their weight.

Those tissues which lose *more than 40 per cent.* are fat, 93.3; blood, 75; spleen, 71.4; pancreas, 64.1; liver, 52; heart, 44.8; intestines, 42.4; muscles, 42.3. Those which lose *less than 40 per cent.* are the muscular coat of the stomach, 39.7; pharynx and œsophagus, 34.2; skin, 33.3; kidneys, 31.9; respiratory apparatus, 22.2; bones, 16.7; eyes, 10; nervous system, 1.9.

The *Fat* entirely disappears, with the exception of a small quantity which remains in the posterior portion of the orbits and around the kidneys. The *Blood* diminishes in volume and loses its nutritive properties. The *Muscles* undergo a marked diminution in volume and become soft and flabby. The *Nervous system* is last to suffer, not more than two per cent. disappearing before death occurs.

The *appearances* presented by the body after death from starvation are those of anæmia and great emaciation; almost total absence of fat; bloodlessness; a diminution in the volume of the organs; an empty condition of the stomach and bowels, the coats of which are thin and transparent. There is a marked disposition of the body to undergo decomposition, giving rise to a very fetid odor.

The *duration of life* after a complete deprivation of food varies from eight to thirteen days, though life can be maintained much longer if a quantity of water be obtained. The water is more essential under these circumstances than the solid matters, which can be supplied by the organism itself.

The different *alimentary principles* which are appropriated by the system are combined in different proportions in the various articles of food, and are separated from the innutritious substances during the process of digestion. They belong to the organic and inorganic worlds, and may be classified, according to their chemical composition, as follows:—

CLASSIFICATION OF ALIMENTARY PRINCIPLES.

1. Albuminous group—nitrogenized, C. O. H. N. S. P.

PRINCIPLE.	WHERE FOUND.
<i>Myosin, syntonin</i>	Flesh of animals.
<i>Vitellin, albumen</i>	Yolk of egg, white of egg.
<i>Fibrin, globulin</i>	Blood contained in meat.
<i>Casein</i>	Milk, cheese.
<i>Gluten</i>	Grain of wheat and other cereals.
<i>Vegetable albumen</i>	Soft growing vegetables.
<i>Legumin</i>	Peas, beans, lentils, etc.
<i>Gelatin</i>	Bones.

2. Saccharine group—non-nitrogenized, C. O. H.

<i>Cane sugar, beet root sugar</i>	Sugar cane, beets, etc.
<i>Glucose, grape sugar</i>	Fruits.
<i>Inosite, liver sugar, glycogen</i>	Muscles, liver, etc.
<i>Lactose or milk sugar</i>	Milk.
<i>Starch</i>	Cereals, tuberous roots and leguminous plants.

3. Oleaginous group—non-nitrogenized, C. O. H.

<i>Animal fats and oils</i>	} Found in the adipose tissue of animals, seeds, grains, nuts, fruits, and other vegetable tissues.
<i>Stearin, olein</i>	
<i>Palmitin, fatty acids</i>	

4. **Inorganic group.** Water, sodium and potassium chlorides, sodium, calcium, magnesium and potassium phosphates, calcium carbonate and iron.

5. **Vegetable acid group.** Malic, citric, tartaric and other acids, found principally in fruits.

6. **Accessory foods.** Tea, coffee, alcohol, cocoa, etc.

The *Albuminous principles* enter largely into the composition of the body, and constitute the organic bases of the different tissues; they are mainly required for the growth and repair of the tissues. There is good reason to believe that the albuminous principles are decomposed in the body into fat and urea, and the former when oxidized gives rise to the evolution of heat and force, while the latter is eliminated by the kidneys. Muscular work, however, does not result from a destruction of the albuminous compounds. The oxidation of the carbonaceous compounds, sugars and oils, furnishing the force which is transformed by the muscular system into motor power. When employed exclusively as food for any length of time, the albuminous substances are incapable of supporting life.

The *Saccharine principles* are important to the process of nutrition, but the changes which they undergo are not fully understood; they form but a small proportion of the animal tissues, and by oxidation generate heat and force. *Starch* undergoes conversion into dextrin and grape sugar.

The *Oleaginous principles* form a large part of the tissues of the body. They are introduced into the system as food, and are formed also from a transformation of albuminous matter during the nutritive process; they enter into the composition of nervous and muscular tissue, and are stored up as adipose tissue in the visceral cavities and subcutaneous connective tissue, thus giving roundness to the form and preventing, to some extent, the radiation of heat. While they aid in the reconstruction of tissue, they mainly undergo oxidation, giving rise to the production of heat and the evolution of muscular and nervous force.

The *Inorganic principles* constitute an essential part of all animal tissues, and are introduced with the food.

Water is present in all fluids and solids of the body, holding their ingredients in solution, promoting the absorption of new material into the blood and tissues, and the removal of waste ingredients.

Sodium chloride is an essential constituent of all tissues, regulating the passage of fluids through animal membranes (endosmosis and exosmosis).

Calcium phosphate gives solidity to bones and teeth, constituting more than one-half their substance.

Iron is a constituent of the coloring matter of the blood.

The *Vegetable acids* are important to nutrition, and tend to prevent the scorbutic diathesis.

The *Accessory foods* also influence the process of nutrition. *Tea* excites the respiratory function, increasing the elimination of carbonic acid. *Coffee* is a stimulant to the nervous system; increases the force of the heart's action, increases the arterial tension and retards waste.

Alcohol, when introduced into the system in *small quantities*, undergoes oxidation and contributes to the production of force, and is thus far a food. It excites the gastric glands to increased secretion, improves the digestion, accelerates the action of the heart and stimulates the activities of the nervous centres. In zymotic diseases, and all cases of depression of the vital powers, it is most useful as a restorative agent. When taken in *excessive quantities*, it is eliminated by the lungs and kidneys. The metamorphosis of the tissues is retarded, the elimination of urea and carbonic acid is lessened, the temperature lowered, the muscular powers impaired and the resistance to depressing external influences diminished. When taken through a long period of time, alcohol impairs digestion, produces gastric catarrh, disorders the secreting power of the hepatic cells. It also diminishes the muscular power and destroys the structure and composition of the cells of the brain and spinal cord. The connective tissue of the body increases in amount, and subsequently contracting, gives rise to sclerosis.

A proper combination of different alimentary principles is essential for healthy nutrition; no one class being capable of maintaining life for any definite length of time.

The *Albuminous food* in excess promotes the arthritic diathesis, manifesting itself as gout, gravel, etc.

The *Oleaginous food* in excess gives rise to the bilious diathesis, while a deficiency of it promotes the scrofulous.

The *Farinaceous food*, when long continued in excess, favors the rheumatic diathesis by the development of lactic acid.

The Alimentary Principles are not introduced into the body as such, but are combined in proper proportions to form compound substances, termed *foods*, *e. g.*, bread, milk, eggs, meat, etc., the nutritive value of each depending upon the extent to which these principles exist.

PERCENTAGE COMPOSITION OF DIFFERENT FOODS.

	WATER.	ALBUMEN.	STARCH.	SUGAR.	FATS.	SALTS.
Bread.....	37	8.1	47.4	3.6	1.6	2.3
Milk.....	86	4.1	...	5.2	3.9	0.8
Eggs.....	74	14.0	10.5	1.5
Meat.....	54	27.6	15.45	2.95
Potatoes.....	75	2.1	18.8	3.2	0.2	0.7
Corn.....	14	11.1	64.7	0.4	8.1	1.7
Oatmeal.....	15	12.6	58.4	5.4	5.6	3
Turnips.....	91	1.2	5.1	2.1	...	6
Carrots.....	83	1.3	8.4	6.1	0.2	1.0
Rice.....	13	6.3	79.1	0.4	0.7	0.5

The Amount of food required in 24 hours has been estimated at about $5\frac{1}{2}$ pounds, comprising meat, 16 oz.; bread, 19 oz.; fat, $3\frac{1}{2}$ oz.; water, 52 fluid oz.

In the excreta of the body the normal proportion of nitrogen to carbon is 1 to 15. To maintain this relation, a proper proportion of nitrogenized to carbonaceous food should be observed in diet. The proportion in the excreta would be—

On a meat diet.....	nitrogen 1 to carbon	3
On a bread diet.....	" 1 "	30
On a mixed diet.....	" 2 "	33
Or.....	" 1 "	16

The amount of nitrogen and carbon necessary to compensate for the loss would be contained in 16 oz. of meat and 2 lbs. of bread; as, however, usually 3 oz. of oil are consumed in 24 hours, only 19 oz. of bread are required.

COMPARISON OF INGESTA AND EGESTA IN 24 HOURS.

FOOD, DRINK, AIR.	OZ.	EXCRETIONS.	OZ.
Albumen.....	4.23	Breath { carbonic acid }	43.40
Starch.....	11.63	{ watery vapor }	
Fats.....	3.17	Perspiration { carbonic acid }	23.62
Salts.....	1.13	{ watery vapor }	
Water (6 pints).....	93.00	Urine.....	66.31
Oxygen.....	26.24	Solid excreta.....	6.07
Total.....	139.40	Total.....	139.40

DIGESTION.

Digestion is a physical and chemical process, by which the food is changed, by the action of solvent fluids, into a form capable of being absorbed into the blood.

The **Digestive Apparatus** consists of the alimentary canal and its appendages, viz.: teeth, salivary, gastric and intestinal glands, liver and pancreas.

Digestion may be divided into *seven stages*: prehension, mastication, insalivation, deglutition, gastric and intestinal digestion and defecation.

Prehension, the act of conveying food into the mouth, is accomplished by the hands, lips and teeth.

Mastication is the trituration of the food, and is accomplished by the teeth and lower jaw, under the influence of muscular contraction. When thoroughly divided, the food presents a greater surface for the solvent action of the digestive fluids, thus aiding the general process of digestion.

The **Teeth** are thirty-two in number, sixteen in each jaw, and divided into four incisors or cutting teeth, two canines, four bicuspid, and six molars or grinding teeth; each tooth consists of a crown covered by enamel, a neck, and a root surrounded by the crista petrosa, and imbedded in the alveolar process; a section through a tooth shows that its substance is made up of *dentine*, in the centre of which is the pulp cavity, containing blood vessels and nerves.

The *lower jaw* is capable of making a downward and an upward, a lateral and an antero-posterior movement, dependent upon the construction of the temporo-maxillary articulation.

The jaw is *depressed* by the contraction of the *digastric*, *genio-hyoid*, *mylo-hyoid* and *platysma myoides* muscles; *elevated* by the *temporal*, *masseter* and *internal pterygoid* muscles; moved *laterally* by the alternate contraction of the *external pterygoid* muscles; moved *anteriorly* by the *pterygoid* and *posteriorly* by the united actions of the *genio-hyoid*, *mylo-hyoid* and posterior fibres of the *temporal* muscle.

The food is kept between the teeth by the *intrinsic* and *extrinsic* muscles of the tongue from within, and the *orbicularis oris* and *buccinator* muscles from without.

The **Movements of Mastication** are called forth by *reflex action* through the medulla oblongata, induced by the presence of food in the mouth.

NERVOUS CIRCLE OF MASTICATION.

AFFERENT OR EXCITOR NERVES.

1. Lingual branch of 5th pair.
2. Glosso-pharyngeal.


Efferent or Motor.

1. 3d branch of 5th pair.
2. Hypo-glossal.
3. Facial.

Insalivation is the incorporation of the food with the saliva secreted by the *parotid*, *sub-maxillary* and *sub-lingual glands*; the *parotid* saliva, thin and watery, is poured into the mouth through Steno's duct; the *sub-maxillary* and *sub-lingual* salivas, thick and viscid, are poured into the mouth through Wharton's and Bartholini's ducts.

In a living serous gland, *e. g.*, parotid, during rest, the secretory cells lining the acini of the gland are seen to be filled with fine granules, which are often so abundant as to obscure the nucleus and enlarge the cells until

FIG. 1.



Cells of the alveoli of a serous or watery salivary gland. A. After rest. B. After a short period of activity. C. After prolonged period of activity.—*From Yeo's Text-Book of Physiology.*

the lumen of the acinus is almost obliterated (Fig. 1). When the gland begins to secrete the saliva, the granules disappear from the outer boundary of the cells, which then become clear and distinct. At the end of the secretory activity, the cells have become free of granules, have become smaller and more distinct in outline. It would seem that the granular matter is formed in the cells during the rest, and discharged into the ducts during the activity of the gland.

In the mucous glands, *e. g.*, sub-maxillary and sub-lingual, the changes that occur in the cells are somewhat different (Fig. 2). During the intervals of digestion, the cells lining the gland are large, clear and highly refractive, and contain a large quantity of mucigen. After secretion has taken place, the cells exhibit a marked change. The mucigen cells have disappeared, and in their place are cells which are small, dark and composed of protoplasm. It would appear that the cells, during rest, elaborate the


mucigen which is discharged into the tubules during secretory activity, to become part of the secretion.

Saliva is an opalescent, slightly viscid, alkaline fluid, having a specific gravity of 1.005. Microscopical examination reveals the presence of salivary corpuscles and epithelial cells. Chemically it is composed of water, proteid matter, a ferment (*ptyalin*) and inorganic salts. The amount secreted in 24 hours is about $2\frac{1}{2}$ lbs. Its function is twofold:—

1. *Physical*.—Softens and moistens the food, glues it together, and facilitates swallowing.


2. *Chemical*.—Converts *starch* into *grape sugar*. This action is due to

FIG. 2.



Section of a "mucous" gland. A. In a state of rest. B. After it has been or some time actively secreting.—After Lavdowsky.

the presence of the organic ferment, ptyalin. The change consists in the assumption of a molecule of water.



NERVOUS CIRCLE OF INSALIVATION.

AFFERENT OR EXCITOR NERVES.

1. Lingual branch of 5th pair.
2. Glosso-pharyngeal.

EFFERENT OR MOTOR NERVES.

1. Auriculo-temporal branch of 5th pair, for parotid gland.
2. Chorda tympani, for sub-maxillary and sub-lingual glands.

The centres regulating the secretion are two, viz.: The medulla oblongata and the sub-maxillary ganglion of the sympathetic; the latter acting antagonistically to the former. Impressions excited by the food in the

mouth reach the medulla oblongata through the afferent nerves; motor impulses are there generated which pass outward through the efferent nerves.

Stimulation of the auriculo-temporal branch increases the flow of saliva from the parotid gland; *division* arrests it.

Stimulation of the chorda tympani is followed by a dilation of the blood vessels of the sub-maxillary gland, increased flow of blood (thus acting as a vaso-dilator nerve) and an abundant discharge of a thin saliva; *division* of the nerve arrests the secretion.

Stimulation of the cervical sympathetic is followed by a contraction of the blood vessels, diminishing the flow of blood (thus acting as a vaso-constrictor nerve) and a diminution of the secretion, which now becomes thick and viscid; *division* of the sympathetic does not, however, completely dilate the vessels. There is evidence of the existence of a local vaso-motor mechanism, which is *inhibited* by the chorda tympani; *exalted* by the sympathetic.

Deglutition is the act of transferring food from the mouth into the stomach, and may be divided into *three* stages:—

1. The passage of the bolus from the mouth into the pharynx.
2. From the pharynx into the cesophagus.
3. From the cesophagus into the stomach.

In the *1st stage*, which is entirely voluntary, the mouth is closed and respiration momentarily suspended; the tongue, placed against the roof of the mouth, arches upward and backward, and forces the bolus into the fauces.

In the *2d stage*, which is entirely reflex, the palate is made tense and directed upward and backward by the levatores-palati and tensores-palati muscles; the bolus is grasped by the superior constrictor muscle of the pharynx and rapidly forced into the cesophagus.

The food is prevented from entering the *posterior nares* by the uvula and the closure of the posterior half-arches (the palato-pharyngei muscles); from entering the *larynx* by its ascent under the base of the tongue and the action of the epiglottis.

In the *3d stage*, the longitudinal and circular muscular fibres, contracting from above downward, *strip* the bolus into the stomach. [For nervous mechanism of Deglutition, see Medulla Oblongata.]

Gastric Digestion. The stomach is a dilatation of the alimentary canal, 13 inches long, 5 inches deep, having a capacity of about 5 pints; there can be distinguished a cardiac and pyloric orifice, a greater and lesser curvature, a greater and lesser pouch.

It possesses three coats:—

1. Serous, a reflection of the peritoneum.
2. Muscular, the fibres of which are arranged longitudinally, transversely and obliquely.
3. Mucous, thrown into folds, forming the rugæ.

Imbedded in the mucous coat are immense numbers of *mucous* and *true gastric glands*. In the pyloric end of the stomach are found the

FIG. 3.




Diagram showing the relation of the ultimate twigs of the blood vessels, V and A, and of the absorbent radicles to the glands of the stomach and the different kinds of epithelium, viz., above cylindrical cells; small, pale cells in the lumen, outside which are the dark ovoid cells.—From *Yeo's Text-Book of Physiology*.

mucous glands, which are lined with columnar epithelium throughout their extent. In the cardiac end are found the true *gastric glands* (Fig. 3), the ducts of which are also lined with columnar cells, while the secretory parts are lined with two distinct varieties of cells. One variety consists of small, spheroidal, granular cells, which border the lumen of the gland, and are known as the *chief cells*; the other variety consists of large, oval, well-defined granular cells, much less

abundant, and are situated between the basement membrane of the gland and the chief cells. From their position they have been termed *parietal cells*. During the intervals of digestion the chief cells are pale, and the hyaline substance of which they are composed is finely granular. During the stage of active secretion the cells become swollen and turbid, and are then said to be rich in pepsin. Toward the end of digestion the granules disappear, the cells become pale and return to their former size.

During the intervals of digestion, the mucous membrane of the stomach is pale and covered with a layer of mucus. Upon the introduction of food, the blood vessels dilate and become filled with blood, and the mucous membrane becomes red. At the same time small drops of a fluid, the gastric juice, begin to exude upon its surface, which gradually run together and trickle down the sides of the stomach.

The Gastric Juice is a secretion of the true peptic cells, and when obtained from the stomach through a fistulous opening, is a clear, straw-colored fluid, decidedly acid, with a specific gravity of 1.005 to 1.010.

COMPOSITION OF GASTRIC JUICE.

Water.....	975.00
Pepsin.....	15.00
Hydrochloric acid.....	4.78
Inorganic salts.....	5.22
	<hr/>
	1000.00

The water forms the largest part of the fluid, and holds in solution the other ingredients. It results from a transudation from the blood vessels under the increased blood supply. Of the inorganic salts the chlorides of sodium and potassium are the most abundant.

Pepsin is the organic nitrogenized ferment of the gastric juice, and is formed, during the intervals of digestion, by the peptic cells. In the presence of a small per cent. of an acid, it acquires the property of converting the albumen of the food into albuminose or peptones.

Hydrochloric acid is present in small quantity, and gives the juice its acidity. In all probability, its production is due to the activity of the parietal cells.

These two characteristic ingredients of the gastric juice exist in a state of combination as hydrochloro-peptic acid, and the presence of both is absolutely essential for the complete digestion of the food. When the food enters the stomach, it is subjected to the peristaltic action of the muscular coat, and thoroughly incorporated with the gastric juice. This

fluid possesses the property of attacking the connective tissues of the food, disintegrating it and dissolving out the alimentary principles.

The Principal Action of the gastric juice, however, is to transform the different albuminous principles of the food into *peptones* or *albuminose*, the different stages of which are due to the acid and pepsin respectively. When freed from its combination, the hydrochloric acid converts the albumen into *acid albumen* or *para-peptone*; while this intermediate product is being formed, the pepsin converts it at once into *peptone*. In order that the digestion of albumen may be complete, it is necessary that both the acid and pepsin be present in proper quantity. Before digestion, the albuminous principles are insoluble in water and incapable of being absorbed. After digestion, they become soluble and are readily absorbed. Peptones differ from the albumins in being—

1. *Diffusible*, passing rapidly through the mucous membrane and walls of the blood vessels.
2. *Non-coagulable* by heat, nitric or acetic acids; but are readily precipitated by tannic acid,
3. *Soluble* in water and saline solutions.
4. *Assimilable* by the blood; when injected into it, they do not reappear in the urine.

Gastric juice exerts no influence either upon grape sugar, cane sugar, starch or fat.

Gastric Digestion occupies on the average from 3 to 5 hours, but varies in duration according to the nature and quantity of the food, exercise, temperature, etc.

The *Amount* of gastric juice secreted in 24 hours varies, under normal conditions, from 8 to 14 pounds.

Movements of the Stomach. As soon as digestion commences, the cardiac and pyloric orifices are closed; the walls of the stomach contract upon the food, and a peristaltic action begins, which carries the food along the greater and lesser curvatures, and thoroughly incorporates it with the gastric juice. As soon as any portion of the food is digested, it passes through the pylorus into the intestine.

Vomiting. The act of vomiting is usually preceded by nausea and a discharge of saliva into the mouth. This is then swallowed, and carries into the stomach a quantity of air which facilitates the ejection of the contents of the stomach by aiding the relaxation of the cardiac sphincter. A deep inspiration is then taken, during which the lower ribs are drawn in and the diaphragm descends and remains contracted. At the same


time the glottis is closed. A sudden expiratory effort is now made, and the cardiac orifice being open, the abdominal muscles contracting, press upon the stomach and forcibly eject its contents into the mouth.

Intestinal Digestion. The intestine is about 20 feet long, $1\frac{1}{2}$ inches in diameter, and possesses three coats:—

1. Serous (peritoneal).
2. Muscular, the fibres of which are arranged longitudinally and transversely.
3. Mucous, thrown into folds, forming the *valvulae conniventes*.

This stage of digestion is probably the most complex and important; here the different alimentary principles are further elaborated and prepared

FIG. 4.



One saccule of the pancreas of the rabbit in different states of activity. *A.* After a period of rest, in which case the outlines of the cells are indistinct, and the inner zone, *i. e.*, the part of the cells (*a*) next the lumen (*c*), is broad and filled with fine granules. *B.* After the gland has poured out its secretion, when the cell outlines (*d*) are clearer, the granular zone (*a*) is smaller, and the clear outer zone is wider.—*From Yeo's Text-Book of Physiology, after Kühne and Lea.*

for absorption into the blood by being acted upon by the *intestinal juice*, *pancreatic juice* and *bile*.

Throughout the mucous coat are imbedded the intestinal follicles, the glands of Brunner and Lieberkühn. They secrete the true *intestinal juice*, which is an alkaline, viscid fluid, composed of water, organic matter and salts. Its *function* is to convert *starch* into glucose, and assist in the digestion of the albuminoids.

The Pancreatic Juice is secreted by the pancreas, a flattened gland about six inches long, running transversely across the posterior wall of the abdomen, behind the stomach; its duct opens into the duodenum.

The pancreas is similar in structure to the salivary glands, consisting of a system of ducts terminating in acini. The acini are tubular or flask-shaped, and consist of a basement membrane lined by a layer of cylindrical, conical cells, which encroach upon the lumen of the acini. The cells exhibit a difference in their structure (Fig. 4), and may be said to consist of two zones, viz., an *outer parietal zone*, which is transparent and apparently homogeneous, staining readily with carmine; an *inner zone*, which borders the lumen, and is distinctly granular and stains but slightly with carmine. These cells undergo changes similar to those exhibited by the cells of the salivary glands during and after active secretion. As soon as the secretory activity of the pancreas is established, the granules disappear, and the inner granular layer becomes reduced to a very narrow border while the outer zone increases in size and occupies nearly the entire cell. During the intervals of secretion, however, the granular layer reappears and increases in size until the outer zone is reduced to a minimum. It would seem that the granular matter is formed by the nutritive processes occurring in the gland during rest, and is discharged during secretory activity into the ducts and takes part in the formation of the pancreatic secretion.

The pancreatic juice is transparent, colorless, strongly alkaline and viscid, and has a specific gravity of 1.040. It is one of the most important of the digestive fluids, as it exerts a transforming influence upon all the classes of alimentary principles, and has been shown to contain at least three distinct ferments. It has the following composition :—

COMPOSITION OF PANCREATIC JUICE.

Water	900.76
Albuminoid substances.....	90.44
Inorganic salts.....	8.80
	<hr/>
	1000.00

The pancreatic juice is characterized by its action: 1st. Upon *starch*. When starch is subjected to the action of the juice, it is at once transformed into glucose; the change takes place more rapidly than when saliva is added. This action is caused by the presence of a special ferment, *amyl-opsin*. 2d. Upon *albumen*. The albuminous bodies are changed by the juice into, first, an alkali albumen, and then into peptone. The albumen does not swell up, as is the case in gastric digestion, but is gradually corroded and dissolved. This change is due to the presence of the ferment, *trypsin*. Long-continued action of trypsin converts the peptones into two crystalline bodies, leucine and tyrosin. 3d. Upon *fats*. The most striking action of the pancreatic juice is the *emulsification* of the fats or their sub-

division into minute particles of microscopic size. This change takes place rapidly and depends upon the alkalinity of the fluid and the quantity of albumen present, combined with the intestinal movements. The *neutral fats* are also decomposed into their corresponding *fatty acids* and *glycerine*; the acids thus set free unite with the alkaline bases present in the intestine and form soaps. This decomposition of the neutral fats is caused by the ferment, *steapsin*. 4th. Upon *cane sugar* the juice also exerts a special influence, converting it readily into glucose.

The total quantity of this fluid secreted in twenty-four hours has not been accurately determined; varies from one to two pounds. It is poured out most abundantly an hour after meals.

The **Bile** has an important influence in the elaboration of the food and its preparation for absorption. It is a golden-brown, viscid fluid, having a neutral or slightly alkaline reaction and a specific gravity of 1.020.

COMPOSITION OF BILE.

Water.....	859.2
Sodium glycocholate }	91.4
Sodium taurocholate }	
Fat	9.2
Cholesterine.....	2.6
Mucus and coloring matter.....	29.8
Salts	7.8
	<hr/>
	1000.00

The *Biliary salts*, sodium glycocholate and taurocholate, are characteristic ingredients, and are formed in the liver by the process of secretion, from materials furnished by the blood. It is probable that they are derived from the nitrogenized compounds, though the stages in the process are unknown. They are reabsorbed from the small intestine to play some ulterior part in nutrition.

Cholesterine is a product of waste taken up by the blood from the nervous tissues and excreted by the liver. It crystallizes in the form of rhombic plates, which are quite transparent. When retained within the blood, it gives rise to the condition of *cholesteræmia*, attended with severe nervous symptoms. It is given off in the fæces under the form of stercorine.

The *Coloring matters* which give the tints to the bile are *biliverdin* and *bilirubin*, and are probably derived from the coloring matter of the blood. Their presence in any fluid can be recognized by adding to it nitric acid containing nitrous acid, when a play of colors is observed, beginning with green, blue, violet, red and yellow.

The Bile is both a secretion and an excretion; it is constantly being formed and discharged by the hepatic ducts into the gall bladder, in which it is stored up, during the intervals of digestion. As soon as food enters the intestines, it is poured out abundantly, by the contraction of the walls of the gall bladder.

The *Amount* secreted in 24 hours is about $2\frac{1}{2}$ pounds.

Functions of the Bile. (1) It assists in the *emulsification* of the fats and promotes their absorption. (2) It tends to prevent putrefactive changes in the food. (3) Stimulates the secretions of the intestinal glands, and excites the normal peristaltic movements of the bowels.

The digested food, the *chyme*, is a grayish, pultaceous mass, but as it passes through the intestines it becomes yellow, from admixture with the bile. It is propelled onward by vermicular motion; by the contraction of the circular and longitudinal muscular fibres.

As the digested food passes through the intestines, the nutritious matters are absorbed into the blood, and the residue enters the large intestine.

The **Fæces** consist chiefly of indigestible matters, *excretin*, *stercorin* and salts; varying in amount from 4 to 7 oz. in 24 hours.

Defecation is the voluntary act of extruding the fæces from the body; accomplished by a relaxation of the sphincter muscle, the contraction of the walls of the rectum, assisted by the abdominal muscles.

The **Gases** contained in the stomach and small intestine are oxygen, nitrogen, hydrogen and carbonic acid. In the large intestine, carbonic acid, sulphuretted and carburetted hydrogen. They are introduced with the food, and also developed by chemical changes in the alimentary canal. They distend the intestines, aid capillary circulation, and tend to prevent pressure.

ABSORPTION.

Absorption has for its object the introduction of new materials into the blood, and takes place mainly from the alimentary tract; but also to some extent from the skin, respiratory surface and closed cavities of the body.

The *Agents of Absorption* are the veins and lymphatics.

As a result of the process of digestion, the different alimentary substances are converted into forms which are capable of being absorbed into the blood, *e.g.*, *albuminose*, *glucose* and *fatty emulsion*, water and inorganic matters undergoing no change, being already in a condition to be absorbed and to play a part in the nutritive process.

The blood vessels which are most active as absorbents, are the *gastric, superior and inferior mesenteric veins*. They arise in the coats of the alimentary canal, and as they converge, unite with the splenic vein to form the *portal vein* which enters the liver (Fig. 5).

As the digested mass of food, the *chyme*, passes through the alimentary canal, a large portion of it disappears; the veins absorb *water*,

FIG. 5.




Diagram of the portal vein (*pv*) arising in the alimentary tract and spleen (*s*), and carrying the blood from these organs to the liver.—From Yeo's *Text-Book of Physiology*.

albuminose, glucose, and inorganic salts, and convey them directly into the liver; the blood of the portal vein being especially rich in these substances.

At times, after the ingestion of large quantities of oleaginous food, the blood vessels take up, in addition, a certain quantity of fatty matter; but this is not usually the case; the *fats* being absorbed by special vessels, the *lymphatics or lacteals*.

General Anatomy of the Lymphatic System. The lymphatics constitute a system of minute, delicate, transparent vessels, which, having their origin at the periphery of the body, pass forward toward the centre and empty into the veins at the base of the neck, by means of the thoracic duct. In their course they pass through small ovoid bodies, the *lymphatic glands*.

Origin of the Lymphatics. The lymphatic vessels arise in several distinct ways: 1. In *lymph spaces* or *juice canals*. Throughout the connective tissues of the body are found numbers of small, irregular, stellate spaces, which communicate freely with each other, and represent the ultimate radicles of the lymphatic vessels. They frequently contain lymph corpuscles. These spaces communicate with the lymph trunks, through the medium of the plexuses of lymph capillaries, which are much larger than capillary blood vessels, and are lined with endothelial cells with sinuous margins. 2. In *openings on the surface of serous membranes*. The surfaces of the serous membranes are covered with a layer of endothelial cells. At intervals between these cells are found openings—the *stomata*. These stomata communicate directly, through short canals, with the lymph capillaries. The serous cavities may be thus regarded as large lymph spaces. 3. In *perivascular lymph spaces*. In the brain and spinal cord the capillary blood vessels are surrounded by a sheath, formed of endothelial cells, and which contains lymph. This space is in free communication with the lymphatics. By this means the blood vessel is bathed by the lymph stream.

The lymphatic vessels of the small intestine (*the lacteals*) arise within the villous processes which project from the inner surface of the intestine throughout its entire extent. The wall of the villus is formed by an elevation of the basement membrane, and covered by a layer of columnar epithelial cells. The basis of the villus consists of adenoid tissue, fine plexus of blood vessels, unstriated muscular fibres and the lacteal vessel. The adenoid tissue consists of a number of intercommunicating spaces, containing leucocytes. The lacteal vessel possesses a thin, but distinct wall, composed of endothelial plates, with here and there openings, which bring the interior of the villus into communication with the spaces of the adenoid tissue.

The *structure* of the larger vessels resembles that of the veins, consisting of three coats—

1. *External*, composed of fibrous tissue and muscular fibres, arranged longitudinally.
2. *Middle*, consisting of white fibrous and yellow elastic tissue, non striated muscular fibres, arranged transversely.
3. *Internal*, composed of an elastic membrane, lined by endothelial cells.

Throughout their course are found numerous *semilunar valves*, looking toward the larger vessels, formed by a folding of the inner coat and strengthened by connective tissue.

Lymphatic Glands consist of an external fibrous covering, from the inner surface of which partitions of fibrous tissue, *the trabeculae*, pass into the substance of the gland, forming a *stroma* or network, in the meshes of which are found the true *lymph corpuscles*.

The lymphatics which enter the gland are called the *afferent* vessels; those which leave it, the *efferent* vessels.

The **Thoracic Duct** is the general trunk of the lymphatic system, into which the vessels of the lower extremities, of the abdominal organs, of the left side of the head and left arm empty their contents. It is about twenty inches in length, arises in the abdomen, opposite the third lumbar vertebra, by a dilatation, the *receptaculum chyli*; ascends along the vertebral column to the seventh cervical vertebra, and terminates in the venous system at the junction of the internal jugular and subclavian veins on the left side. The lymphatics of the right side of the head, of the right arm and the right side of the thorax, terminate in the *right thoracic duct*, about one inch in length, which joins the venous system at the junction of the internal jugular and subclavian on the right side.

Lymph is a clear, transparent fluid, slightly alkaline, having a saline taste and a specific gravity of 1.022. It is found in the lymphatic vessels throughout the body.

Lymph contains a number of corpuscles (the *leucocytes*) resembling the white corpuscles of the blood, which increase in number as it passes through the lymphatic glands. They are about $\frac{1}{2500}$ of an inch in diameter and somewhat granular; they are discharged into the blood, but their function is obscure. When withdrawn from the vessels, lymph undergoes spontaneous coagulation, separating into serum and clot, as in the case of the blood.

COMPOSITION OF LYMPH.

	DR. OWEN REES.
Water.....	96.536
Proteids (serum-albumen, fibrin, globulin).....	1.320
Extractives (urea, sugar, cholesterine).....	1.559
Fatty matter.....	a trace
Salts.....	0.585
	<hr/>
	100.000

Origin of Lymph. Lymph is undoubtedly a transudation from the capillary blood vessels, occurring during the process of nutrition, and is

identical, for the most part, with the liquor sanguinis, or plasma. As new material is constantly exuded, the old is absorbed by the lymphatics, and returned again to the circulation.

Excrementitious matters, as *urea*, *cholesterine*, etc., are also taken up from the tissues by the lymphatics and emptied into the blood.

FIG. 6.




Diagram showing the course of the lacteals through the mesentery, and their termination in the thoracic duct. The course of the thoracic duct and its termination at the junction of the internal jugular and subclavian veins.

The *total quantity* of lymph poured into the thoracic duct in 24 hours has been estimated at $3\frac{1}{2}$ lbs.

Chyle. As a result of the process of digestion, the oleaginous matters which have been acted upon by the *pancreatic juice* and *bile* are trans-

formed into a condition of *emulsion*, forming an opaque, milky fluid, termed *chyle*, which adheres to the folds of the mucous membrane and villi.

The Molecules of the fat are first absorbed by the epithelial cells upon the surface of the villi, through which they pass and enter the lymphatics.

Absorption by the Lacteals. The *lacteals*, or lymphatics of the small intestine, have their origin in the interior of the villi, from which they emerge and form a lymphatic plexus; the larger branches of which pass through the layers of the mesentery, and finally terminate in the thoracic duct (Fig. 6).

In the *intervals* of digestion the lacteals contain clear, transparent lymph, and are invisible on account of their small size and delicacy. But *during digestion* these vessels become filled, from absorption of the chyle, and form a visible network of white vessels ramifying through the mesentery, and converging toward the *receptaculum chyli*.

The lacteal vessels also absorb a small quantity of *water, albuminose, glucose and salts*.

COMPOSITION OF CHYLE.

Water	902.37
Albumen	35.16
Fibrin.....	3.70
Extractives.....	15.65
Fatty matters.....	36.01
Salts	7.11
	<hr/>
	1000.00

The *Products of digestion* find their way into the general circulation by two routes:—

1. *Water, albuminose, glucose and salts* are mainly absorbed by the *gastric and mesenteric veins*, carried into the liver, through the capillaries of which they pass, and enter the inferior vena cava by the hepatic veins.

2. The *Fats* are absorbed by the lacteals, emptied into the thoracic duct, and enter the blood at the junction of the internal jugular and subclavian veins.

Forces aiding the movements of Lymph and Chyle.

1. *Endosmosis.* The continued transudation of matter from the capillaries, and its absorption into the lymphatics by *endosmosis*, constitutes the main cause, the *vis-a-tergo*, of the movement of the lymph; it is so considerable as to rupture the walls of the vessels if they are ligated.

2. *Contraction* of the *non-striated muscular* fibres in the walls of the lymphatic vessels, especially when fully distended, aided by the action of the valves, promotes the onward flow of the fluids.

3. *Muscular contraction* in all parts of the body, by exerting intermittent pressure upon the lymphatic vessels, hastens the current onward; regurgitation being prevented by the closure of the valves.

4. *The inspiratory movement*, by expanding the chest, causes a dilation of the thoracic duct, and a rapid flow of lymph and chyle into it; during expiration it is compressed, and the fluids forcibly expelled into the venous system.

BLOOD.

The Blood is a nutritive fluid containing all the elements necessary for the repair of the tissues; it also contains principles of waste absorbed from the tissues, which are conveyed to the various excretory organs and by them eliminated from the body.

The *total amount* of blood in the body is estimated to be about one-eighth of the body weight; from 16 to 18 pounds in an individual of average physical development. The quantity varies during the 24 hours; the *maximum* being reached in the afternoon, the *minimum* in the early morning hours.

Blood is a heterogeneous, opaque red fluid, having an alkaline reaction, a *saline taste*, and a *specific gravity* of 1.055.

The *opacity* is due to the refraction of the rays of light by the elements of which the blood is composed. The *color* varies in hue, from a bright scarlet in the arteries to a deep purple in the veins, due to the presence of a coloring matter, *hemoglobin*, in different degrees of oxidation.

The *alkalinity* is constant, and depends upon the presence of the alkaline sodium phosphate, $\text{Na}_2 \text{H P O}_4$.

The *saline taste* is due to the amount of sodium chloride present.

The *specific gravity* ranges within the limits of health, from 1.045 to 1.075.

The *odor* of the blood is characteristic, and varies with the animal from which it is drawn, due to the presence of caproic acid.

The *temperature* of the blood ranges from 98° Fahr. at the surface to 107° Fahr. in the hepatic vein; it loses heat by radiation and evaporation as it approaches the extremities, and as it passes through the lungs.

Blood consists of two portions :—

1. The *Liquor Sanguinis* or *Plasma*, a transparent, colorless fluid, in which are floating—

2. *Red and white corpuscles*; these constituting by weight less than one-half, 40 per cent., of the entire amount of blood.

COMPOSITION OF PLASMA.		DALTON.
Water.....		902.00
Albumen.....		53.00
Paraglobulin.....		22.00
Fibrinogen.....		3.00
Fatty matters.....		2.50
Crystallizable nitrogenous matters.....		4.00
Other organic matter.....		5.00
Mineral salts.....		8.50
		<hr/> 1000.00

Water acts as a solvent for the inorganic matters and holds in suspension the corpuscular elements.

Albumen is the nutritious principle of the blood; it is absorbed by the tissues to repair their waste, and is transformed into the organic basis characteristic of each structure.

Paraglobulin or *fibrinoplastin* is a soft amorphous substance precipitated by sodium chloride in excess, or by passing a stream of carbonic acid through dilute serum.

Fibrinogen can also be obtained by *strongly diluting* the serum and passing carbonic acid through it for a long time, when it is precipitated as a viscous deposit.

Fatty matters exist in small proportion, except in pathological conditions and after the ingestion of food rich in oleaginous matters; it soon disappears, undergoing oxidation, generating heat and force, or is deposited as adipose tissue.

Sugar is represented by glucose, a product of the digestion of saccharine matter and starches in the alimentary canal; glycogenic matter is derived from the liver.

The *Saline constituents* aid the process of osmosis, give alkalinity to the blood, promote the absorption of carbonic acid from the tissues into the blood, and hold other substances in solution; the most important are the sodium and potassium chlorides, the calcium and magnesium phosphates.

Excrementitious matters are represented by carbonic acid, urea, creatin, creatinin, urates, oxalates, etc.; they are absorbed from the tissues by the blood and conveyed to the excretory organs, lungs, kidneys, etc.

Gases. Oxygen, nitrogen, and carbonic acid exist in varying proportions.

BLOOD CORPUSCLES.

The corpuscular elements of the blood occur under two distinct forms, which from their color are known as the *red* and *white* corpuscles.

The *Red Corpuscles*, as they float in a thin layer of the *Liquor Sanguinis*, are of a pale straw color; it is only when aggregated in masses that they assume the bright red color. In form they are circular and biconcave; they have an average diameter of the $\frac{1}{3200}$ of an inch.

In mammals, birds, reptiles, amphibia and fish the corpuscles vary in size and number, gradually becoming larger and less numerous as the scale of animal life is descended, *e. g.*:—

TABLE SHOWING COMPARATIVE DIAMETER OF RED CORPUSCLES.

<i>Mammals.</i>	<i>Birds.</i>	<i>Reptiles.</i>	<i>Amphibia.</i>	<i>Fish.</i>
Man, 3200.	Eagle, 1812.	Turtle, 1231.	Frog, 1108.	Perch, 2005.
Chimpanzee, 3112.	Owl, 1703.	Tortoise, 1252.	Toad, 1043.	Carp, 2141.
Ourang, 3343.	Sparrow, 2140.	Lizard, 1555.	Proteus, 400.	Pike, 2000.
Dog, 3542.	Swallow, 2133.	Viper, 1274.	Siren, 420.	Eel, 1745.
Cat, 4404.	Pigeon, 1873.		Amphiuma, 383.	
Hog, 3730.	Turkey, 3043.			
Horse, 4000.	Goose, 1808.			
Ox, 4707.	Swan, 1808.			

In man and the mammals the red corpuscles present neither a nucleus nor a cell wall, and are universally of a small size. They can be readily distinguished from the corpuscles of birds, reptiles and fish, in which they are larger, oval in shape and possess a well-defined nucleus.

The red corpuscles are exceedingly numerous, amounting to about 5,000,000 in a cubic millimetre of blood. In structure they consist of a firm, elastic, colorless framework, the *stroma*, in the meshes of which is entangled the coloring matter, the *hæmoglobin*.

CHEMICAL COMPOSITION OF RED CORPUSCLES.

Water.....	688.00
Globulin.....	282.22
Hæmoglobin.....	16.75
Fatty matter.....	2.31
Extractives.....	2.60
Mineral salts.....	8.12

1000.00

Hæmoglobin, the coloring matter of the corpuscles, is an albuminous compound, composed of C. O. H. N. S. and iron. It may exist either in an amorphous or crystalline form. When deprived of all its oxygen, except the quantity entering into its intimate composition, the hæmoglobin becomes darker in color, somewhat purple in hue, and is known as *reduced hæmoglobin*. When exposed to the action of oxygen, it again absorbs a definite amount and becomes scarlet in color, and is known as *oxy-hæmoglobin*. The amount of oxygen absorbed is 1.76 c.cm. ($\frac{7}{10}$ cubic inch) for 1 milligramme ($\frac{1}{64}$ grain) of hæmoglobin.

It is this substance which gives the color to the venous and arterial blood. As the venous blood passes through the capillaries of the lungs, the *reduced hæmoglobin* absorbs the oxygen from the pulmonary air and becomes *oxy-hæmoglobin*, scarlet in color, and the blood becomes arterial. When the arterial blood passes into the systemic capillaries, the oxygen is absorbed by the tissues, the hæmoglobin becomes reduced, purple in color, and the blood becomes venous. A dilute solution of oxy-hæmoglobin gives two absorption bands between the lines D and E of the solar spectrum. Reduced hæmoglobin gives but one absorption band, occupying the space existing between the two bands of the oxy-hæmoglobin spectrum.

The *Function* of the red corpuscles is, therefore, to absorb oxygen and carry it to the tissues; the smaller the corpuscles, and the greater the number, the greater is the quantity of oxygen absorbed; and, consequently, all the vital functions of the body become more active.

The *White Corpuscles* are far less numerous than the red, the proportion being, on an average, about 1 white to 350 or 400 red; they are globular in shape, and measure the $\frac{1}{2500}$ of an inch in diameter, and consist of a soft, granular, colorless substance, containing several nuclei.

The *white corpuscles* possess the power of spontaneous movement, alternately contracting and expanding, throwing out processes of their substance and quickly withdrawing them, thus changing their shape from moment to moment. These movements resemble those of the amœba, and for this reason are termed *amœboid*. They also possess the capability of moving from place to place. In the interior of the vessels they adhere to the inner surface, while the red corpuscles move through the centre of the stream.

The white corpuscles are identical with the leucocytes, and are found in milk, lymph, chyle and other fluids.

Origin of Corpuscles. The red corpuscles take their origin from the mesoblastic cells in the vascular area of the developing embryo.

In the adult they are produced from colorless nucleated corpuscles

resembling the white corpuscles. The spleen is the organ in which they are finally destroyed.

The white corpuscles originate from the leucocytes of the adenoid tissue, and subsequently give rise to the red corpuscles and partly to new tissues that result from inflammatory action.

COAGULATION OF THE BLOOD.

When blood is withdrawn from the body and allowed to remain at rest, it becomes somewhat thick and viscid in from three to five minutes; this viscosity gradually increases until the entire volume of blood assumes a jelly-like consistence, which occupies from five to fifteen minutes.

As soon as coagulation is completed, a second process begins, which consists in the contraction of the coagulum and the oozing of a clear, straw-colored liquid, the *serum*, which gradually increases in quantity as the *clot* diminishes in size, by contraction, until the separation is completed, which occupies from 12 to 24 hours.

The changes in the blood are as follows:—

Before coagulation.

Living blood.	{	Liq. Sanguinis	}	Consisting of	{	Water.
		or				Albumen.
		Plasma.				Fibrinogen.
		Corpuscles. Red and white.				Salts.

After coagulation.

Dead blood.	{	Crassamentum.	}	Containing	{	Fibrin.
		Clot or coagulum.				Corpuscles.
		Serum.		Containing	{	Water.
						Albumen.
						Salts.

The serum, therefore, differs from the *Liquor Sanguinis* in not containing fibrin.

In from 12 to 24 hours the upper surface of the clot presents a grayish appearance, the *buffy coat*, which is due to the rapid sinking of the red corpuscles beneath the surface, permitting the fibrin to coagulate without them, which then assumes a grayish-yellow tint. Inasmuch as the white corpuscles possess a lighter specific gravity than the red, they do not sink so rapidly, and becoming entangled in the fibrin, assist in forming the buffy coat. Continued contraction gives a *cupped* appearance to the surface of the clot.

Inflammatory states of the blood produce a marked increase in the buffed and cupped condition, on account of the aggregation of the corpuscles and their tendency to rapid sinking.

Nature of Coagulation. Coagulated fibrin does not pre-exist in the blood, but is formed at the moment blood is withdrawn from the vessels. According to Denis, a liquid substance, *plasmin*, exists in the blood, which, when withdrawn from the circulation, decomposes into *fibrin* and *met-albumen*.

According to Schmidt, fibrin results from the union of *fibrinoplastin* (paraglobulin) and *fibrinogen*, brought about by the presence of a third substance, the fibrin ferment.

According to Hammersten and others, the fibrin obtained from the blood after coagulation, comes from the fibrinogen alone, the conversion being brought about by the presence of a ferment substance. Paraglobulin in this case having nothing to do with the change. This view is supported by the fact that the quantity of fibrin obtained from the blood is never greater than the quantity of fibrinogen previously present. The origin of the ferment is obscure, but there is reason to believe that it comes from the injured vascular coats or from the breaking up of the white corpuscles.

Conditions Influencing Coagulation. The process is *retarded* by cold, retention within living vessels, neutral salts in excess, inflammatory conditions of the system, imperfect aeration, exclusion from air, etc.

It is *hastened* by a temperature of 100° F., contact with air, rough surfaces and rest.

Blood coagulates in the body after the arrest of the circulation in the course of 12 to 24 hours; local arrest of the circulation, from compression or a ligature, will cause coagulation, thus preventing hemorrhages from wounded vessels.

The Composition of the Blood varies in different portions of the body. The *arterial* differs from the *venous*, in being more coagulable, in containing more oxygen and less carbonic acid, in having a bright scarlet color, from the union of oxygen with hæmoglobin; the purple hue of venous blood results from the deoxidation of the coloring matter.

The blood of the *portal vein* differs in constitution, according to different stages of the digestive process; during digestion it is richer in water, albuminous matter and sugar; occasionally it contains fat; corpuscles are diminished, and there is an absence of biliary substances.

The blood of the *hepatic vein* contains a larger proportion of red and

white corpuscles; the sugar is augmented, while albumen, fat and fibrin are diminished.

Pathological conditions of the Blood.

1. *Plethora*—increase in the volume or quantity of blood.
2. *Anæmia*—deficiency of red globules with increase of water.
3. *Leucocythemia*—increase of white and diminution of red corpuscles.
4. *Glycohæmia*—excess of sugar in the blood.
5. *Uræmia*—increase in the amount of urea.
6. *Cholesteræmia*—an excess of cholesterine in the blood.
7. *Thrombosis* and *embolism*—clotting of blood in the vessels and dissemination of coagula.
8. *Lipæmia*—an excess of fat.
9. *Melanæmia*—pigment in the blood.

CIRCULATION OF THE BLOOD.

The Object of the Circulation is to distribute nutritious blood to all portions of the system and to carry waste materials to the various eliminating organs.

The Circulatory Apparatus consists of the heart, arteries, capillaries and veins.

The Heart is a hollow, muscular organ, pyramidal in shape, measuring $5\frac{1}{2}$ inches in length and weighing from 10 to 12 oz. in the male, and 8 to 10 oz. in the female. It is invested externally by a closed fibro-serous sac, the *pericardium*, containing a small amount of fluid, which prevents friction as the visceral and parietal layers glide over each other, during the movements of the heart and lungs.

The heart consists of four cavities, a right auricle and ventricle, and a left auricle and ventricle, completely separated by a vertical partition. The *right* is the *venous side*, receiving the blood from the *venæ cavæ*, and propelling it through the pulmonary artery into the lungs; the *left* is the *arterial side*, receiving the arterial blood from the lungs by the pulmonary veins, and propelling it through the aorta to the system at large.

The *Auriculo-ventricular orifices* are guarded on the right and left sides by the *tricuspid* and *mitral* valves respectively, while they are so arranged as to permit the flow of blood in the forward direction only; the orifices of the *pulmonary artery* and *aorta* are guarded by the *semi-lunar* valves.

The *Endocardium* is a delicate, shining membrane, lining the interior of the heart, and continuous with the lining membrane of the blood vessels.

The walls of the left ventricle are nearly half an inch in diameter, being

two or three times thicker than the walls of the right; the force of its contraction is, therefore, much greater.

The Function of the Heart is to propel the blood to all portions of the vascular system; accomplished by successive alternate contractions and relaxations of its muscular walls, constituting the *systole* and *diastole*.

Course of the blood through the Heart. The venous blood returned to the heart by the superior and inferior venæ cavæ is emptied, during the diastole, into the right auricle, on the contraction of which it is forced through the right auriculo-ventricular opening into the right ventricle and distends it. Upon contraction of the ventricle, the blood is propelled through the pulmonary artery into the lungs, where it undergoes aeration and is changed in color.

The arterial blood is now collected by the pulmonary veins and poured into the left auricle; thence it passes into the left ventricle, which becomes fully distended. Upon the contraction of the ventricle, the blood is propelled into the aorta, and by it distributed to the system at large, to be again returned to the heart by the veins.

Regurgitation from the ventricles into the auricles during the systole is prevented by the closure of the tricuspid and mitral valves; regurgitation from the pulmonary artery and aorta into the ventricles during the diastole is prevented by the closure of the semi-lunar valves.

Movements of the Heart. At each revolution, during the systole, the heart hardens and becomes shortened in its long diameter; its apex is raised up, rotated on its axis from left to right and thrown forward against the walls of the chest. The *impulse* of the heart, observed about two inches below the nipple, and one inch to the sternal side, between the fifth and sixth ribs, is caused mainly by the apex of the heart striking against the chest walls, assisted by the distention of the great vessels about the base of the heart.

Sounds of the Heart. If the ear be placed over the cardiac region, two distinct sounds are heard during each revolution of the heart, closely following each other and which differ in character.

The sound coinciding with the systole in point of time, the *first sound*, is long and dull, and caused by the closure and vibration of the auriculo-ventricular valves, the contraction of the walls of the ventricles and the apex beat; the *second sound*, occurring during the *diastole*, is short and sharp, and caused by the closure of the semi-lunar valves.

The *capacity* of the left ventricle when fully distended is estimated at from four to seven ounces.

The frequency of the heart's action varies at different periods of life, but in the adult male it beats about 72 times per minute. It is influenced by age, exercise, posture, digestion, etc.

Age.	Before birth the <i>number of pulsations</i> per minute averages	140
	During the first year it diminishes to.....	128
	During the third year diminishes to.....	95
	From the eighth to the fourteenth years averages.....	84
	In adult life the average is.....	72

Exercise and digestion increase the frequency of the heart's action.

Posture influences the number of pulsations per minute; in the male, standing, the average is 81; sitting, 71; lying, 66; independent, for the most part, of muscular effort.

The Rhythmical movements of the heart are dependent upon—1. An inherent irritability of the muscular fibre, which manifests itself as long as the nutrition is maintained. 2. The continuous flow of blood through its cavities distending them and stimulating the endocardium.

The force exerted by the left ventricle at each contraction has been estimated at 52 pounds. If a tube be inserted into the aorta, the pressure there will be sufficient to support a column of blood nine feet or a column of mercury six inches in height, the weight in either case being about four pounds. The estimation of the force which the heart is required to exert to support this column of blood, is arrived at by multiplying the pressure in the aorta (4 pounds) by the area of the internal surface of the left ventricle (about 13 inches²). Each inch of the ventricle being capable of supporting a downward pressure of 4 pounds.

Work done by the Heart. The work done by the heart is estimated by multiplying the amount of blood sent out from the right and left ventricles at each contraction, by the pressure in the pulmonary artery and aorta respectively, *e. g.*, when the right ventricle contracts, it forces out one quarter pound of blood, and in so doing must overcome a pressure in the pulmonary artery sufficient to support a column of blood three feet in height; that is, must exert energy sufficient to raise $\frac{1}{4}$ lb. 3 feet, or $\frac{1}{4} \times 3$ or $\frac{3}{4}$ lb. one foot. When the left ventricle contracts, it sends out $\frac{1}{4}$ lb. of blood, and in so doing, the left ventricle must overcome a pressure in the aorta sufficient to support a column of blood nine feet in height; that is must exert energy sufficient to raise $\frac{1}{4}$ lb. 9 feet, or $\frac{1}{4} \times 9$ or $2\frac{1}{4}$ lbs. one foot. Work done is estimated by the amount of energy required to raise a definite weight a definite height, the unit, the foot pound, being that required to raise one pound one foot.

The heart, therefore, at each systole exerts energy sufficient to raise 3 foot pounds, and as it contracts 72 times per minute, it would raise in that time 3×72 or 216 foot pounds; and in one hour 216×60 or 12,960 foot pounds; and in 24 hours $12,960 \times 24$ or 311,040 foot pounds or 138.5 foot tons.

Influence of the Nervous System upon the Heart. When the heart of a frog is removed from the body, it continues to beat for a variable length of time, depending upon the nature of the conditions surrounding it. The heart of warm-blooded animals continues to beat but for a very short time. The cause of the continued pulsations of the frog heart is the presence of nervous ganglia in its substance. These ganglia have not been shown to exist in the mammalian heart, but there is reason to believe that the nervous mechanism is fundamentally the same.

The ganglia of the heart are three in number, one situated at the opening of the inferior vena cava (the ganglion of Remak), a second situated in the auriculo-ventricular septum (the ganglion of Bidder), and a third situated in the inter-auricular septum (the ganglion of Ludwig). The first two are motor in function and excite the pulsations of the heart; the third is inhibitory in function and retards the action of the heart. The actions of these ganglia, though for the most part automatic, are modified by impressions coming through nerves from the medulla oblongata. When the inhibitory centre is stimulated by muscarin, the heart is arrested in diastole; when atropia is applied, the heart recommences to beat, because atropia paralyzes the inhibitory centre.

The nerves modifying the action of the heart are the Pneumogastric (Vagus) and the Accelerator nerves.

The *Pneumogastric nerve*, after emerging from the medulla, receives motor fibres from the spinal accessory nerve. It then passes downward, giving off branches, some of which terminate in the inhibitory ganglion. *Stimulation* of the vagus by increasing the activity of the inhibitory centre arrests the heart in diastole with its cavities full of blood; but as the stimulation is only temporary, after a few seconds the heart recommences to beat; at first the pulsations are weak and feeble, but soon regain their original vigor. After the administration of atropia in sufficient doses to destroy the termination of the pneumogastric, stimulation of its trunk has no effect upon the heart. The inhibitory fibres in the vagus are constantly in action, for *division* of the nerve on both sides is always followed by an increase in the frequency of the heart's pulsations.

The *Accelerator fibres* arise in the medulla, pass down the cord, emerge in the cervical region, pass to the last cervical and first dorsal ganglia of the sympathetic, and thence to the heart. *Stimulation* of these fibres

causes an increased frequency of the heart's pulsations, but they are diminished in force.

ARTERIES.

The **Arteries** are a series of branching tubes conveying blood to all portions of the body. They are composed of three coats—

1. *External*, formed of areolar and elastic tissue.
2. *Middle*, contains both elastic and muscular fibres, arranged transversely to the long axis of the artery. The elastic tissue is more abundant in the larger vessels, the muscular in the smaller.
3. *Internal*, composed of a thin homogeneous membrane, covered with a layer of elongated endothelial cells.

The arteries possess both *elasticity* and *contractility*.

The *Property of Elasticity* allows the arteries already full to accommodate themselves to the incoming amount of blood, and to convert the intermittent acceleration of blood in the large vessels into a steady and continuous stream in the capillaries.

The *Contractility* of the smaller vessels equalizes the current of blood, regulates the amount going to each part, and promotes the onward flow of blood.

Blood Pressure. Under the influence of the ventricular systole, the recoil of the elastic walls of the arteries, and the resistance offered by the capillaries, the blood is constantly being subjected to a certain amount of pressure. If a large artery of an animal be divided, and a glass tube of the same calibre be inserted into its orifice, the blood will rise to a height of about nine feet; or if it be connected with a mercurial manometer, the mercury will rise to a height of six inches. This height will be a measure of the pressure in the vessel. The absolute quantity of mercury sustained by an artery can be arrived at by multiplying the height of the column by the area of a transverse section of that artery.

The pressure of the blood is greatest in the large arteries, but gradually decreases toward the capillaries.

The blood pressure is increased or diminished by influences acting upon the heart or upon the peripheral resistance of the capillaries, viz. :—

If, while the force of the heart remains the same, the number of pulsations per minute increases, thus increasing the volume of blood in the arteries, the pressure rises. If the rate remains the same, but the force increases, the pressure again rises. Causes that increase the peripheral resistance by contracting the arterioles, *e. g.*, vaso-motor nerves, cold, etc., produce an increase of the pressure.

On the other hand, influences which *diminish* either the volume of the blood, or the number of pulsations, or the force of the heart, or the peripheral resistance, *lower* the pressure.

The Pulse is the sudden distention of the artery in a transverse and longitudinal direction, due to the injection of a volume of blood into the arteries at the time of the ventricular systole. As the vessels are already full of blood, they must expand in order to accommodate themselves to the incoming volume of blood. The blood pressure is thus increased, and the pressure originating at the ventricle excites a *pulse wave*, which passes from the heart toward the capillaries at the rate of about twenty-nine feet per second. It is this wave that is appreciated by the finger.

The Velocity with which the blood flows in the arteries diminishes from the heart to the capillaries, owing to an increase of the united sectional area of the vessels, and increases in rapidity from the capillaries toward the heart. It moves most rapidly in the large vessels, and especially under the influence of the ventricular systole. From experiments on animals, it has been estimated to move in the carotid of man at the rate of sixteen inches per second, and in the large veins at the rate of four inches per second.

The Calibre of the blood vessels is regulated by the *vaso-motor* nerves, which have their origin in the gray matter of the medulla oblongata. They issue from the spinal cord through the anterior roots of spinal nerves, pass through the sympathetic ganglia, and ultimately are distributed to the coats of the blood vessels. They exert, at different times, a constricting and dilating action upon the vessels, thus keeping up the arterial tonus.

Capillaries. The capillaries constitute a network of vessels of microscopic size, which distribute the blood to the inmost recesses of the tissues, inosculating with the arteries on the one hand and the veins on the other; they branch and communicate in every possible direction.

The diameter of a capillary vessel varies from the $\frac{1}{80000}$ to the $\frac{1}{30000}$ of an inch; their walls consist of a delicate homogeneous membrane, the $\frac{1}{80000}$ of an inch in thickness, lined by flattened, elongated, endothelial cells, between which, here and there, are observed *stomata*.

It is through the agency of the capillary vessels that the phenomena of nutrition and secretion takes place, for here the blood flows in an equable and continuous current, and is brought into intimate relationship with the tissues, two of the essential conditions for proper nutrition.

The *rate of movement* in the capillary vessels is estimated at one inch in thirty seconds.

In the capillary current the red corpuscles may be seen hurrying down the centre of the stream, while the white corpuscles in the still layer adhere to the walls of the vessel, and at times can be seen to pass through the walls of the vessel by amœboid movements.

The passage of the blood through the capillaries is mainly due to the force of the ventricular systole and the elasticity of the arteries; but it is probably also aided by a power resident in the capillaries themselves, the result of a vital relation between the blood and the tissues.

The Veins are the vessels which return the blood to the heart; they have their origin in the venous radicles, and as they approach the heart, converge to form larger trunks, and terminate finally in the venæ cavæ.

They possess three coats—

1. *External*, made up of areolar tissue.
2. *Middle*, composed of non-striated muscular fibres, yellow, elastic and fibrous tissue.
3. *Internal*, an endothelial membrane, similar to that of the arteries.

Veins are distinguished by the possession of valves throughout their course, which are arranged in pairs, and formed by a reflection of the internal coat, strengthened by fibrous tissue; they always look toward the heart, and when closed prevent a return of blood in the veins. Valves are most numerous in the veins of the extremities, but are entirely absent in many others.

The onward flow of blood in the veins is mainly due to the action of the heart; but is assisted by the contraction of the voluntary muscles and the force of aspiration.

Muscular contraction, which is intermittent, aids the flow of blood in the veins, by compressing them. As regurgitation is prevented by the closure of the valves, the blood is forced onward toward the heart.

Rhythmical movements of veins have been observed in some of the lower animals, aiding the onward current of blood.

During the movement of *inspiration* the thorax is enlarged in all its diameters, and the pressure on its contents at once diminishes. Under these circumstances a suction force is exerted upon the great venous trunks, which causes the blood to flow with increased rapidity and volume toward the heart.

Venous pressure. As the force of the heart is nearly expended in driving the blood through the capillaries, the pressure in the venous system is not very marked, not amounting in the jugular vein of a dog to more than $\frac{1}{12}$ that of the carotid artery.

The time required for a complete circulation of the blood throughout the vascular system has been estimated to be from 20 to 30 seconds, while for the *entire mass* of blood to pass through the heart 58 pulsations would be required, occupying 48 seconds.

The Forces keeping the blood in circulation are—

1. Action of the heart.
2. Elasticity of the arteries.
3. Capillary force.
4. Contraction of the voluntary muscles upon the veins.
5. Respiratory movements.

RESPIRATION.

Respiration is the function by which oxygen is absorbed into the blood and carbonic acid exhaled. The appropriation of the oxygen and the evolution of carbonic acid takes place in the tissues as a part of the general nutritive process; the blood and respiratory apparatus constituting the media by means of which the interchange of gases is accomplished.

The Respiratory Apparatus consists of the larynx, trachea and lungs.

The Larynx is composed of firm cartilages, united together by ligaments and muscles; running antero-posteriorly across the upper opening are four ligamentous bands, the two superior or *false* vocal cords, and the two inferior or *true* vocal cords, formed by folds of the mucous membrane. They are attached anteriorly to the thyroid cartilages and posteriorly to the arytenoid cartilages, and are capable of being separated by the contraction of the posterior crico-arytenoid muscles, so as to admit the passage of air into and from the lungs.

The Trachea is a tube from four to five inches in length, three-quarters of an inch in diameter, extending from the cricoid cartilage of the larynx to the third dorsal vertebra, where it divides into the right and left bronchi. It is composed of a series of cartilaginous rings, which extend about two-thirds around its circumference, the posterior third being occupied by fibrous tissue and non-striated muscular fibres which are capable of diminishing its calibre.

The trachea is covered externally by a tough, fibro-elastic membrane, and internally by mucous membrane, lined by columnar ciliated epithelial cells. The cilia are always waving from within outward. When the two bronchi enter the lungs they divide and subdivide into numerous and

smaller branches, which penetrate the lung in every direction until they finally terminate in the *pulmonary lobules*.

As the bronchial tubes become smaller their walls become thinner; the cartilaginous rings disappear, but are replaced by irregular angular plates of cartilage; when the tube becomes less than the $\frac{1}{50}$ of an inch in diameter they wholly disappear, and the fibrous and mucous coats blend together, forming a delicate, elastic membrane, with circular muscular fibres.

The Lungs occupy the cavity of the thorax, are conical in shape, of a pink color and a spongy texture. They are composed of a great number of distinct lobules, the *pulmonary lobules*, connected together by inter-lobular connective tissue. These lobules vary in size, are of an oblong shape, and are composed of the ultimate ramifications of the bronchial tubes, within which are contained the *air vesicles or cells*. The walls of the air vesicles, exceedingly thin and delicate, are lined internally by a layer of tessellated epithelium, externally covered by elastic fibres, which give the lungs their elasticity and distensibility.

The Venous Blood is distributed to the lungs for aeration by the pulmonary artery, the terminal branches of which form a rich plexus of capillary vessels surrounding the air cells; the air and blood are thus brought into intimate relationship, being separated only by the delicate walls of the air cells and capillaries.

The Pleura. Each lung is surrounded by a closed serous membrane, the pleura, one layer of which, the *visceral*, is reflected over the lung, the other, the *parietal*, reflected over the wall of the thorax; between the two layers is a small amount of fluid which prevents friction during the play of the lungs in respiration.

The lungs are nourished by blood from the bronchial arteries ramifying in the walls of the bronchial tubes and interlobular connective tissue.

FIG 7.




Diagram of the respiratory organs. The windpipe leading down from the larynx is seen to branch into two large bronchi, which subdivide after they enter their respective lungs.

Respiratory movements. The movements of respiration are two, and consist of an alternate dilatation and contraction of the chest, known as inspiration and expiration.

1. *Inspiration* is an active process, the result of the expansion of the thorax, whereby air is introduced into the lungs.

2. *Expiration* is a partially passive process, the result of the recoil of the elastic walls of the thorax, and the recoil of the elastic tissue of the lungs, whereby the carbonic acid is expelled.

In Inspiration the chest is enlarged by an increase in all its diameters, viz.:—

1. The *vertical* is increased by the contraction and descent of the diaphragm when it approximates a straight line.

2. The *antero-posterior* and *transverse* diameters are increased by the elevation and rotation of the ribs upon their axes.

In *ordinary tranquil inspiration* the muscles which elevate the ribs and thrust the sternum forward, and so increase the diameters of the chest, are the *external intercostals*, running from above downward and forward, the *sternal portion* of the *internal intercostals* and the *levator costarum*.

In the *extraordinary efforts* of inspiration certain auxiliary muscles are brought into play, viz.; the *sterno-mastoid*, *pectorales*, *serratus magnus*, which increase the capacity of the thorax to its utmost limit.

In Expiration the diameters of the chest are all diminished, viz. :

1. The *vertical*, by the ascent of the diaphragm.

2. The *antero-posterior*, by a depression of the ribs and sternum.

In *ordinary tranquil expiration* the diameters of the thorax are diminished by the recoil of the elastic tissue of the lungs and the ribs; but in forcible expiration the muscles which depress the ribs and sternum, and thus further diminish the diameter of the chest, are the *internal intercostals*, the *infracostals*, and the *triangularis sterni*.

In the *extraordinary efforts of expiration* certain auxiliary muscles are brought into play, viz.: the *abdominal* and *sacro-lumbalis* muscles, which diminish the capacity of the thorax to its utmost limit.

Expiration is aided by the recoil of the elastic tissue of the lungs and ribs and the pressure of the air.

Movements of the Glottis. At each inspiration the *rima glottidis* is dilated by a separation of the vocal cords, produced by the contraction of the *crico-arytenoid* muscles, so as to freely admit the passage of air into the lungs; in expiration they fall passively together, but do not interfere with the exit of the air from the chest.

Nervous Mechanism of Respiration. The movements of respiration are involuntary and reflex, and are under the control of the *medulla oblongata*.

This centre may be stimulated—

1. *Directly*, by the condition of the blood. An increase of carbonic acid or a diminution of oxygen in the blood causes an acceleration of the respiratory movements; the reverse of these conditions causes a diminution of the respiratory movements.

2. *Indirectly*, by reflex action. The medulla may be excited to action through the pneumogastric nerve, by the presence of carbonic acid in the lungs irritating its terminal filaments; through the fifth nerve, by irritation of the terminal branches; and through the nerves of general sensibility. In either case this centre reflects *motor* impulses to the respiratory muscles through the *phrenic, intercostals, inferior laryngeal* and other nerves.

Types of Respiration. The *abdominal type* is most marked in young children, irrespective of sex; the respiratory movements being effected by the diaphragm and abdominal muscles.

In the *superior costal type*, exhibited by the adult female, the respiratory movements are more marked in the upper part of the chest, from the 1st to the 7th ribs, permitting the uterus to ascend in the abdomen during pregnancy without interfering with respiration.

In the *inferior costal type*, manifested by the male, the movements are largely produced by the muscles of the lower portion of the chest, from the 7th rib downward, assisted by the diaphragm.

The respiratory movements vary according to age, sleep and exercise, being most frequent in early life, but averaging 20 per minute in adult life. They are diminished by sleep and increased by exercise. There are about four pulsations of the heart to each respiratory act.

During inspiration *two sounds* are produced; the one, heard in the thorax, in the trachea and larger bronchial tubes, is *tubular* in character; the other, heard in the substance of the lungs, is *vesicular* in character.

AMOUNT OF AIR EXCHANGED IN RESPIRATION, AND CAPACITY OF LUNGS.

The *Tidal* or *breathing volume* of air, that which passes in and out of the lungs at each inspiration and expiration, is estimated at from 20 to 30 cubic inches.

The *Complemental air* is that amount which can be taken into the lungs by a forced inspiration, in addition to the ordinary tidal volume, and amounts to about 110 cubic inches.

The *Reserve air* is that which usually remains in the chest after the ordinary efforts of expiration, but which can be expelled by forcible expiration. The volume of reserve air is about 100 cubic inches.

The *Residual air* is that portion which remains in the chest and cannot be expelled after the most forcible expiratory efforts, and which amounts, according to Dr. Hutchinson, to about 100 cubic inches.

The Vital Capacity of the chest indicates the amount of air that can be forcibly expelled from the lungs after the deepest possible inspiration, and is an index of an individual's power of breathing in disease and prolonged severe exercise. The combined amounts of the tidal, the complementary and reserve air, 230 cubic inches, represents the vital capacity of an individual 5 feet 7 inches in height. The vital capacity varies chiefly with stature. It is increased 8 cubic inches for every inch in height above this standard, and diminishes 8 cubic inches for each inch below it.

The Tidal Volume of air is carried only into the trachea and larger bronchial tubes by the inspiratory movements. It reaches the deeper portions of the lungs in obedience to the law of diffusion of gases, which is inversely proportionate to the square root of their densities.

The *ciliary action* of the columnar cells lining the bronchial tubes also assists in the interchange of air and carbonic acid.

The *entire volume of air* passing in and out of the thorax in 24 hours is subject to great variation, but can be readily estimated from the tidal volume and the number of respirations per minute. Assuming that an individual takes into the chest 20 cubic inches at each inspiration, and breathes 18 times per minute, in 24 hours there would pass in and out of the lungs 518,400 cubic inches or 300 cubic feet.

Composition of Air: Oxygen, 20.81 parts; nitrogen, 79.19, forming a mechanical mixture in which exist traces of carbonic acid and watery vapor.

The changes in the air effected by respiration are—

Loss of oxygen, to the extent of 5 cubic inches per 100 of air, or 1 in 20.

Gain of carbonic acid, to the extent of 4.66 cubic inches per 100 of air or .93 inch in 20.

Increase of watery vapor and organic matter.

Elevation of temperature.

Increase and at times decrease of nitrogen.

Gain of ammonia.

The *total quantity of oxygen* withdrawn from the air and consumed by

the body in 24 hours amounts to 15 cubic feet, and can be readily estimated from the amount consumed at each respiration. Assuming that one inch of oxygen remains in the lungs at each respiration, in one hour there are consumed 18 inches, and in 24 hours, 25,920 cubic inches or 15 cubic feet, weighing 18 oz. To obtain this quantity, 300 cubic feet of air are necessary.

The *quantity of carbonic acid* exhaled in 24 hours varies greatly. It can be estimated in the same way. Assuming that an individual exhales .93 + cubic inch at each respiration, in one hour there are eliminated 1008 cubic inches, and in 24 hours 24,192 cubic inches or 14 cubic feet, containing 7 oz. of pure carbon.

As oxygen and carbon unite to form an equal volume of carbonic acid gas, there disappears daily in the body, one cubic foot of oxygen, which in all probability unites with the surplus hydrogen of the food to form water.

The exhalation of carbonic acid is *increased* by muscular exercise; nitrogenous food; tea, coffee and rice; age, and by muscular development; *decreased* by a lowering of temperature; repose; gin and brandy, and a dry condition of the air.

Condition of the Gases in the Blood.

Oxygen is absorbed from the lungs into the arterial blood by the coloring matter, *hæmoglobin*, with which it exists in a state of loose combination, and is disengaged during the process of nutrition.

Carbonic acid, arising in the tissues, is absorbed into the blood, in consequence of its alkalinity; where it exists in a state of simple solution and also in a state of feeble combination with the carbonates, soda and potassa, forming the bicarbonates; it is liberated by pneumatic acid in the pulmonary tissue.

Nitrogen is simply held in solution in the plasma.

The *amount of watery vapor* thrown off from the lungs daily is about one pound, with which is mingled organic matter and ammonia.

Changes in the Blood during Respiration.

As the blood passes through the lungs it is changed in *color*, from the dark purple hue of venous blood to the bright scarlet of arterial blood.

The heterogeneous composition of venous blood is exchanged for the uniform composition of the arterial.

It gains oxygen and loses carbonic acid.

Its coagulability is increased. Temperature is diminished.

Asphyxia. If the supply of oxygen to the lungs be diminished and the carbonic acid retained in the blood, the normal respiratory movements cease, the condition of asphyxia ensues, which soon terminates in death.

The *phenomena* of asphyxia are, violent spasmodic action of the respiratory muscles, attended by convulsions of the muscles of the extremities, engorgement of the venous system, lividity of the skin, abolition of sensibility and reflex action, and death.

The *cause of death* is a paralysis of the heart, from over distention by blood. The passage of the blood through the capillaries is prevented by contraction of the smaller arteries, from irritation of the vaso-motor centre. The heart is enfeebled by a want of oxygen and inhibited in its action by the inhibitory centres.

ANIMAL HEAT.

The Functional Activity of all the organs and tissues of the body is attended by the evolution of heat, which is independent, for the most part, of external conditions. Heat is a *necessary condition* for the due performance of all vital actions; though the body constantly loses heat by *radiation* and *evaporation*, it possesses the capability of renewing it and maintaining it at a fixed standard. The *normal temperature* of the body in the adult, as shown by means of a delicate thermometer placed in the axilla, ranges from 97.25° Fahr. to 99.5° Fahr., though the *mean normal temperature* is estimated by Wunderlich at 98.6° Fahr.

The temperature varies in different portions of the body, according to the degree in which oxidation takes place; being the highest in the muscles during exercise, in the brain, blood, liver, etc.

The conditions which produce variations in the normal temperature of the body are: age, period of the day, exercise, food and drink, climate, season and disease.

Age. At birth the temperature of the infant is about 1° F. above that of the adult, but in a few hours falls to 95.5° F., to be followed in the course of 24 hours by a rise to the normal or a degree beyond. During childhood the temperature approaches that of the adult; in aged persons the temperature remains about the same, though they are not as capable of resisting the depressing effects of external cold as adults. A *diurnal variation* of the temperature occurs from 1.8° F. to 3.6° F. (Jürgensen); the *maximum* occurring late in the afternoon, from 4 to 9 P.M., the *minimum*, early in the morning, from 1 to 7 A.M.

Exercise. The temperature is raised from 1° to 2° F. during active

contractions of the muscular masses, and is probably due to the increased activity of chemical changes; a rise beyond this point being prevented by its diffusion to the surface, consequent on a more rapid circulation, radiation, more rapid breathing, etc.

Food and drink. The ingestion of a hearty meal increases the temperature but slightly; an absence of food, as in starvation, produces a marked decrease. Alcoholic drinks, in large amounts, in persons unaccustomed to their use, cause a depression of the temperature, amounting from 1° to 2° F. Tea causes a slight elevation.

External temperature. Long continued exposure to cold, especially if the body is at rest, diminishes the temperature from 1° to 2° F., while exposure to a great heat slightly increases it.

Disease frequently causes a marked variation in the normal temperature of the body, rising as high as 107° F. in typhoid fever, and 105° F. in pneumonia; in cholera it falls as low as 80° F. Death usually occurs when the heat remains high and persistent, from 106° to 110° F.; the increase of heat in disease is due to excessive production rather than to diminished elimination.

The source of heat is to be sought for in the chemical combinations taking place during the general process of nutrition, and the amount of its production is in proportion to the activity of the internal changes.

Every contraction of a muscle, every act of secretion, each exhibition of nerve force, is accompanied by a change in the chemical composition of the tissues and an evolution of heat. The reduction of the disintegrated tissues to their simplest form by oxidation; the combination of the oxygen of the inspired air with the carbon and hydrogen of the blood and tissues, results in the formation of carbonic acid and water and the generation of a large amount of heat.

Certain elements of the food, particularly the *non-nitrogenized* substances, undergo oxidation without taking part in the formation of the tissues, being transformed into carbonic acid and water, and thus increase the sum of heat in the body.

Heat-producing Tissues. All the tissues of the body add to the general amount of heat, according to the degree of their activity. But special structures, on account of their mass and the large amount of blood they receive, are particularly to be regarded as *heat producers*; e. g.:

1. During mental activity the *brain* receives nearly one-fifth of the entire volume of blood, and the venous blood returning from it is charged with waste matters, and its temperature is increased.

2. The *muscular tissue*, on account of the many chemical changes occurring during active contractions, must be regarded as the chief heat-producing tissue.

3. The *secreting glands*, during their functional activity, add largely to the amount of heat.

Of the entire quantity of heat generated in the body, it is estimated that only a small proportion is utilized, as five-sixths escape by radiation and evaporation, the remaining one-sixth being utilized in keeping the body at the normal temperature standard, 98.6° F., and in the production of muscular force.

The *body loses heat* by radiation and evaporation from the general cutaneous surface, the respiratory passages and by the urine and fæces. About 75 per cent. of all the heat lost escapes from the skin. In passing through the lungs the temperature of the blood is lowered by about 1° Fahr.

The *nervous system* influences the production of heat in a part, by increasing the amount of blood going through it by its action upon the vaso-motor nerves. Whether there exists a special heat centre has not been satisfactorily determined, though this is probable.

SECRETION.

The **Process of Secretion** consists in the separation of materials from the blood, which are either to be again utilized to fulfill some special purpose in the economy, or are to be removed from the body as excrementitious matter; in the former case they constitute the *secretions*, in the latter, the *excretions*.

The *materials* which enter into the composition of the *secretions* are derived from the nutritive principles of the blood, and require special organs, *e. g.*, gastric glands, mammary glands, etc., for their proper elaboration.

The *materials* which compose the *excretions* pre-exist in the blood, and are the results of the activities of the nutritive process; if retained within the body they exert a deleterious influence upon the composition of the blood.

Destruction of a secreting gland abolishes the secretion peculiar to it, and it cannot be formed by any other gland; but among the excreting organs there exists a complementary relation, so that if the function of one organ be interfered with, another performs it, to a certain extent.

CLASSIFICATION OF THE SECRETIONS.

PERMANENT FLUIDS.

Serous fluids.	Vitreous humor of the eye.
Synovial fluid.	Fluid of the labyrinth of the internal ear.
Aqueous humor of the eye.	Cerebro-spinal fluid.

TRANSITORY FLUIDS.

Mucus.	Gastric juice.
Sebaceous matter.	Pancreatic juice.
Cerumen (external meatus).	Secretion from Brunner's glands.
Meibomian fluid.	Secretion from Leiberkühn's glands.
Milk and colostrum.	Secretion from follicles of the large intestine.
Tears.	Bile (also an excretion).
Saliva.	

EXCRETIONS.

Perspiration and the secretion of the axillary glands.	Urine.
	Bile (also a secretion).

FLUIDS CONTAINING FORMED ANATOMICAL ELEMENTS.

Seminal fluid, containing spermatozoids. Fluid of the Graafian follicles.

The essential apparatus for secretion is a delicate, homogeneous, structureless *membrane*, on one side of which, in close contact, is a capillary plexus of *blood vessels*, and on the other side a layer of *cells* whose physiological function varies in different situations.

Secreting organs may be divided into *membranes* and *glands*.

Serous membranes usually exist as closed sacs, the inner surface of which is covered by pale, nucleated epithelium, containing a small amount of secretion.

The serous membranes are the *pleura*, *peritoneum*, *pericardium*, *synovial sacs*, etc.

The *serous fluids* are of a pale amber color, somewhat viscid, alkaline, coagulable by heat, and resemble the serum of the blood; their *amount* is but small; the pleural varies from 4 to 7 drachms; the peritoneal from 1 to 4 ounces; the pericardial from 1 to 3 drachms.

The *synovial fluid* is colorless, alkaline, and extremely viscid, from the presence of *synovine*.

The *function* of serous fluids is to moisten the opposing surfaces, so as to prevent friction during the play of the viscera.

The *mucous membranes* are soft and velvety in character, and line the cavities and passages leading to the exterior of the body, *e. g.*, the *gastro*

intestinal, pulmonary and genito-urinary. They consist of a primary basement membrane covered with epithelial cells, which, in some situations, are tessellated, in others, columnar.

Mucus is a pale, semi-transparent, alkaline fluid, containing epithelial cells and leucocytes. It is composed, chemically, of water, an albuminous principle, *mucosine*, and mineral salts; the principal varieties are nasal, bronchial, vaginal and urinary.

Secreting Glands are formed of the same elements as the secreting membranes; but instead of presenting flat surfaces, are involuted, forming tubules, which may be *simple follicles*, *e. g.*, mucous, uterine or intestinal; or *compound follicles*, *e. g.*, gastric glands, mammary glands; or *racemose glands*, *e. g.*, salivary glands and pancreas. They are composed of a basement membrane, enveloped by a plexus of blood vessels, and are lined by epithelial and *true secreting cells*, which in different glands possess the capability of elaborating elements characteristic of their secretions.

In the production of the secretions two essentially different processes are concerned:—

1. *Chemical.* The formation and elaboration of the characteristic organic ingredients of the secreting fluids, *e. g.*, pepsin, pancreatin, takes place during the intervals of glandular activity, as a part of the general function of nutrition. They are formed by the cells lining the glands, and can often be seen in their interior with the aid of the microscope, *e. g.*, bile in the liver cells, fat in the cells of the mammary gland.

2. *Physical.* Consisting of a transudation of water and mineral salts from the blood into the interior of the gland.

During the intervals of glandular activity, only that amount of blood passes through the gland sufficient for proper nutrition; when the gland begins to secrete, under the influence of an appropriate stimulus, the blood vessels dilate and the quantity of blood becomes greatly increased beyond that flowing through the gland during its repose.

Under these conditions a transudation of water and salts takes place, washing out the characteristic ingredients, which are discharged by the gland ducts. The *discharge* of the secretions is intermittent; they are retained in the glands until they receive the appropriate stimulus, when they pass into the larger ducts by the *vis-a-tergo*, and are then discharged by the contraction of the muscular walls of the ducts.

The *activity* of glandular secretion is hastened by an increase in the blood pressure and retarded by a diminution.

The *nervous centres* in the medulla oblongata influence secretion, (1) by

increasing or diminishing the amount of blood entering a gland ; (2) by exerting a direct influence upon the secreting cells themselves, the centres being excited by reflex irritation, mental emotion, etc.

MAMMARY GLANDS.

The Mammary Glands secrete the milk, and undergo at different periods of life remarkable changes in structure. Though rudimentary in childhood, they gradually increase in size as the young female approaches puberty.

The gland presents, at its convexity, a small prominence of skin, the *nipple*, surrounded by an *areola* of a deeper tint. It is covered anteriorly by a layer of adipose tissue and posteriorly by a fibrous structure which attaches it loosely to the *pectoralis* muscle.

During *utero-gestation* the mammæ become large, firm, well-developed and lobulated ; the areola becomes darker and the veins more prominent. In the intervals of lactation the glands gradually shrink in size to their original condition, undergo involution, and become non-secreting organs.

Structure of the Mammæ. The mamma is a conglomerate gland, consisting of a number of *lobes*, from 15 to 20 in number, each of which is subdivided into *lobules* made up of *gland vesicles* or *acini*. The ducts which convey the secretion to the exterior, the *lactiferous ducts*, open by 15 to 20 orifices upon the surface of the nipple, at the base of which they are dilated to form little reservoirs in which the milk collects during the periods of active secretion.

The *walls* of the lacteal duct consist of white, fibrous tissue, and non-striated muscular fibres, lined by short columnar cells, which disappear during active lactation. The ducts measure about the $\frac{1}{16}$ of an inch in diameter ; as they pass into the substance of the gland, each duct divides into a number of branches, which are distributed to distinct lobules and terminate in the acini.

An *acinus* is made up of a number of *vesicles* composed of a homogeneous membrane, lined by pavement epithelium. The gland vesicles are held together by white, fibrous tissue, which unites the lobules into lobes.

MILK.

Milk has a pale, blue color, is almost inodorous, of a sweetish taste, an alkaline reaction, and a specific gravity varying from 1.025 to 1.046. Examined microscopically it is seen to contain an immense number of globules, measuring the $\frac{1}{10000}$ of an inch in diameter, suspended in a clear

fluid; these are the *milk globules*, formed of a small mass of oily matter covered by a layer of albumen.

The *quantity* of milk secreted by the human female in 24 hours, during the period of lactation, is about two to three pints; the quantity removed by the infant from a full breast at one time being about two ounces.

COMPOSITION OF MILK.

Water.....	890.00
Proteids, including casein and serum albumen.....	35.00
Fatty matter (butter).....	25.00
Sugar (lactose) with extractives.....	48.00
Salts.....	2.00
	<hr/>
	1000.00

Casein is the nutritive principle of milk, and constitutes its most important ingredient. It is held in solution by an alkali, but upon the addition of an acid it undergoes coagulation, passing into a semi-solid form. The presence of lactic acid, resulting from a transformation of milk sugar, causes spontaneous coagulation to take place.

The *Fatty matter* is more or less solid at ordinary temperature, and consists of margarine and oleine; when subjected to the churning process the globules run together and form a coherent mass, the *butter*.

When milk is allowed to stand for a varying length of time the fat globules rise to the surface, forming a layer more or less thick, the *cream*.

Milk sugar or lactose is an important ingredient in the food of the young child; it is readily transformed into lactic acid in the presence of nitrogenized ferments.

Influences modifying the secretion. During lactation there is a demand for an increased amount of fluid, and if not supplied, the amount of milk secreted is diminished. Good food in sufficient quantity is necessary for the proper elaboration of milk, though no particular article influences its production.

Mental emotion at times influences the character of the milk, decreasing the amount of its different constituents.

Mechanism of Secretion. The water and salts pre-exist in the blood and pass into the gland vesicles by *osmosis*. The casein, fatty matter and sugar appear only in the mammary gland, but the mechanism of their formation is not understood.

Colostrum is a yellowish, opaque fluid, formed in the mammary glands towards the latter period of utero-gestation; it consists of water, albumen, fat, sugar and salts, and acts as a laxative to the newly-born infant.

VASCULAR OR DUCTLESS GLANDS.

The **Vascular Glands** are regarded as possessing the power of acting upon certain elements of the food and aiding the process of sanguinification; of modifying the composition of the blood as it flows through their substance, by some act of secretion.

The vascular glands are the *spleen*, *suprarenal capsules*, *thyroid* and *thymus* glands.

The **Spleen** is about 5 inches in length, 6 ounces in weight, of a dark bluish color, and situated in the left hypochondriac region. It is covered externally by a reflection of the peritoneum, beneath which is the proper fibrous coat, composed of areolar and elastic tissue and non-striated muscular fibres. From the inner surface of the fibrous envelope processes or trabeculae are given off, which penetrate the substance of the gland, forming a network, in the meshes of which is contained the *spleen pulp*. The *splenic artery* divides into a number of branches, some of which, when they become very minute, pass directly into veins, while others terminate in true capillaries.

As the capillary vessels ramify through the substance of the gland, their walls frequently disappear and the blood passes from the arteries into the veins through *lacunæ* (Gray).

The *splenic* or *Malpighian corpuscles* are small bodies, spherical or ovoid in shape, the $\frac{1}{10}$ of an inch in diameter, situated upon the sheaths of the small arteries. They consist of a delicate membrane, containing a semi-fluid substance composed of numerous small cells resembling lymph corpuscles. The *spleen pulp* is a dark red, semi-fluid substance, of a soft consistence, contained in the meshes of the trabeculae. In it are found numerous corpuscles, like those observed in the Malpighian bodies, blood corpuscles in a natural and altered condition, nuclei and pigment granules.

Function of the Spleen. Probably influences the preparation of the albuminous food for nutrition; during digestion the spleen becomes larger, its contents are increased in amount, and after digestion it gradually diminishes in size, returning to the normal condition.

The *red corpuscles* are here disintegrated, after having fulfilled their function in the blood; the splenic venous blood containing relatively a small quantity.

The *white corpuscles* appear to be increased in number, the blood of the splenic vein containing an unusually large proportion.

The *spleen* serves also as a reservoir for blood when the portal circulation becomes obstructed.

The *nervous system* controls the enlargement of the spleen; division of the nerve produces dilatation of the vessels, stimulation contracts them.

The Supra-renal Capsules are triangular, flattened bodies, situated above the kidney. They are invested by a fibrous capsule sending in trabeculae, forming the framework. The glandular tissue is composed of two portions, a *cortical* and *medullary*. The cortical being made up of small cylinders lined by cells and containing an opaque mass, nuclei and granular matter. The medullary consists of a fibrous network containing in the alveoli nucleated protoplasm.

The Thyroid gland consists of a fibrous stroma, containing ovoid closed sacs, measuring on the average $\frac{1}{40}$ of an inch, formed of a delicate membrane lined by cells; the contents of the sacs consist of yellowish albuminous fluid.

The Thymus gland is most developed in early life and almost disappears in the adult. It is divided by processes of fibrous tissue into lobules, and these again into follicles which contain lymphoid corpuscles.

The *functions* of the vascular glands appear to be the more complete elaboration of the blood necessary for proper nutrition; they are most highly developed during infancy and embryonic life, when growth and development are most active.

EXCRETION.

The Principal Excrementitious Fluids discharged from the body are the urine, perspiration and bile; they hold in solution principles of waste which are generated during the activity of the nutritive process, and are the ultimate forms to which the organic constituents are reduced in the body. They also contain inorganic salts.

The Urinary Apparatus consists of the kidneys, ureters and bladder.

KIDNEYS.


The Kidneys are the organs for the excretion of urine; they resemble a bean in shape, are from four to five inches in length, two in breadth, and weigh from four to six ounces.

They are situated in the lumbar region, one on each side of the vertebral column, behind the peritoneum, and extend from the 11th rib to the crest of the ilium; the anterior surface is convex, the posterior concave, and presents a deep notch, the *hilum*.

The kidney is surrounded by a thick layer of fat, beneath which is the

fibrous coat, thin and smooth, composed of dense white fibrous tissue with which are intermingled elastic fibres. It is adherent to the surface of the organ, but can easily be removed by dissection.

FIG. 8.



Longitudinal section through the kidney, the pelvis of the kidney, and a number of renal calyces. A, branch of the renal artery; U, ureter; C, renal calyx; 1, cortex; 1', medullary rays; 1'', labyrinth, or cortex proper; 2, medulla; 2', papillary portion of medulla, or medulla proper; 2'', border layer of the medulla; 3, 3, transverse section through the axes of the tubules of the border layer; 4, fat of the renal sinus; 5, 5, arterial branches; *, transversely coursing medulla rays.—*Tyson, after Henle.*

The Substance of the Kidney is dense, but friable; upon making a longitudinal section, and dividing it, there is presented a cavity, the

pelvis, lined by the proper fibrous coat and occupied by the expanded portion of the *ureter*.

The kidney exhibits two structures, viz.:—

1. *An external or cortical portion*, about $\frac{1}{6}$ of an inch in diameter, of a reddish color, and somewhat granular.

2. *An internal or medullary portion*, of a dark red color, arranged in the form of *pyramids*, the bases of which are directed toward the cortical portion, and the apices toward the *pelvis*, into which they project, and are covered by the *calyces*.

The **Cortical portion** of the kidney consists of a delicate matrix containing an immense number of tubules, having a markedly convoluted appearance, and interlacing in every direction (the tubules of Ferrein). Throughout its structure are numerous ovoid bodies, the *Malpighian bodies*, which are the flask-like terminations of the convoluted tubules;

FIG. 9.



Diagrammatic exposition of the method in which the uriniferous tubes unite to form primitive cones.—*Tyson, after Ludwig.*

these tubes are composed of a delicate homogeneous membrane lined by nucleated cells. After pursuing a most intricate course in the cortical portion, they become narrower and form loops which dip into the pyramidal portion (Henle's tubules), returning upon themselves, to finally terminate in the straight tubes of the pyramids.

The **Malpighian bodies**, the dilated extremities of the convoluted tubes, consist of a little sac (the capsule of Müller), which is ovoid in shape, measuring about the $\frac{1}{200}$ of an inch in diameter, and contains a tufted mass of minute blood vessels, over the surface of which is reflected a layer of cells.

Medullary Substance. The conical masses, the *pyramids of Malpighi*, consist of a number of straight tubes, which commence at the apex by from 10 to 20 openings; and as they pass toward the cortical portion, they divide and subdivide at acute angles, until a large mass of tubes is

produced. These tubes are on the average about $\frac{1}{500}$ of an inch in diameter, and composed of a thin, but firm, elastic, structureless membrane, lined by polygonal nucleated cells, which reduce the diameter of the lumen of the tube about two-thirds; these are the *straight tubes* of Bellini.

Blood vessels of the Kidney. *The renal artery* is of large size and enters the organ at the hilum; it divides into several large branches, which penetrate the substance of the kidney, between the pyramids, at the base of which they form an anastomosing plexus, which completely surrounds them. From this plexus vessels follow the straight tubes toward the apex, while others entering the cortical portion, divide into small twigs which enter the Malpighian body and form a mass of convoluted vessels, the *glomerulus*. After circulating through the Malpighian tuft the blood is gathered together by two or three small veins, which again subdivide and form a fine capillary plexus, which envelops the convoluted tubules; from this plexus the veins converge to form the emulgent vein, which empties into the vena cava.

The nerves of the kidney follow the course of the blood vessels and are derived from the renal plexus.

The Ureter is a membranous tube, situated behind the peritoneum, about the diameter of a goose quill, 18 inches in length, and extends from the pelvis of the kidney to the base of the bladder, which it perforates in an oblique direction. It is composed of 3 coats, fibrous, muscular and mucous.

The Bladder is a temporary reservoir for the reception of the urine prior to its expulsion from the body; when fully distended it is ovoid in shape, and holds about *one pint*. It is composed of four coats, *serous*, *muscular*, the fibres of which are arranged longitudinally and circularly, *areolar* and *mucous*. The orifice of the bladder is controlled by the *sphincter vesicæ*, a muscular band, about half an inch in width.

As soon as the urine is formed it passes through the *tubuli uriniferi* into the pelvis, and from thence through the ureters into the bladder, which it enters at an irregular rate. Shortly after a meal, after the ingestion of large quantities of fluid, and after exercise, the urine flows into the bladder quite rapidly, while it is reduced to a few drops during the intervals of digestion. It is prevented from regurgitating into the ureters on account of the oblique direction they take between the mucous and muscular coats.

Nervous Mechanism of Urination. When the urine has passed into the bladder it is there retained by the sphincter vesicæ muscle kept in a state of tonic contraction by the action of a nerve centre in the lumbar region of the spinal cord. This centre can be inhibited and the sphincter relaxed, either *reflexly*, by impressions coming through sensory nerves from the mucous membrane of the bladder, or *directly*, by a voluntary impulse descending the spinal cord. When the desire to urinate is experienced, impressions made upon the vesical sensory nerves are carried to the centres governing the *sphincter* and *detrusor urinæ* muscles and to the brain. If now the act of urination is to take place, a voluntary impulse, originating in the brain, passes down the spinal cord and still further inhibits the sphincter vesicæ centre, with the effect of relaxing the muscle, and of stimulating the centre governing the detrusor muscle, with the effect of contracting the muscle and expelling the urine. If the act is to be suppressed voluntary impulses inhibit the *detrusor* centre and possibly stimulate the *sphincter* centre.

The *genito-spinal* centre controlling these movements is situated in that portion of the spinal cord corresponding to the origin of the 3d, 4th and 5th sacral nerves.

URINE.

Normal Urine is of a pale yellow or amber color, perfectly transparent, with an aromatic odor, an acid reaction, a specific gravity of 1.020, and a temperature when first discharged of 100° Fahr.

The *color* varies considerably in health, from a pale yellow to a brown hue, due to the presence of the coloring matter, *urobilin* or *urochrome*.

The *transparency* is diminished by the presence of mucus, the calcium and magnesium phosphates and the mixed urates.

The *reaction* is slightly acid, caused by the acid phosphate of sodium. After standing for a short time, an increased acidity is observed, due to an acid fermentation, from the presence of mucus. The urea is converted into ammonium carbonate, giving rise to a strong ammoniacal odor.

The *specific gravity* varies from 1.010 to 1.025.

The *quantity* of urine excreted in 24 hours is between 40 and 50 fluid ounces, but ranges above and below this standard.

The *odor* is characteristic, and caused by the presence of taurylic and phenylic acids, but is influenced by vegetable foods and other substances eliminated by the kidneys.

COMPOSITION OF URINE.

COMPOSITION OF URINE.	
Water.....	967.
Urea.....	14.230
Other nitrogenized crystalline bodies, uric acid, principally in the form of alkaline urates.	10.635
Creatin, creatinin, xanthin, hypoxanthin.	
Hippuric acid, leucin, tyrosin, taurin, cystin, all in small amounts, and not constant.	
Mucus and pigment.	
Salts:—	
<i>Inorganic</i> , principally sodium and potassium sulphates, phosphates and chlorides, with magnesium and calcium phosphates, traces of silicates and chlorides.	8.135
<i>Organic</i> : lactates, hippurates, acetates, formates, which appear only occasionally.	
Sugar.....	a trace.
Gases (nitrogen and carbonic acid principally).	
<hr/>	
1000.00	

The Average Quantity of the principal constituents excreted in 24 hours is as follows:—

Water.....	52 fluid oz.
Urea.....	512.4 grains.
Uric acid.....	8.5 “
Phosphoric acid.....	45.0 “
Sulphuric acid.....	31.11 “
Inorganic salts.....	323.25 “
Lime and magnesia.....	6.5 “

To Determine the amount of solid matters in any given amount of urine, multiply the last two figures of the specific gravity by the coefficient of Haeser, 2.33; *e.g.*, in 1000 grains of urine having a specific gravity 1.022, there are contained $22 \times 2.33 = 51.26$ grains of solid matter.

The Elimination of the urinary constituents is accomplished by the two processes of filtration and secretion.

1. By **Filtration** the water and mineral salts are removed from the blood, and takes place, for the most part, in the Malpighian corpuscles, by the process of osmosis. The *amount* of these constituents eliminated varies with the pressure of blood in the renal arteries. All of the agencies which increase the general blood pressure increase the quantity of urine.

Season. In summer, while the capillary vessels of the skin are dilated,

and perspiration is abundant, there is a diminished blood pressure, and a consequent diminution in the amount of urine; in winter the reverse takes place.

During sleep the renal excretion is diminished, but increased in the morning hours, and especially after the ingestion of hearty meals.

The nervous system influences the secretion of urine. Irritation of the medulla oblongata, a little above the origin of the pneumogastric and auditory nerves, increases the quantity; division of the renal nerves destroys the nutrition of the kidney, and thus interferes with the elimination of the urine. Mental emotion, fear, anxiety, etc., increase the amount secreted.

2. Secretion. While it is established that the Malpighian corpuscles permit the filtration of water and salts, it has also been shown that the renal epithelial cells lining the convoluted tubes are the agencies by which the solid matters, *urea*, *creatin*, etc., are removed from the blood, by a process of *true secretion*, which is independent of blood pressure and caused by the pressure of these ingredients in the blood.

Urea is the most important of the organic constituents of the urine. It is a colorless, neutral substance, crystallizing in four-sided prisms, soluble in boiling alcohol and water; when subjected to prolonged boiling it is decomposed, with the production of ammonium carbonate.

Urea is not formed in the kidneys, but pre-exists in the blood.

The Amount of Urea excreted in 24 hours is estimated at about 500 grains; it is *increased* during the waking hours, by an animal diet, and by prolonged muscular exertion; *diminished* during sleep and by non-nitrogenized food.

Source. Urea results from an imperfect oxidation of the albuminous principles of the food, and from a disintegration of the organic constituents of the tissues.

Uric acid, or *lithic acid*, is a constant ingredient of the urine; the amount excreted daily is about 8 grains; it is increased by nitrogenized, decreased by non-nitrogenized food. It exists in the urine in a free state, and as the urate of soda. It arises from the disassimilation of albuminous compounds, and when secreted in excess is deposited in a crystalline form, as a brown or "brick-red" sediment, with the sodium and ammonium urates.

Creatin is a colorless, transparent substance, crystallizing in prisms; found in blood, kidneys, and muscular tissue; by boiling in acid solutions it is transformed into

Creatinin, which resembles creatin chemically. It is soluble in water and alcohol, and crystallizes in colorless prisms. About 15 grains are excreted daily.

The *Earthy phosphates* are insoluble in water but held in solution in the urine by the acid reaction. If the urine becomes alkaline, they are deposited copiously, and yet may not be increased in quantity; from 15 to 25 grains are excreted in 24 hours. The *sulphates* are those of sodium and potassium; they are very soluble and do not appear as a precipitate; the average quantity excreted in 24 hours is about 60 grains.

Abnormal ingredients appear in the urine at times, in pathological conditions, *e. g.*, sugar, albumen, biliary salts, etc.

The *Gases* of the urine are carbonic acid and nitrogen.

LIVER.

The **Liver** is a highly vascular, conglomerate gland, appended to the alimentary canal, and performs the triple office of (1) *excreting bile*, (2) *elaborating blood* and (3) *secreting glycogen*.

It is the largest gland in the body, weighing about $4\frac{1}{2}$ pounds; it is situated in the right hypochondriac region, and retained in position by five ligaments, four of which are formed by duplicatures of the peritoneal investment.

The *proper coat* of the liver is a thin but firm fibrous membrane, closely adherent to the surface of the organ, which it penetrates at the transverse fissure, and follows the vessels in their ramifications through its substance, constituting *Glisson's capsule*.

Structure of the Liver. The liver is made up of a large number of small bodies, the *lobules*, rounded or ovoid in shape, measuring the $\frac{1}{25}$ of an inch in diameter, separated by a space in which are situated blood vessels, nerves, hepatic ducts and lymphatics.

The *lobules* are composed of cells, which, when examined microscopically, exhibit a rounded or polygonal shape, and measure, on the average, the $\frac{1}{1000}$ of an inch in diameter; they possess one, and at times two, nuclei; they also contain globules of fat, pigment matter, and animal starch. The cells constitute the *secreting* structure of the liver, and are the *true hepatic cells*.

The **Blood vessels** which enter the liver are (1) The *portal vein*, made up of the *gastric, splenic, superior* and *inferior mesenteric veins*; (2) the *hepatic artery*, a branch of the coeliac axis; both of which are invested by a sheath of areolar tissue; the vessels which leave the liver

are the *hepatic veins*, originating in its interior, collecting the blood distributed by the portal vein and hepatic artery, and conducting it to the *ascending vena cava*.

Distribution of Vessels. The *portal vein* and *hepatic artery*, upon entering the liver, penetrate its substance, divide into smaller and smaller branches, occupy the spaces between the lobules, completely surrounding and limiting them, and constitute the *inter-lobular vessels*. The *hepatic artery*, in its course, gives off branches to the walls of the portal vein and Glisson's capsule, and finally empties into the small branches of the *portal vein* in the interlobular spaces.

The *interlobular vessels* form a rich plexus around the lobules, from which branches pass to neighboring lobules and enter their substance, where they form a very fine network of capillary vessels, ramifying over the hepatic cells, in which the various functions of the liver are performed. The blood is then collected by small veins, converging toward the centre of the lobule, to form the *intra-lobular vein*, which runs through its long axis and empties into the *sub-lobular vein*. The *hepatic veins* are formed by the union of the sub-lobular veins, and carry the blood to the ascending vena cava; their walls are thin and adherent to the substance of the hepatic tissue.

The Hepatic Ducts or Bile Capillaries originate within the lobules, in a very fine plexus lying between the hepatic cells; whether the smallest vessels have distinct membranous walls, or whether they originate in the spaces between the cells by open orifices, has not been satisfactorily determined.

The *Bile Channels* empty into the interlobular ducts, which measure about $\frac{1}{2000}$ of an inch in diameter, and are composed of a thin homogeneous membrane lined by flattened epithelial cells.

As the interlobular bile ducts unite to form larger trunks, they receive an external coat of fibrous tissue, which strengthens their walls; they finally unite to form one large duct, the *hepatic duct*, which joins the *cystic duct*; the union of the two forms the *ductus communis choledochus*, which is about three inches in length, the size of a goose quill, and opens into the duodenum.

The Gall Bladder is a pear-shaped sack, about four inches in length, situated in a fossa on the under surface of the liver. It is a reservoir for the bile, and is capable of holding about one ounce and a half of fluid. It is composed of three coats, (1) serous, a reflection of the peritoneum, (2) fibrous and muscular, (3) mucous.

(1) **Bile. Mechanism of its Secretion.** Bile does not preëxist in the blood, but is formed in the interior of the hepatic cells, from materials derived from the venous as well as arterial blood. The secreted bile is then taken up by the delicate plexus of vessels, from which it passes into the larger ducts, and finally either empties into the intestine or is regurgitated backward into the gall bladder, in which it is stored up during the intervals of digestion.

Although the secretion of bile is constantly taking place, it is only when the food passes into the intestinal canal that this fluid is discharged abundantly, under the influence of the contraction of the walls of the gall bladder; it increases in amount during the period of active digestion, from the 2d to the 8th hour, and then gradually diminishes.

The Bile is both a *secretion* and an *excretion*; it contains new constituents which are formed only in the substance of the liver, and are destined to play an important part ultimately in nutrition; it contains also waste ingredients which are discharged into the intestinal canal and eliminated from the body.

The physical properties and functions of bile have been considered under the head of digestion (see page 32).

(2) **Elaboration of Blood.** Besides the capability of secreting bile, the liver possesses the property of so acting upon and modifying the chemical composition of the products of digestion, as they traverse its substance, that they readily assimilate with the blood, and are transformed into materials capable of being converted into the elements of the blood and solid tissues.


The *albuminose* particularly requires the modifying influence of the liver; for if it be removed from the portal vein and introduced into the jugular vein, it is at once removed from the blood by the action of the kidneys.

The blood of the *hepatic vein* differs from the blood of the *portal vein*, in being richer in blood corpuscles, both red and white; its plasma is more dense, containing a less percentage of water and a greater amount of solid constituents, but no fibrin; its serum contains less albumen, fat and salts, but its sugar is increased.

(3) **Glycogenic Function.** In addition to the two preceding functions, Bernard, in 1848, demonstrated the fact that the liver, during life, normally produces a sugar-forming substance, analogous in its chemical composition to starch, which he termed glycogen; also that when the liver is removed from the body, and its blood vessels thoroughly washed out, after a few hours, sugar again makes its appearance, in abundance.

It can be shown to exist in the blood of the hepatic vein as well as in a decoction of the liver substance, by means of either Trommer's or Fehling's tests, even when the blood of the portal vein does not contain a trace of sugar.

Origin and Destination of Glycogen. Glycogen appears to be formed *de novo* in the liver cells, from materials derived from the food, whether the diet be animal or vegetable, though a larger per cent. is formed when the animal is fed on starchy and saccharine, than when fed on animal food. The *glucose*, which is one of the products of digestion, is absorbed by the blood vessels, and carried directly into the liver; as it does not appear in the urine, as it would if injected at once into the general circulation, it is probable that it is detained in the liver, dehydrated and stored up as *glycogen*. The change is shown by the following formula:—



The glycogen thus formed is stored up in the hepatic cells for the future requirements of the system. When it is carried from the liver it is again transformed into *glucose* by the agency of a ferment. Glycogen does not undergo oxidation in the blood; this takes place in the tissues, particularly in the muscles, where it generates heat and contributes to the development of muscular force.

Glycogen, when obtained from the liver, is an amorphous, starch-like substance, of a white color, tasteless and odorless, and soluble in water; by boiling with dilute acids, or subjected to the action of an animal ferment, it is easily converted into *glucose*. When an excess of sugar is generated by the liver, it can be found, not only in the blood of the hepatic vein, but also in other portions of the body; under these circumstances it is eliminated by the kidneys, appearing in the urine, constituting the condition of *glycosuria*.

The nervous system influences the production of the glycogenic matter; irritation of the medulla oblongata, between the auditory and pneumogastric nerves, is followed by an increase in the production of sugar, and its appearance in the urine, which, however, is only temporary.

SKIN.

The Skin, the external investment of the body, is a most complex and important structure, serving (1) as a *protective covering*; (2) an organ for *tactile sensibility*; (3) an organ for the *elimination of excrementitious matters*.

The Amount of Skin investing the body of a man of average size is about twenty feet, and varies in thickness, in different situations, from the $\frac{1}{8}$ to the $\frac{1}{100}$ of an inch.

The skin consists of two principal layers, viz., a deeper portion, the *Corium*, and a superficial portion, the *Epidermis*.

The Corium, or Cutis Vera, may be subdivided into a *reticulated* and a *papillary* layer. The *former* is composed of white fibrous tissue, non-striated muscular fibres and elastic tissue, interwoven in every direction, forming an areolar network, in the meshes of which are deposited masses of fat, and a structureless amorphous matter; the *latter* is formed mainly of club-shaped elevations or projections of the amorphous matter, constituting the *papillæ*; they are most abundant, and well developed, upon the palms of the hands and the soles of the feet; they average the $\frac{1}{100}$ of an inch in length, and may be simple or compound; they are well supplied with nerves, blood vessels and lymphatics.

The Epidermis or scarf skin is an extra-vascular structure, a product of the true skin, and composed of several layers of cells. It may be divided into two layers, the *rete mucosum* or the *Malpighian layer*, and the *horny or corneous*.

The *former* closely applies itself to the papillary layer of the true skin, and is composed of large, nucleated cells, the lowest layer of which, the "prickle cells," contain pigment granules, which give to the skin its varying tints in different individuals and in different races of men; the more superficial cells are large, colorless, and semi-transparent. The *latter*, the corneous layer, is composed of flattened cells, which, from their exposure to the atmosphere, are hard and horny in texture; it varies in thickness from $\frac{1}{8}$ of an inch on the palms of the hands and feet, to the $\frac{1}{100}$ of an inch in the external auditory canal.

APPENDAGES OF THE SKIN.

Hairs are found in almost all portions of the body, and can be divided into (1) long, soft hairs, on the head; (2) short, stiff hairs, along the edges of the eyelids and nostrils; (3) soft, downy hairs, on the general cutaneous surface. They consist of a *root* and a *shaft*, which is oval in shape, and about the $\frac{1}{100}$ of an inch in diameter; it consists of fibrous tissue, covered externally by a layer of imbricated cells, and internally by cells containing granular and pigment material.

The *Root* of the hair is embedded in the hair follicle, formed by a tubular depression of the skin, extending nearly through to the subcutaneous tissue;

its walls are formed by the layers of the corium, covered by epidermic cells. At the bottom of the follicle is a papillary projection of amorphous matter, corresponding to a papilla of the true skin, containing blood vessels and nerves, upon which the hair root rests. The investments of the hair roots are formed of epithelial cells, constituting the *internal* and *external* root sheaths.

The hair protects the head from the heat of the sun and cold, retains the heat of the body, prevents the entrance of foreign matter into the lungs, nose, ears, etc. The *color* is due to the pigment matter, which, in old age, becomes more or less whitened.

The Sebaceous Glands, imbedded in the true skin, are simple and compound racemose glands, opening, by a common excretory duct, upon the surface of the epidermis or into the hair follicle. They are found in all portions of the body, most abundantly in the face, and are formed by a delicate, structureless membrane, lined by flattened polyhedral cells. The sebaceous glands secrete a peculiar oily matter, the *sebum*, by which the skin is lubricated and the hairs softened; it is quite abundant in the region of the nose and forehead, which often present a greasy, glistening appearance; it consists of water, mineral salts, fatty globules, and epithelial cells.

The *Vernix caseosa* which frequently covers the surface of the foetus at birth consists of the residue of the sebaceous matters, containing epithelial cells and fatty matters; it seems to keep the skin soft and supple, and guards it from the effects of the long continued action of water.

The Sudoriparous Glands excrete the sweat; they consist of a mass or coil of a tubular gland duct, situated in the derma and in the subcutaneous tissue; average the $\frac{1}{8}$ of an inch in diameter, and are surrounded by a rich plexus of capillary blood vessels. From this coil the duct passes in a straight direction up through the skin to the epidermis, where it makes a few spiral turns and opens obliquely upon the surface. The sweat glands consist of a delicate homogeneous membrane lined by epithelial cells, whose function is to extract from the blood the elements existing in the perspiration.

The glands are very abundant all over the cutaneous surface, as many as 3528 to the square inch, according to Erasmus Wilson.

The Perspiration is an excrementitious fluid, clear, colorless, almost odorless, slightly acid in reaction, with a specific gravity of 1.003 or 1.004.

The total quantity of perspiration excreted daily has been estimated at about two pounds, though the amount varies with the nature of the food and drink, exercise, external temperature, season, etc.

The elimination of the sweat is not intermittent, but continuous; but it takes place so gradually that as fast as it is formed it passes off by evaporation as *insensible* perspiration. Under exposure to great heat and exercise the evaporation is not sufficiently rapid, and it appears as *sensible* perspiration.

COMPOSITION OF SWEAT.

Water.....	995.573
Urea.....	0.043
Fatty matters.....	0.014
Alkaline lactates.....	0.317
Alkaline sudorates.....	1.562
Inorganic salts.....	2.491
	<hr/>
	1000.000

Urea is a constant ingredient.

Carbonic acid is also exhaled from the skin, the amount being about $\frac{1}{100}$ of that from the lungs.

Perspiration regulates the temperature, and removes waste matters from the blood; it is so important, that if elimination be prevented death occurs in a short time.

The Nervous System influences the secretion of watery vapor by causing a dilatation of the capillary blood vessels around the tubular coil. It is increased by mental emotions; section of the sympathetic fibres in the neck is followed by a copious perspiration; stimulation of the nerves, producing contraction of the vessels, is followed by an arrestation of the elimination of the sweat.

NERVOUS SYSTEM.

The Nervous System co-ordinates all the various organs and tissues of the body, and brings the individual into conscious relationship with external nature by means of sensation, motion, language, mental and moral manifestations.

The Nervous Tissue may be divided into two systems, viz: the *Cerebro-spinal* and the *Sympathetic*.

(1) **The Cerebro-spinal System** occupies the cavities of the cranium and spinal canal, and consists of the brain, the spinal cord, the cranial and spinal nerves. It is the system of animal life, and presides over the functions of sensation, motion, etc.

(2) **The Sympathetic System**, situated along each side of the spinal column, consists (1) of a double chain of ganglia, united together by nerve cords, which extends from the base of the cranium to the coccyx; (2) of various ganglia, situated in the head and face, thorax, abdomen, pelvis, etc. All the ganglia are united together by numerous communicating fibres, many of which anastomose with the fibres of the cerebro-spinal system. It is the nervous system of organic life, and governs the functions of nutrition, growth, etc.

Nervous Tissue is composed of two kinds of matter, the *gray* and *white*, which differ in their color, structure and physiological endowments; the former consists of *vesicles* or *cells* which receive and generate nerve force; the latter consists of *fibres* which simply conduct it, either from the periphery to the centre or the reverse.

Structure of Gray Matter. The gray matter, found on the surface of the brain in the convolutions, in the interior of the spinal cord, and in the various ganglia of the cerebro-spinal and sympathetic nervous systems, consists of a fine connective-tissue stroma, the *neuroglia*, in the meshes of which are embedded the *gray cells* or *vesicles*.

The *cells* are grayish in color, and consist of a delicate investing capsule containing a soft, granular, albuminous matter, a *nucleus*, and sometimes a *nucleolus*. Some of the cells are spherical or oval in shape, while others have an interrupted outline, on account of having one, two, or more processes issuing from them, constituting the *uni-polar*, *bi-polar* or *multi-polar* nerve cells. Cells vary in size; the smallest being found in the brain, the largest in the anterior horns of gray matter of the spinal cord. Some of

the cell processes become continuous with the fibres of the white matter, while others anastomose with those of adjoining cells and form a *plexus*.

Structure of the White Matter. The white matter, found for the most part in the interior of the brain, on the surface of the spinal cord, and in almost all the nerves of the cerebro-spinal and sympathetic systems, consists of minute tubules or fibres, the *ultimate nerve filaments*, which in the perfectly fresh condition, are apparently structureless and homogeneous; but when carefully examined after death are seen to consist of three distinct portions, (1) a tubular membrane; (2) the white substance of Schwann; (3) the axis cylinder.

The *Tubular membrane*, investing the nerve filament, is thin, homogeneous, and lined by large, oval nuclei, and presents, in its course, annular constrictions; it serves to keep the internal parts of the fibre in position, and protects them from injury.

The *White substance* of Schwann, or the *medullary layer*, is situated immediately within the tubular membrane, and gives to the nerves their peculiar white and glistening appearance. It is composed of oleaginous matter in a more or less fluid condition; after death it undergoes coagulation, giving to the fibre a knotted or varicose appearance. It serves to insulate the axis cylinder, and prevents the diffusion of the nerve force.

The *Axis cylinder* occupies the centre of the medullary substance. In the natural condition it is transparent and invisible, but when treated with proper reagents, it presents itself as a pale, granular, flattened band, albuminous in character, more or less solid, and somewhat elastic. It is composed of a number of minute fibrillæ united together to form a single bundle. (Schultze.)

Nerve fibres in which these three structural elements coexist are known as the *medullated* nerve fibres. In the sympathetic system, and in the gray substance of the cerebro-spinal system, many nerves are destitute of a medullary layer, and are known as the *non-medullated* nerve fibres.

Gray or Gelatinous nerve fibres, found principally in the sympathetic system, are gray in color, semi-transparent, flattened, with distinct borders, finely granular, and present oval nuclei.

The diameter of the gelatinous fibres is about the $\frac{1}{8000}$ of an inch; of the medullated fibres, from $\frac{1}{2500}$ to $\frac{1}{1500}$ of an inch.

Ganglia are small bodies, varying considerably in size, situated on the posterior roots of spinal nerves, on the sensory cranial nerves, alongside of the vertebral column, forming a connected chain, and in the different viscera. They consist of a dense, investing, fibrous membrane, containing

in its interior *gray* or *vesicular* cells, among which are found *white* and *gelatinous* nerve fibres. They may be regarded as independent nerve centres.

Structures of Nerves. Nerves are rounded or flattened cords extending from the centres to the periphery; they are surrounded externally by a sheath, the *neurilemma*, composed of fibrous and elastic tissue forming a stroma, in which blood vessels ramify, from which the nerves derive their nourishment.

A *Nerve* consists of a greater or less number of *ultimate nerve filaments*, separated into bundles by fibrous septa given off from the neurilemma. The nerve filaments pursue an uninterrupted course, from their origin to their termination; branches pass from one nerve trunk into the sheath of another, but there is no anastomosis or coalescence with adjoining nerve fibres.

A **Plexus** is formed by a number of branches of different nerves interlacing in every direction, in the most intricate manner, but from which fibres are again given off to pursue their independent course, *e. g.*, brachial, cervical, lumbar, sacral, cardiac plexuses, etc.

SPINAL NERVES.

Origin. The spinal nerves are thirty-one in number on each side of the spinal cord, and arise by two roots, an *anterior* and *posterior*, from the anterior and posterior aspects of the cord respectively; the *posterior roots* present near their emergence from the cord a small ganglionic enlargement; outside of the spinal canal the two roots unite to form a main trunk, which is ultimately distributed to the skin, muscles and viscera.

The Function of the Anterior Roots is to transmit *motor* impulses from the centres outward to the periphery. *Irritation* of these roots, from whatever cause, excites convulsive movements in the muscles to which they are distributed; *disease* or *division* of these roots induces a condition of paresis or paralysis.

The Function of the Posterior Roots is to transmit the impressions made upon the periphery to the centres in the spinal cord, where they excite motor impulses, or to the brain, in which they are translated into conscious sensations. *Irritation* of these roots gives rise to painful sensations; *division* of the roots abolishes all sensation in the parts to which they are distributed.

The *ganglion* on the posterior root influences the nutrition of the sensory nerve; for if the nerve be separated from the ganglion, it undergoes degeneration in the course of a few days, in the direction in which it carries impressions, *i. e.*, from the periphery to the centres; if the nerve be

divided between the ganglion and the cord, the central end only undergoes degeneration. The nutrition of the *anterior root* is governed by nerve cells in the gray matter of the cord; for if these cells undergo *atrophy*, or if the nerve be *divided*, it undergoes degeneration outward.

Nerve Terminations. (1) *Central.* Both motor and sensory nerve fibres, as they enter the spinal cord and brain, lose their external investments, and retaining only the axis cylinder, ultimately become connected with the processes of the gray cells.

(2) *Peripheral.* As the nerves approach the tissues to which they are to be distributed, they inosculate freely, forming a *plexus* from which the ultimate fibres proceed to individual tissues.

Motor Nerves. In the *voluntary* or *striped* muscles the motor nerves are connected with the contractile substance by means of the "*motorial end plates*;" when the nerve enters the muscular fibre the tubular membrane blends with the sarcolemma, the medullary layer disappears, and the axis cylinder spreads out into the form of a little plate, granular in character, and containing oval nuclei.

In the *unstriped* or *involuntary* muscles, the terminal nerve fibres form a plexus on the muscular fibre cells, and become connected with the granular contents of the nuclei.

In the *glands* nerve fibres have been traced to the glandular cells, where they form a branching plexus from which fibres pass into their interior and become connected with their substance, and thus influence secretion.

Sensitive Nerves terminate in the skin and mucous membranes, in three distinct modes, *e. g.*, as tactile corpuscles, Pacinian corpuscles, and as end bulbs.

The *tactile corpuscles* are found in the papillæ of the true skin, especially on the palmar surface of the hands and fingers, feet and toes; they are oblong bodies, measuring about $\frac{1}{80}$ of an inch in length, consisting of a central bulb of homogeneous connective tissue surrounded by elastic fibres and elongated nuclei. The nerve fibre approaches the base of the corpuscle, makes two or three spiral turns around it, and terminates in loops. They are connected with the sense of touch.

The *Pacinian corpuscles* are found chiefly in the subcutaneous cellular tissue, on the nerves of the hands and feet, the intercostal nerves, the cutaneous nerves, and in many other situations. They are oval in shape, measure about the $\frac{1}{10}$ of an inch in length on the average, and consist of concentric layers of connective tissue; the nerve fibre penetrates the corpuscle and terminates in a rounded knob in the central bulb. Their *function* is unknown.

The *end bulbs* of Krause are formed of a capsule of connective tissue in which the nerve fibre terminates in a coiled mass or bulbous extremity; they exist in the conjunctiva, tongue, glans penis, clitoris, etc.

Many *sensitive* nerves terminate in the papillæ at the base of the hair follicle; but in the skin, mucous membranes, and organs of special sense their mode of termination is not well understood.

PROPERTIES AND FUNCTIONS OF NERVES.

Classification. Nerves may be divided into two groups, viz.:—

(1) *Afferent* or *centripetal*, as when they convey to the nerve centres the impressions which are made upon their peripheral extremities or parts of their course. They may be *sensitive*, when they transmit impressions which give rise to sensations; *reflective* or *excitant*, when the impression carried to the nerve centre is reflected outward by an efferent nerve and produces motion or some other effect in the part to which the nerve is distributed.

(2) *Efferent* or *centrifugal*, as when the impulses generated in the centres are transmitted outward to the muscles and various organs. They may be *motor*, as when they convey impulses to the voluntary and involuntary muscles; *vaso-motor*, when they regulate the calibre of the small blood vessels, increasing or diminishing the amount of blood to a part; *secretory*, when they influence secretion; *trophic*, when they influence nutrition; *inhibitory*, when they conduct impulses which produce a restraining or inhibiting action.

The *Axis Cylinder* is the essential conducting agent, the white substance of Schwann and tubular membrane being probably accessory structures, protecting the axis from injury, and preventing the diffusion of nerve force to adjoining nerves.

The properties of *sensation* and *motion* reside in different nerve fibres. *Motor* nerves can be destroyed or paralyzed by the introduction of woorara under the skin, without affecting sensation; the *sensibility* of nerves can be abolished by the employment of anæsthetics without destroying motion.

Irritability. Nerves conduct peripheral impressions to the centres, and motor impulses to the periphery, in virtue of their possessing an ultimate and inherent property, denominated *neurility*, *nervous irritability*, or *excitability*, which is manifested as long as the physical and chemical integrity of the nerve is maintained.

Nerve degeneration. When nerves are separated from their trophic or nutritive centres, they degenerate progressively in the direction in which they conduct impressions. In *motor* nerves, from the centre to the periphery; in sensory nerves, from the periphery to the centres.

Nerve-force is not identical with electricity. Nerves do not possess the power of generating force, or of originating impulses within themselves, but propagate only the nervous impulses which are called forth by chemical, physical and mechanical stimuli from without, and by volitional acts, normal and pathological conditions from within.

Phenomena of Muscles and Nerves. The muscles are the motor organs of the body and constitute a large per cent. of the body weight. Muscles are of two kinds, striated and non-striated or involuntary. The striated muscles consist of bundles of fibres, the *fasciculi*, held together by connective tissue. Each muscle fibre is about $\frac{1}{2}$ to $1\frac{1}{2}$ inches long, and possesses a delicate homogeneous membrane, the *sarcolemma*, in the interior of which is contained the contractile substance, which presents a striated appearance. During life this substance is in a fluid condition, but after death undergoes stiffening.

The non-striated muscles form membranes which surround cavities, *e. g.*, stomach, arteries, bladder, etc. They are composed of elongated cells without striations and contain in their interior one or more nuclei.

Muscular tissue is composed of water, an organic contractile substance, *myosin*, non-nitrogenized substances, such as glycogen, inosite, fat, and inorganic salts. When at rest the muscle is alkaline in reaction, but during and after contraction it becomes acid.

Muscles possess the properties of (1) *Contractility*, which is the capability of shortening themselves in the direction of their long axis, and at the same time becoming thicker and more rigid. (2) *Extensibility*, by means of which they are lengthened in proportion to weights attached. (3) *Elasticity*, in virtue of which they return to their original shape when the force applied is removed.

The contractility of muscles is called forth mainly by nervous impulses, descending motor nerves, which originate in the central nervous system; but it can also be excited by the electric current, the application of strong acids, heat, or by mechanical means.

Phenomena of a Muscular Contraction. When a single induction shock is propagated through a nerve, the muscle to which it is distributed undergoes a quick pulsation, and speedily returns to its former condition. As is shown by the muscle curve, the contraction, which is at first slow, increases in rapidity to its maximum, gradually relaxes and is again at rest, the entire pulsation not occupying more than the $\frac{1}{10}$ of a second.

The muscular contraction does not instantly follow the induction shock, even when the electrodes are placed directly upon the muscular fibres themselves; an appreciable period intervenes before the contraction,

during which certain chemical changes are taking place preparatory to the manifestation of force. This is the "latent period," which has an average duration of the $\frac{1}{100}$ of a second, but varies with the temperature, the strength of the stimulus, the animal, etc. The muscular movements of the body, however, are occasioned by contractions of a much longer duration, depending upon the number (the average, 20) of nervous impulses passing to the muscles in a second.

During the muscular contraction the following phenomena are observed, viz: a change in form, a rise in temperature, a consumption of oxygen and an evolution of carbonic acid; the production of a distinct musical sound, a change from an alkaline to an acid reaction, from the development of sarcolactic acid; a disappearance of the natural muscle currents, which undergo a *negative variation* in the "latent period," just after the nervous impulse reaches the termination of the nerve, and before the appearance of the muscular contraction wave.

Electrical Properties of Nerves. When a galvanic current is made to flow along a motor nerve from the centre to the periphery, from the positive to the negative pole, it is known as the *direct, descending* or *centrifugal* current. When it is made to flow in the reverse direction it is known as the *inverse, ascending* or *centripetal* current.

The passage of a direct current enfeebls the excitability of a nerve; the passage of the inverse current increases it. The excitability of a nerve may be exhausted by the repeated applications of electricity; when thus exhausted it may be restored by repose, or by the passage of the inverse current if the nerve has been exhausted by the direct current or vice versa.

During the actual passage of a feeble constant current in either direction neither pain nor muscular contraction is ordinarily manifested; if the current be very intense the nerve may be disorganized and its excitability destroyed.

Electrotonus. The passage of a direct galvanic current through a portion of a nerve excites in the parts beyond the electrodes a condition of electric tension or *electrotonus*, during which the *excitability* of the nerve is *decreased* near the anode or positive pole, and *increased* near the kathode or negative pole; the increase of excitability in the *kathoelectrotonic area*, that nearest the muscle, being manifested by a more marked contraction of the muscle than the normal, when the nerve is irritated in this region. The passage of an *inverse* galvanic current excites the same condition of electrotonus; and the *diminution* of excitability near the anode, the *anelectrotonic area*, that now nearest the muscle, being manifested by a less marked contraction than the normal when the nerve is stimulated in

this region. Between the electrodes is a neutral point where the kath-electrotonic area emerges into the anelectrotonic area. If the current be a strong one, the neutral point approaches the kathode; if weak, it approaches the anode.

When a nervous impulse passes along a nerve, the only appreciable effect is a change in its electrical condition, there being no change in its temperature, chemical composition or physical condition. The natural nerve currents, which are always present in a living nerve as a result of its nutritive activity, in great part disappear during the passage of an impulse, undergoing a *negative variation*.

The *rapidity* with which nervous impulses are propagated along a nerve has been estimated at about 100 feet in a second for both motor and sensory nerves, but varies according to the temperature, the degree of excitability, the strength of the stimulus, etc.

Law of Contraction. If a *feeble* galvanic current be applied to a recent and excitable nerve, contraction is produced in the muscles only upon the *making* of the circuit with both the direct and inverse current.

If the current be *moderate* in intensity, the contraction is produced in the muscle both upon the *making* and *breaking* of the circuit, with both the direct and inverse currents.

If the current be *intense*, contraction is produced only when the circuit is *made* with the direct current, and only when it is *broken* with the inverse current.

CRANIAL NERVES.

The **Cranial Nerves** come off from the base of the brain, pass through the foramina in the walls of the cranium, and are distributed to the skin, muscles and organs of sense in the face and head.

According to the classification of Sæmmering, there are 12 pairs of nerves, enumerating them from before backward, as follows, viz.:—

1st Pair, or Olfactory.	7th Pair, or Facial, Portio dura.
2d Pair, or Optic.	8th Pair, or Auditory, Portio mollis.
3d Pair, or Motor oculi communis.	9th Pair, or Glosso-pharyngeal.
4th Pair, or Patheticus, Trochlearis.	10th Pair, or Pneumogastric.
5th Pair, or Trifacial, Trigemini.	11th Pair, or Spinal accessory.
6th Pair, or Abducens.	12th Pair, or Hypoglossal.

The **Cranial Nerves** may also be classified physiologically, according to their function, into three groups: 1. Nerves of special sense. 2. Nerves of motion. 3. Nerves of general sensibility.

1st Pair. Olfactory.

Apparent Origin. From the inferior and internal portion of the anterior lobes of the cerebrum by three roots, viz.: an *external white root*, which passes across the fissure of Sylvius to the middle lobe of the cerebrum; an *internal white root*, from the most posterior part of the anterior lobe; a *gray root*, from the gray matter in the posterior and inner portion of the inferior surface of the anterior lobe.

Deep Origin. Not satisfactorily determined.

Distribution. The olfactory nerve, formed by the union of the three roots, passes forward along the under surface of the anterior lobe to the ethmoid bone, where it expands into the olfactory bulb. This bulb contains ganglionic cells, is grayish in color and soft in consistence; it gives off from its under surface from 15 to 20 nerve filaments, the *true olfactory nerves*, which pass through the cribriform plate of the ethmoid bone, and are distributed to the schneiderian mucous membrane. This membrane extends from the cribriform plate of the ethmoid bone downward, about one inch.

Properties. The olfactory nerves give rise to neither motor nor sensory phenomena when stimulated. They carry simply the special impressions of odorous substances. Destruction or injury of the olfactory bulbs is attended by a loss of the sense of smell.

Function. Governs the sense of smell. Conducts the impressions which give rise to odorous sensations.

2d Pair. Optic.

Apparent Origin. From the anterior portion of the optic commissure.

Deep Origin. An *external white root*, from the corpus geniculatum externum; an *internal white root*, from the corpus geniculatum internum and the anterior tubercula quadrigemina; a *gray root*, from the gray matter in the floor of the 3d ventricle. Filaments also come from the optic thalami and cerebral peduncles.

Distribution. The two roots unite to form a flattened band, the optic tract, which winds around the crus cerebri to decussate with the nerve of the opposite side, forming the *optic chiasm*. The decussation of fibres is not complete; some of the fibres of the left optic tract going to the outer half of the eye of the same side, and to the inner half of the eye of the opposite side; the same holds true for the right optic tract.

The *optic nerves proper* arise from the commissure, pass forward through the optic foramina, and are finally distributed in the *retina*.

Pages 89-104 missing

Division of the *posterior* columns impairs the power of muscular co-ordination, such as is witnessed in locomotor ataxia.

The *gray matter* is probably both *insensible* and *inexcitable* under the influence of direct stimulation.

A *transverse section* of one lateral half of the cord produces:—

(1) On the *same* side, paralysis of voluntary motion and a relative or absolute elevation of temperature and an increased flow of blood in the paralyzed parts; hyperæsthesia for the sense of contact, tickling, pain and temperature.

(2) On the *opposite* side, complete anæsthesia as regards contact, and tickling and temperature, in the parts corresponding to those which are paralyzed in the opposite side. Complete preservation of voluntary power and of the muscular sense.

A *vertical section* through the middle of the *gray matter* results in the loss of sensation on both sides of the body below the section, but no loss of voluntary power.

FUNCTIONS OF THE SPINAL CORD.

1. As a Conductor. The *Lateral columns*, particularly the posterior portions, the “pyramidal tracts,” and the columns of Türek, are the channels through which pass the voluntary motor impulses from the brain to the large multipolar nerve cells in the anterior cornuæ of gray matter, and through them become connected with the anterior roots which transmit the motor stimuli to the muscles.

The *Anterior columns*, especially the portion surrounding the anterior cornuæ, the “anterior radicular zones,” are composed of short longitudinal commissural fibres, which serve to connect together different segments of the spinal cord, a condition required for the coördination of muscular movements.

The *Posterior columns* are composed of short and long commissural fibres which connect together different segments of the cord. They are insensible to direct irritation, but aid in the coördination of muscular movements in walking, standing, running, etc. Degeneration of the posterior columns gives rise to the lack of muscular coördination observed in locomotor ataxia.

The *Gray matter*, and especially that portion immediately surrounding the central canal, transmits the sensory nerve fibres from the posterior roots up to the brain. Decussation of the sensory fibres takes place throughout the whole length of the gray matter.

The *Multipolar cells* of the *anterior cornuæ* are connected with the generation and transmission of motor impulses outward; are centres for reflex movements; are the trophic centres for the motor nerves and muscular fibres to which they are distributed. The anterior roots give passage to the vaso-constrictor and vaso-dilator fibres which exert an influence upon the calibre of the blood vessels. *Complete destruction* of the anterior horns is followed by a paralysis of motion, degeneration of the anterior roots, atrophy of muscles and bones and an abolition of reflex movements.

2. As an Independent Nerve Centre.

The spinal cord, by virtue of its containing ganglionic nerve matter, is capable of transforming impressions made upon the centripetal nerves into motor impulses, which are reflected outward through centrifugal nerves to muscles, producing movements. These reflex movements taking place through the gray matter, are independent of sensation and volition.

The mechanism involved in every reflex act is a sentient surface, a sensory nerve, a nerve centre, a motor nerve and muscle.

The *reflex excitability* of the cord may be—

(1) *Increased* by disease of the lateral columns, the administration of strychnia, and in frogs, by a separation of the cord from the brain, the latter apparently exerting an inhibitory influence over the former and depressing its reflex activity.

2. *Inhibited* by destructive lesions of the cord, *e. g.*, locomotor ataxia, atrophy of the anterior cornuæ, the administration of various drugs, and, in the frog, by irritation of certain regions of the brain. When the cerebrum alone is removed and the optic lobes stimulated, the time elapsing between the application of an irritant to a sensory surface and the resulting movement will be considerably prolonged. The optic lobes (Setchenow's centre) apparently generating impulses which, descending the cord, retard its reflex movements.

All movements taking place through the nervous system are of this reflex character, and may be divided into *excito-motor*, *sensory-motor* and *ideo-motor*.

Classification of Reflex Movements. (Küss.) They may be divided into four groups, according to the route through which the centripetal and centrifugal impulses pass.

1. Those normal reflex acts, *e. g.*, deglutition, coughing, sneezing, walking, etc., pathological reflex acts, *e. g.*, tetanus, vomiting, epilepsy, which take place both centripetally and centrifugally, through spinal nerves.

2. Reflex acts which take place in a centripetal direction through a

cerebro-spinal sensory nerve, and in a centrifugal direction through a sympathetic motor nerve, usually a vaso-motor nerve, *e.g.*, the normal reflex acts, which give rise to most of the secretions, pallor and blushing of the skin, certain movements of the iris, certain modifications in the beat of the heart; the pathological, which, on account of the difficulty in explaining their production, are termed *metastatic*, *e.g.*, ophthalmia, coryza, orchitis, which depend on a reflex hyperæmia; amaurosis, paralysis, paraplegia, etc., due to a reflex anæmia.

3. Reflex movements, in which the centripetal impulse passes through a sympathetic nerve, and the centrifugal through a cerebro-spinal nerve; most of these phenomena are pathological, *e.g.*, convulsions from intestinal irritation produced by the presence of worms, eclampsia, hysteria, etc.

4. Reflex actions, in which both the centripetal and centrifugal impulses pass through filaments of the sympathetic nervous system, *e.g.*, those obscure reflex actions which preside over the secretions of the intestinal fluids, which unite the phenomena of the generative organs, the dilatation of the pupils from intestinal irritation (worms), and many pathological phenomena.

Laws of Reflex Action. (Pflüger.)

1. *Law of Unilaterality.* If a feeble irritation be applied to one or more sensory nerves, movement takes place usually on one side only, and that upon the same side as the irritation.

2. *Law of Symmetry.* If the irritation becomes sufficiently intense, motor reaction is manifested, in addition, in corresponding muscles of the opposite side of the body.

3. *Law of Intensity.* Reflex movements are usually more intense on the side of the irritation; at times the movements of the opposite side equal them in intensity, but they are usually less pronounced.

4. *Law of Radiation.* If the excitation still continues to increase, it is propagated upward, and motor reaction takes place through centrifugal nerves coming from segments of the cord higher up.

5. *Law of Generalization.* When the irritation becomes very intense, it is propagated to the medulla oblongata; motor reaction then becomes general, and is propagated up and down the cord, so that all the muscles of the body are thrown into action, the medulla oblongata acting as a focus whence radiate all reflex movements.

Special Centres in the Spinal Cord.

Genito-spinal centre. In the lower portion of the spinal cord are located the centres which control the sphincter muscles of the rectum and

bladder, the erection of the penis, the emission of the semen, the action of the uterus during parturition, etc.

Cilio-spinal centre. Situated in the spinal cord between the 6th cervical and 2d dorsal nerves; stimulation of the cord in this situation produces a dilatation of both pupils through filaments of the sympathetic, which take their origin from this region of the cord.

Throughout the spinal cord are situated numerous centres which preside over the following reflexes, viz:—

The *patellar tendon reflex* takes place through the segments from which arise the 2d, 3d and 4th lumbar nerves; the *cremasteric reflex* through the segment from which arise the 1st and 2d lumbar nerves; the *abdominal reflex* through the segments between the 8th and 12th dorsal nerves; the *epigastric reflex* through the segments from which arise the 4th, 5th and 6th dorsal nerves.

Paralysis from Disease of the Spinal Cord.

Seat of Lesion. If it be in the *lower part* of the *sacral canal*, there is paralysis of the compressor urethræ, accelerator urinæ, and sphincter ani muscles; no paralysis of the muscles of the leg.

At the upper limit of the sacral region. Paralysis of the muscles of the bladder, rectum and anus; loss of sensation and motion in the muscles of the legs, except those supplied by the anterior crural and obturator, viz: psoas iliacus, Sartorius, pectineus, adductor longus, magnus and brevis, obturator, vastus externus and internus, etc.

At the upper limit of the lumbar region. Sensation and motion paralyzed in both legs; loss of power over the rectum and bladder; paralysis of the muscular walls of the abdomen interfering with expiratory movements.

At the lower portion of the cervical region. Paralysis of the legs. etc., as above; in addition, paralysis of all the intercostal muscles and consequent interference with respiratory movements; paralysis of muscles of the upper extremities, except those of the shoulders.

Above the middle of the cervical region. In addition to the preceding, difficulty of deglutition and vocalization, contraction of the pupils, paralysis of the diaphragm, scalene muscles, intercostals, and many of the accessory respiratory muscles; death resulting immediately, from arrest of respiratory movements.

Anterior half of spinal cord. Paraplegia developing symmetrically.

Posterior half of spinal cord. Characteristic symptoms of locomotor ataxia or tabes dorsalis.

In the gray substance in the vicinity of the central canal and anterior horns. If the lesion be acute, symptoms characteristic of acute spinal paralysis manifest themselves; if chronic, symptoms characteristic of progressive muscular atrophy.

MEDULLA OBLONGATA.

The *Medulla Oblongata* is the expanded portion of the upper part of the spinal cord. It is pyramidal in form and measures one and a half inches in length, three-quarters of an inch in breadth, half an inch in thickness, and is divided into two lateral halves by the anterior and pos-



View of Cerebellum in section, and of Fourth Ventricle, with the neighboring parts. (From Sappey.)

1. Median groove fourth ventricle, ending below in the *calamus scriptorius*, with the longitudinal eminences formed by the fasciculi teretes, one on each side. 2. The same groove, at the place where the white streaks of the auditory nerve emerge from it to cross the floor of the ventricle. 3. Inferior peduncle of the cerebellum, formed by the restiform body. 4. Posterior pyramid, above this is the *calamus scriptorius*. 5. Superior peduncle of cerebellum, or *processus e cerebello ad testes*. 6 6. Fillet to the side of the *crura cerebri*. 7 7. Lateral grooves of the *crura cerebri*. 8. *Corpora quadrigemina*.—After Hirschfeld and Leveille.

terior median fissures, which are continuous with those of the cord. Each half is again subdivided by minor grooves, into four columns, viz: *anterior pyramid*, *lateral tract* and *olivary body*, *restiform body* and *posterior pyramid*.

1. The *anterior pyramid* is composed partly of fibres continuous with those of the anterior column of the spinal cord; but mainly of fibres derived from the lateral tract of the opposite side, by *decussation*. The united fibres then pass upward through the pons Varolii and crura cerebri, and for the most part terminate in the corpus striatum and cerebrum.

2. The *lateral tract* is continuous with the lateral columns of the cord; its fibres in passing upward take three directions, viz: an external bundle joins the restiform body, and passes into the cerebellum; an internal bundle decussates at the median line and joins the opposite anterior pyramid; a middle bundle ascends beneath the olivary body, behind the pons, to the cerebrum, as the *fasciculus teres*.

The *olivary body* of each side is an oval mass, situated between the anterior pyramid and restiform body; it is composed of white matter externally and gray matter internally, forming the *corpus dentatum*.

3. The *restiform body*, continuous with the posterior column of the cord, also receives fibres from the lateral column. As the restiform bodies pass upward they diverge and form a space, the 4th ventricle, the floor of which is formed by gray matter, and then turn backward and enter the cerebellum.

4. The *posterior pyramid* is a narrow, white cord bordering the posterior median fissure; it is continued upward, in connection with the *fasciculus teres*, to the cerebrum.

The Gray Matter of the medulla is continuous with that of the cord. It is arranged with much less regularity, becoming blended with the white matter of the different columns, with the exception of the anterior. By the separation of the posterior columns, the transverse commissure is exposed, forming part of the floor of the 4th ventricle; special collections of gray matter are found in the posterior portions of the medulla, connected with the roots of origin of different cranial nerves.

Properties and Functions. The medulla is excitable anteriorly, and sensitive posteriorly to direct irritation. It serves (1) as a conductor of sensitive impressions upward from the cord, through the gray matter to the cerebrum; (2) as a conductor of voluntary impulses from the brain to the spinal cord and nerves, through its anterior pyramids; (3) as a conductor of coördinating impulses from the cerebellum, through the restiform bodies to the spinal cord.

As an Independent Reflex Centre. The medulla oblongata contains special collections of gray matter, which constitute independent nerve centres which preside over different functions, some of which are as follows, viz:—

1. A centre which controls the movements of *mastication*, through afferent and efferent nerves. (See page 24.)
2. A centre reflecting impressions which influence the *secretion* of *saliva*. (See page 25.)
3. A *centre for deglutition*, whence are derived motor stimuli exciting to action and coördinating the muscles of the palate, pharynx and œsophagus, necessary for the swallowing of the food.

NERVOUS CIRCLE OF DEGLUTITION. (2d and 3d Stages.)

Excitor or Centripetal Nerves.	{ Palatal branch of 5th pair. Pharyngeal branches of the glosso-pharyngeal. Superior laryngeal branches of the pneumogastric. (Esophageal branches of the pneumogastric.
Motor or Centrifugal Nerves.	{ Pharyngeal branches of the pneumogastric, derived from the spinal accessory. Hypoglossal and branches of the cervical plexus. Inferior or recurrent laryngeal. Motor filaments of the 3d division of the 5th pair. Portio dura.

4. A centre which coördinates the muscles concerned in the act of *vomiting*.
5. A *Speech centre*, coördinating the various muscles necessary for the accomplishment of articulation through the hypoglossal, facial nerves and the 2d division of the 5th pair.
6. A centre for the harmonization of muscles concerned in *expression*, reflecting its impulses through the facial nerve.
7. A *Cardiac centre*, which exerts (1) an *accelerating* influence over the heart's pulsations through accelerating nerve fibres emerging from the cervical portion of the cord, entering the inferior cervical ganglion, and thence passing to the heart; (2) an *inhibitory* or retarding influence upon the action of the heart, through fibres of the spinal accessory nerve running in the trunk of the pneumogastric.
8. A *Vaso-motor centre*, which, by alternately contracting and dilating the blood vessels through nerves distributed in their walls, regulates the quantity of blood distributed to an organ or tissue, and thus influences nutrition, secretion and calorification. The vaso-motor centre is situated in the medulla oblongata and pons Varolii, between the *corpora quadrigemina* and the *calamus scriptorius*. The vaso-motor fibres having their origin in this centre descend through the interior of the cord, emerge through the anterior roots of spinal nerves, enter the ganglia of the sympathetic, and thence

pass to the walls of the blood vessels, and maintain the *arterial tonus*; they may be divided into two classes, viz: *vaso-dilators*, e. g., chorda tympani, and *vaso-constrictors*, e. g., sympathetic fibres.

Division of the cord at the lower border of the medulla is followed by a dilatation of the entire vascular system and a marked fall of the blood pressure. *Galvanic stimulation* of the divided surface of the cord is followed by a contraction of the blood vessels and a rise in the blood pressure.

9. A *Diabetic centre*, irritation of which causes an increase in the amount of urine secreted, and the appearance of a considerable quantity of sugar.

10. A *Respiratory centre*, situated near the origin of the pneumogastric nerves, presides over the movements of respiration and its modifications, laughing, sighing, sobbing, sneezing, etc. It may be excited *reflexly* by the presence of carbonic acid in the lungs irritating the terminal pneumogastric filaments; or *automatically*, according to the character of the blood circulating through it; an *excess* of carbonic acid or a diminution of oxygen increasing the number of respiratory movements; a reverse condition diminishing the respiratory movements.

11. A *Spasm centre*, stimulation of which gives rise to convulsive phenomena.

NERVOUS CIRCLE OF RESPIRATION (ENTIRELY REFLEX).

Excitor or Centripetal Nerves.	{	Pulmonary branches of the pneumogastric.
		Superior laryngeal.
		Trifacial, or 5th pair.
		Nerves of general sensibility.
		Sympathetic nerve.
Motor or Centrifugal Nerves.	{	Phrenic, distributed to the diaphragm.
		Intercostals, distributed to the intercostal muscles.
		Facial nerve, or portio dura, to the facial muscles.
		External branch of spinal accessory, to the trapezius and sterno-cleido-mastoid muscles.

PONS VAROLII.

The Pons Varolii unites together the cerebrum above, the cerebellum behind, and the medulla oblongata below. It consists of transverse and longitudinal fibres, amidst which are irregularly scattered collections of gray or vesicular nervous matter.

The *transverse fibres* unite the two lateral halves of the cerebellum.

The *longitudinal fibres* are continuous (1) with the anterior pyramids of the medulla oblongata, which interlacing with the deep layers of the transverse fibres, ascend to the crura cerebri, forming their superficial or fasciculated portions; (2) with fibres derived from the olivary fasciculus, some of which pass to the tubercula quadrigemina, while others, uniting with fibres from the lateral and posterior columns of the medulla, ascend in the deep or posterior portions of the crura cerebri.

Properties and Functions. The superficial portion is *insensible* and *inexcitable* to direct irritation; the deeper portions appear to be *excitable*, consisting of descending motor fibres; the posterior portions are *sensible* but *inexcitable* to irritation.

Transmits motor impulses and sensory impressions from and to the cerebrum.

The *gray ganglionic* matter consists of centres which convert impressions into conscious sensations, and originate motor impulses, these taking place independent of any intellectual process; they are the seat of instinctive reflex acts; the centres which assist in the co-ordination of the automatic movements of station and progression.

CRURA CEREBRI.

The **Crura Cerebri** are largely composed of the longitudinal fibres of the pons (anterior pyramids, fasciculi teretes); after emerging from the pons they increase in size, and become separated into two portions by a layer of dark gray matter, the *locus niger*.

The *superficial* portion, the *crusta*, composed of the anterior pyramids, constitutes the *motor tract*, which terminates, for the most part, in the *corpus striatum*, but to some extent, also, in the cerebrum; the *deep portion*, made up of the fasciculi teretes and posterior pyramids and accessory fibres from the cerebellum, constitute the *sensory tract* (the *tegmentum*), which terminates in the *optic thalamus* and cerebrum.

Function. The crura are conductors of motor impulses and sensory impressions; the gray matter, the *locus niger*, assists in the coördination of the complicated movements of the eyeball and iris, through the motor oculi communis nerve. They also assist in the harmonization of general muscular movements; section of one crus giving rise to peculiar movements of rotation and somersaults forward and backward.

CORPORA QUADRIGEMINA.

The *Corpora Quadrigemina* are four small, rounded eminences, two on each side of the median line, situated immediately behind the third ventricle, and beneath the posterior border of the corpus callosum.

The *anterior* tubercles are oblong from before backward, and larger than the *posterior*, which are hemispherical in shape ; they are grayish in color, but consist of white matter externally and gray matter internally.

Both the anterior and posterior tubercles are connected with the optic thalami by commissural bands named the *anterior* and *posterior brachia*, respectively. They receive fibres from the olivary fasciculus and fibres from the cerebellum, which pass upward to enter the optic thalami.

The *corpora geniculata* are situated, one on the inner side and one on the outer side of each optic tract, behind and beneath the optic thalamus, and from their position are named the *corpora geniculata interna* and *externa*; they give origin to fibres of the optic nerve.

Functions. The *Tubercula quadrigemina* are the physical centres of sight, translating the luminous impressions into visual sensations. Destruction of these tubercles is immediately followed by a loss of the sense of sight; moreover, their action in vision is crossed, owing to the decussation of the optic tracts, so that if the tubercle of the right side be destroyed by disease or extirpated, in a pigeon, the sight is *lost* in the eye of the opposite side, and the iris loses its mobility.

The tubercula quadrigemina as nerve centres preside over the reflex movements which cause a dilation or contraction of the iris; irritation of the tubercles causing contraction, destruction causing dilatation. Removal of the tubercles on one side produces a temporary loss of power of the opposite side of the body, and a tendency to move around an axis is manifested, as after a section of one crus cerebri, which, however, may be due to giddiness and loss of sight.

They also assist in the coördination of the complex movements of the eye, and regulate the movements of the iris during the movements of accommodation for distance.

CORPORA STRIATA AND OPTIC THALAMI.

The *Corpora Striata* are two large ovoid collections of gray matter, situated at the base of the cerebrum, the larger portions of which are imbedded in the white matter, the smaller portions projecting into the anterior part of the lateral ventricle. Each striated body is divided, by a narrow band of white matter, into two portions, viz :—

1. The *Caudate nucleus*, the intraventricular portion, which is conical in shape, having its apex directed backward, as a narrow, tail-like process.

2. The *Lenticular nucleus*, imbedded in the white matter, and for the most part external to the ventricle; on the outer side of the lenticular nucleus is found a narrow band of white matter, the *external capsule*; and between it and the convolutions of the island of Reil, a thin band of gray matter, the *claustrum*; the corpora striata are grayish in color, and when divided present transverse striations, from the intermingling of white fibre and gray cells.

The *Optic Thalami* are two oblong masses situated in the ventricles posterior to the corpora striata, and resting upon the posterior portion of the crura cerebri. The internal surface projecting into the lateral ventricles is white, but the interior is grayish, from a commingling of both white fibres and gray cells. Separating the lenticular nucleus from the caudate nucleus and the optic thalamus, is a band of white tissue, the *internal capsule*.

The *internal capsule* is a narrow, bent tract of white matter, and is, for the most part, an expansion of the *motor tract* of the crura cerebri. It consists of two segments, *an anterior*, situated between the caudate nucleus and the anterior surface of the lenticular nucleus, and a *posterior*, situated between the optic thalamus and the posterior surface of the lenticular nucleus. These two segments unite at an obtuse angle, which is directed towards the median line. Pathological observation has shown that the nerve fibres of the direct and crossed pyramidal tracts can be traced upward through the anterior two-thirds of the posterior segment, into the centrum ovale, where, for the most part, they are lost; a portion, however, remaining united, ascend higher and terminate in the paracentral lobule, the superior extremity of the ascending frontal and parietal convolutions. The *sensory tract* can be traced upward, through the posterior third, into the cerebrum, where they probably terminate in the hippocampus major and uncinate convolution.

Functions. The *Corpora striata* are the centres in which terminate some of the fibres of the superficial or *motor tract* of the crura cerebri; others pass upward through the *internal capsule*, to be distributed to the cerebrum. It might be inferred, from their anatomical relations, that they are motor centres. *Irritation* by a weak galvanic current produces muscular movements of the opposite side of the body; *destruction* of their substance by a hemorrhage, as in apoplexy, is followed by a paralysis of motion of the opposite side of the body, but there is no loss of sensation.

When the hemorrhagic destruction involves the fibres of the anterior two-thirds of the posterior segment of the internal capsule, and thus separates them from their trophic centres in the cortical motor region, a descending degeneration is established, which involves the direct pyramidal tract of the same side and the crossed pyramidal tract of the opposite side.

Destruction of the posterior one-third of the posterior segment of the internal capsule is followed by a loss of sensation on the opposite side of the body, and a loss of the senses of smell and vision on the same side (Charcot). The precise function of the corpora striata is unknown, but they are in some way connected with motion.

The *Optic thalami* receive the fibres of the *tegmentum*, the posterior portion of the crura cerebri. They are insensible and inexcitable to direct irritation. Removal of one optic thalamus, or destruction of its substance by disease or hemorrhage, is followed by a loss of sensibility of the opposite side of the body, but there is no loss of motion; their precise function is also unknown, but in some way connected with *sensation*. In both cases their action is crossed.

CEREBELLUM.

The **Cerebellum** is situated in the inferior fossæ of the occipital bone, beneath the posterior lobes of the cerebrum. It attains its maximum weight, which is about 5 oz., between the twenty-fifth and fortieth years; the proportion between the cerebellum and cerebrum being 1 to $8\frac{1}{4}$.

It is composed of *two lateral hemispheres* and a central elongated lobe, the *vermiform process*; the two hemispheres are connected with each other by the fibres of the *middle peduncle* forming the superficial portion of the pons Varolii. It is brought into connection with the *medulla oblongata* and *spinal cord*, through the prolongation of the *restiform* bodies; with the cerebrum, by fibres passing upward beneath the corpora quadrigemina and the optic thalami, and then forming part of the diverging cerebral fibres.

Structure. It is composed of both white and gray matter, the former being internal, the latter external, and convoluted, for economy of space.

The *White matter* consists of a central stem, the interior of which is a dentated capsule of gray matter, the *corpus dentatum*. From the external surface of the stem of white matter processes are given off, forming the *laminae*, which are covered with gray matter.

The *Gray matter* is convoluted and covers externally the laminated processes; a vertical section through the gray matter reveals the following structures:—

1. A delicate *connective tissue layer*, just beneath the pia mater, containing rounded corpuscles, and branching fibres passing toward the external surface.

2. *The cells of Purkinje*, forming a layer of large, nucleated, branched nerve cells sending off processes to the external layer.

3. *A granular layer* of small, but numerous corpuscles.

4. *Nerve fibre layer*, formed by a portion of the white matter.

Properties and Functions. Irritation of the cerebellum is not followed by any evidences either of pain or convulsive movements; it is, therefore, *insensible* and *inexcitable*.

Co-ordination of Movements. Removal of the superficial portions of the cerebellum in pigeons produces *feebleness* and *want of harmony* in the muscular movements; as successive slices are removed, the movements become more irregular, and the pigeon becomes restless; when the last portions are removed, all power of *flying, walking, standing*, etc., is entirely gone, and the *equilibrium* cannot be maintained, the power of *coördinating* muscular movements being entirely gone. The same results have been obtained by operating on all classes of animals.

The following symptoms were noticed by Wagner, after removing the whole or a large part of the cerebellum. 1. A tendency on the part of the animal to throw itself on one side, and to extend the legs as far as possible. 2. Torsion of the head on the neck. 3. Trembling of the muscles of the body, which was general. 4. Vomiting and occasionally liquid evacuations.

Forced Movements. Division of *one crura cerebelli* causes the animal to fall on one side and roll rapidly on its longitudinal axis. According to Schiff, if the peduncle be divided from *behind*, the animal falls on the same side as the injury; if the section be made in *front*, the animal turns to the opposite side.

Disease of the cerebellum partially corroborates the results of experiments; in many cases symptoms of unsteadiness of gait, from a *want of coördination*, have been noticed.

Comparative anatomy reveals a remarkable correspondence between the development of the cerebellum and the complexity of muscular actions. It attains a much greater development, relatively to the rest of the brain, in those animals whose movements are very complex and varied in character, such as the kangaroo, shark and swallow.

The cerebellum may possibly exert some influence over the sexual function, but physiological and pathological facts are opposed to the idea of

its being the seat of the sexual instinct. It appears to be simply a centre for the coördination and equilibration of muscular movements.

CEREBRUM.

The Cerebrum is the largest portion of the encephalic mass, constituting about four-fifths of its weight; the average weight in the adult male is from 48 to 50 oz., or about 3 pounds, while in the adult female it is about 5 oz. less. After the age of 40 the weight of the cerebrum gradually diminishes at the rate of one ounce every ten years. In idiots the brain weight is often below the normal, at times not amounting to more than 20 ounces.

The Blood Supply to the cerebrum is unusually large, considering its comparative bulk; nearly *one-fifth* of the entire volume of blood being distributed to it by the carotid and vertebral arteries. These vessels anastomose so freely, and are so arranged within the cavity of the cranium, that an obstruction in one vessel will not interfere with the regular supply of blood to the parts to which its branches are distributed. A diminished amount, or complete cessation, of the supply of blood is at once followed by a suspension of its functional activity.

The cerebrum is connected with the pons Varolii and medulla oblongata through the crura cerebri, and with the cerebellum, through the superior peduncles. It is divided into two lateral halves, or hemispheres, by the longitudinal fissure running from before backward in the median line; each hemisphere is composed of both *white* and *gray* matter, the former being internal, the latter external; it covers the surfaces of the hemisphere which are infolded, forming convolutions, for economy of space.

Fissures.

1. The *Fissure of Sylvius* is one of the most important; it is the first to appear in the development of the foetal brain, being visible at about the third month; in the adult it is quite deep and well marked, running from the under surface of the brain upward, outward and backward, and forms a boundary between the frontal and temporo-sphenoidal lobes.

2. The *Fissure of Rolando* is second in importance, and runs from a point on the convexity near the median line transversely outward and downward toward the fissure of Sylvius, but does not enter it. It separates the frontal from the parietal lobe.

3. The *Parietal fissure*, arising a short distance behind the fissure of Rolando, upon the convexity of the hemisphere, runs downward and backward to its posterior extremity.

Secondary fissures of importance are found in different lobes of the cerebrum, separating the various convolutions. In the anterior lobe are found the *pre-central*, *superior frontal* and *inferior frontal* fissures; in the occipital lobe, are found the *parieto-occipital* and the *calcarine fissures*.

Convolutions. *Frontal Lobe.*

The *Ascending frontal convolution*, situated in front of the fissure of Rolando, runs downward and forward; it is continuous above with the anterior frontal, and below with the inferior frontal convolution.

The *Superior frontal convolution* is bounded internally by the longitudinal fissure, and externally by the superior frontal fissure; it is connected with the superior end of the frontal convolution, and runs downward and forward to the anterior extremity of the frontal lobe, where it turns backward, and rests upon the orbital plate of the frontal bone.

The *Middle frontal convolution*, the largest of the three, runs from behind forward, along the sides of the lobe, to its anterior part; it is bounded above by the superior and below by the inferior frontal fissures.

The *Inferior frontal convolution* winds around the ascending branch of the fissure of Sylvius, in the anterior and inferior portion of the cerebrum.

Parietal Lobe. The *Ascending parietal convolution* is situated just behind the fissure of Rolando, running downward and forward; above, it becomes continuous with the upper parietal convolution, and below, winds around to be united with the ascending frontal.

The *Upper parietal convolution* is situated between the parietal and longitudinal fissures.

The *Supra-marginal convolution* winds around the superior extremity of the fissure of Sylvius.

The *Angular convolution*, a continuation of the preceding, follows the parietal fissure to its posterior extremity, and then makes a sharp angle downward and forward.

Temporo-sphenoidal Lobe. Contains three well marked convolutions, the *superior*, *middle* and *inferior*, separated by well defined fissures, and continuous posteriorly with the convolutions of the parietal lobe.

The *Occipital Lobe* lies behind the parieto-occipital fissure, and contains the *superior*, *middle* and *inferior* convolutions, not well marked.

The *Central Lobe* or *Island of Reil*, situated at the bifurcation of the fissure of Sylvius, is a triangular-shaped cluster of six convolutions, the *gyri operi*, which are connected with those of the frontal, parietal, and temporo-sphenoidal lobes.

Structure. The *Gray matter* of the cerebrum, about one-eighth of an inch thick, is composed of five layers of nerve cells: (1) a superficial layer, containing few small multipolar ganglion cells; (2) small ganglion cells, pyramidal in shape; (3) a layer of large pyramidal ganglion cells with processes running off superiorly and laterally; (4) the granular formation containing nerve cells; (5) spindle shaped and branching nerve cells of moderate size.

The *White Matter* consists of three distinct sets of fibres—

1. The *diverging* or *peduncular* fibres are mainly derived from the columns of the cord and medulla oblongata; passing upward through the crura cerebri, they receive accessory fibres from the olivary fasciculus, corpora quadrigemina and cerebellum. Some of the fibres terminate in the optic thalami and corpora striata, while others radiate into the anterior, middle and posterior lobes of the cerebrum.

2. The *transverse commissural* fibres connect together the two hemispheres, through the corpus callosum and anterior and posterior commissures.

3. The *longitudinal commissural* fibres connect together different parts of the same hemisphere.

Functions. The cerebral hemispheres are the centres of the nervous system through which are manifested all the phenomena of the mind; they are the centres in which impressions are registered, and reproduced subsequently as ideas; they are the seat of intelligence, reason and will.

However important a centre the cerebrum may be, for the exhibition of this highest form of nervous action, it is not directly essential for the continuance of life; for it does not exert any control over those automatic reflex acts, such as respiration, circulation, etc., which regulate the functions of organic life.

From the study of comparative anatomy, pathology, vivisection, etc., evidence has been obtained which throws some light upon the physiology of the cerebral hemispheres.

1. *Comparative Anatomy* shows that there is a general connection between the size of the brain, its texture, the depth and number of convolutions, and the exhibition of mental power. Throughout the entire animal series, the increase in intelligence goes hand in hand with an increase in the development of the brain. In man there is an enormous increase in size over that of the highest animals, the anthropoids. The most cultivated races of men have the greatest cranial capacity; that of the educated European being about 116 cubic inches, that of the Australian being about

60 cubic inches, a difference of 56 cubic inches. Men distinguished for great mental power usually have large and well developed brains; that of Cuvier weighed 64 oz.; that of Abercrombie 63 oz.; the average being about 48 to 50 oz.; not only the size, but above all, the texture of the brain, must be taken into consideration.

2. *Pathology.* Any severe injury or disease disorganizing the hemispheres is at once attended by a disturbance, or entire suspension of mental activity. A blow on the head producing concussion, or undue pressure from cerebral hemorrhage destroys consciousness; physical and chemical alterations in the gray matter have been shown to coexist with insanity, loss of memory, speech, etc. Congenital defects of organization from imperfect development are usually accompanied by a corresponding deficiency of intellectual power and the higher instincts. Under these circumstances no great advance in mental development can be possible, and the intelligence remains at a low grade. In congenital idiocy not only is the brain of small size, but it is wanting in proper chemical composition; *phosphorus*, a characteristic ingredient of the nervous tissue, being largely diminished in amount.

3. *Experimentation* upon the lower animals by removing the cerebral hemispheres is attended by results similar to those observed in disease and injury. Removal of the cerebrum in pigeons produces complete abolition of intelligence, and destroys the capability of performing spontaneous movements. The pigeon remains in a condition of profound stupor, which is not accompanied, however, by a loss of sensation, or of the power of producing reflex or instinctive movements. The pigeon can be temporarily aroused by pinching the feet, loud noises, light placed before the eyes, etc., but soon relapses into a state of quietude, being unable to remember impressions and connect them with any train of ideas; the faculties of memory, reason and judgment being completely abolished.

LOCALIZATION OF FUNCTIONS.

The *Faculty of articulate language*, by which the individual associates ideas and words comprises two distinct faculties, viz: 1. The power of recalling particular words; 2. The coördination of muscles necessary for their articulation.

Aphasia is a condition in which the power of expressing ideas in words is completely lost. Pathological observation has shown that this condition is frequently associated with disease of the 3d frontal convolution of the left side, and also the convolutions of the island of Reil, parts nourished by the middle cerebral artery. It usually coexists with right hemiplegia; at

times aphasia results from disease of corresponding structures of the right side. The faculty of articulate language may be located in the 3d frontal convolution and the island of Reil, usually on the left side of the brain.

Motor Centres. The gray matter covering the cerebral hemispheres is both insensible and inexcitable to ordinary mechanical and chemical stimuli; but when a galvanic current of low intensity is applied to particular regions of the cerebrum of a monkey, in which the general arrangement of the convolutions approximates that of man, definite co-ordinate movements occur on the opposite side of the body, *e. g.*, rotation, flexion and extension of the limbs, and movements of the facial muscles.

The regions in which the *motor centres* are located are the convolutions around the fissure of Rolando, especially the *ascending frontal* and the *ascending parietal*. Destruction of the gray matter of these convolutions is followed by a paralysis of motion on the opposite side of the body; destruction of circumscribed areas in these convolutions causes paralysis of the groups of muscles excited to action by electrical stimulation of such areas.

The *antero-frontal* and *occipital lobes* are not excitable to electrical stimulation.

Special Centres. The *superior* and *middle frontal* convolutions appear to be connected with the intellectual faculties; ablation of these convolutions causes a marked impairment of the intelligence and of the faculty of attentive observation, without impairing either sensation or motion.

The *Visual centre* is located in the angular convolution. If this centre be destroyed, blindness of the opposite eye results, which is, however, only temporary if the opposite angular convolution be intact; destruction of both causes a complete and permanent loss of visual perceptions.

The *Auditory* and *Taste centres* are located in the superior and inferior temporo sphenoidal convolutions respectively; destruction of these regions abolishes the sense of hearing and taste. The *sense of smell* is situated in the uncinate convolution or cornu ammonis.

Systemic sensations are probably located in the occipital lobes; their ablation is followed by a state of great depression, loss of appetite, and a refusal to take food.

The *centre for sensory impressions* is located in the posterior portion of the cerebrum; destruction of the posterior portion of the internal capsule causes loss of sensation on the opposite side of the body.

SYMPATHETIC NERVOUS SYSTEM.

The **Sympathetic Nervous System** consists of a chain of ganglia connected together by longitudinal nerve filaments, situated on each side of the spinal column, running from above downward. The two ganglionic cords are connected together in the interior of the cranium by the ganglion of Ribes, on the anterior communicating artery, and terminate in the ganglion impar, situated at the tip of the coccyx.

The chain of ganglia is divided into groups and named according to the localities in which they are found, viz: *cranial*, four in number; *cervical*, three; *thoracic*, twelve; *lumbar*, five, *sacral*, five; *coccygeal*, one. Each ganglion consists of a collection of *vesicular nervous matter*, among which are found *tubular* and *gelatinous* nerve fibres. The ganglia are reinforced by motor and sensory fibres from the cerebro-spinal nervous system.

The **Ganglia** are distinct nerve fibres, from which branches are distributed to glands, arteries, muscles and to the cerebral and spinal nerves; many pass, also, to the visceral ganglia, *e. g.*, cardiac, semilunar, pelvic, etc.

Cephalic Ganglia.

1. The *Ophthalmic* or *Ciliary* ganglion is situated in the orbital cavity posterior to the eyeball; it is of small size, and of a reddish-gray color; receives filaments of communication from the motor oculi, ophthalmic branch of the 5th pair, and the carotid plexus. Its filaments of distribution are the ciliary nerves which pass to the iris and ciliary muscle.

Function. It is the centre through which the reflex acts take place by which the pupil is contracted or dilated; controls the movement of accommodation for vision at different distances.

2. The *Spheno-palatine*, or Meckel's ganglion, triangular in shape, is situated in the spheno-maxillary fossa; receives filaments from the facial (Vidian nerve), and the superior maxillary branch of the 5th nerve. Its filaments of distribution pass to the gums, the soft palate, levator palati, and azygos uvulæ muscles.

The *Otic*, or Arnold's ganglion, is of small size, oval in shape, and situated beneath the foramen ovale; receives a motor filament from the facial and sensory filaments from the glosso-pharyngeal and 5th nerve; sends filaments to the mucous membrane of the tympanic cavity and to the tensor tympani muscle.

4. The *Submaxillary* ganglion, situated in the sub-maxillary gland, receives motor filaments from the chorda tympani and sensory filaments from the lingual branch of the 5th nerve. Regulates to some extent the secretion of saliva.

Cervical Ganglia.

The *Superior cervical* ganglion is fusiform in shape, of a grayish-red color, and situate opposite the 2d and 3d cervical vertebræ; it sends branches to form the carotid and cavernous plexuses which follow the course of the carotid arteries to their distribution; also sends branches to join the glosso-pharyngeal and pneumogastric, to form the pharyngeal plexus.

The *Middle cervical* ganglion, the smallest of the three, is occasionally wanting; it is situated opposite the 5th cervical vertebra; sends branches to the superior and inferior cervical ganglion, and to the thyroid artery.

The *Inferior cervical* ganglion, irregular in form, is situated opposite the last cervical vertebræ; it is frequently fused with the first thoracic ganglion.

The *superior, middle and inferior cardiac nerves*, arising from these cervical ganglia, pass downward and forward to form the deep and superficial cardiac plexuses located at the bifurcation of the trachea, from which branches are distributed to the heart, coronary arteries, etc.

The Thoracic Ganglia are usually twelve in number, placed against the heads of the ribs behind the pleura; they are small in size and gray in color; they communicate with the cerebro-spinal nerves by two filaments, one of which is white, the other gray.

The *great splanchnic nerve* is formed by the union of branches from the sixth, seventh, eighth and ninth ganglia; it passes through the diaphragm to the semi-lunar ganglion.

The *lesser splanchnic nerve* is formed by the union of filaments from the tenth and eleventh ganglia, and is distributed to the cœliac plexus.

The *renal splanchnic nerve* arises from the last thoracic ganglion and terminates in the renal plexus.

The *semi-lunar ganglia*, the largest of the sympathetic, are situated by the side of the cœliac axis; they send radiating branches to form the *solar plexus*; from the various plexuses, nerves follow the gastric, splenic, hepatic, renal, etc., arteries, into the different abdominal viscera.

The Lumbar Ganglia, four in number, are placed upon the bodies of the vertebra; they give off branches which unite to form the aortic lumbar plexus and the hypogastric plexus, and follow the blood vessels to their terminations.

The Sacral and Coccygeal Ganglia send filaments of distribution to all the blood vessels of the pelvic viscera.

Properties and Functions. The sympathetic nerve possesses both sensibility and the power of exciting motion, but these properties are much

less decided than in the cerebro-spinal system. Irritation of the ganglia does not produce any evidence of pain until some time has elapsed. If caustic soda be applied to the semilunar ganglia, or a galvanic current be passed through the splanchnic nerves, no instantaneous effect is noticed, as in the case of the cerebro-spinal nerves; but in the course of a few seconds a slow, progressive contraction of the muscular coat of the intestines is established, which continues for some time after the irritation is removed.

Division of the sympathetic nerve in the neck is followed by a vascular congestion of the parts above the section on the corresponding side, attended by an increase in the temperature; not only is there an increase in the amount of blood, but the rapidity of the blood current is very much hastened, and the blood in the veins becomes of a brighter color. *Galvanization* of the upper end of the divided nerve causes all of the preceding phenomena to disappear; the congestion decreases, the temperature falls, and the venous blood becomes dark again.

The sympathetic exerts a similar influence upon the circulation of the limbs and the glandular organs; *destruction* of the first thoracic ganglion and division of the nerves forming the lumbar and sacral plexuses is followed by a dilatation of the vessels, an increased rapidity of the circulation, and an elevation of temperature in the anterior and posterior limbs; galvanization of the peripheral ends of these nerves causes all of these phenomena to disappear. Division of the splanchnic nerve causes a dilatation of the blood vessels of the intestine.

These phenomena of the sympathetic nerve system are dependent upon the presence of vaso-motor nerves which, under normal circumstances, exert a tonic influence upon the blood vessels. These nerves, derived from the cerebro-spinal system, the medulla oblongata, leave the spinal cord by the *rami communicantes*, enter the sympathetic ganglia, and finally terminate in the muscular wall of the blood vessels.

Sleep is a periodical condition of the nervous system, in which there is a partial or complete cessation of the activities of the higher nerve centres. The *cause* of sleep is a diminution in the quantity of blood, occasioned by a contraction of the smaller arteries under the influence of the vaso-motor nerves.

During the waking state the brain undergoes a physiological waste, as a result of the exercise of its functions; after a certain length of time its activities become enfeebled, and a period of repose ensues, during which a regeneration of its substance takes place.

When the brain becomes enfeebled there is a diminished molecular activity and an accumulation of waste products; under these circumstances

it ceases to dominate the medulla oblongata and the spinal cord. These centres then act more vigorously, and diminish the calibre of the cerebral blood vessels through the action of the vaso-motor nerves, producing a condition of physiological anæmia and sleep; during this state waste products are removed, force is stored up, nutrition is restored, and waking finally occurs.

THE SENSE OF TOUCH.

The Sense of Touch is a modification of general sensibility, and located in the skin, which is especially adapted for this purpose, on account of the number of nerves and papillary elevations it possesses. The structures of the skin and the modes of terminations of the sensory nerves have already been considered.

The Tactile Sensibility varies in acuteness in different portions of the body; being most marked in those regions in which the tactile corpuscles are most abundant, *e. g.*, the palmar surface of the third phalanges of the fingers and thumb.

The *relative sensibility* of different portions of the body has been ascertained by means of a pair of compasses, the points of which are guarded by cork, and then determining how closely they could be brought together, and yet be felt at two distinct points. The following are some of the measurements:—

Point of tongue.....	$\frac{1}{2}$ of a line.
Palmar surface of third phalanx.....	1 line.
Red surface of lips.....	2 lines.
Palmar surface of metacarpus.....	3 “
Tip of the nose.....	3 “
Part of lips covered by skin.....	4 “
Palm of hand.....	5 “
Lower part of forehead.....	10 “
Back of hand.....	14 “
Dorsum of foot.....	18 “
Middle of the thigh.....	30 “

The sense of touch communicates to the mind the idea of *resistance* only, and the varying degrees of resistance offered to the sensory nerves enables us to estimate, with the aid of the muscular sense, the qualities of *hardness* and *softness* of external objects. The idea of *space* or *extension* is obtained when the sensory surface or the external object changes its place in regard to the other; the *character* of the surface, its *roughness* or *smoothness*, is estimated by the impressions made upon the tactile papillæ.

Appreciation of Temperature.—The general surface of the body is more or less sensitive to differences of temperature, though this sensation is separate from that of touch; whether there are nerves especially adapted for the conduction of this sensation has not been fully determined. Under pathological conditions, however, the sense of touch may be abolished, while the appreciation of changes in temperature may remain normal.

This cutaneous surface varies in its sensibility to temperature in different parts of the body, and depends, to some extent, upon the thickness of the skin, exposure, habit, etc.; the inner surface of the elbow is more sensitive to changes in temperature than the outer portion of the arm; the left hand is more sensitive than the right; the mucous membrane less so than the skin.

Excessive heat or cold has the same effect upon the sensibility; the temperatures most readily appreciated are those between 50° F. and 115° F.

The sensation of *pain* and *tickling* appear to be conducted to the brain, also, by nerves different from those of touch; in abnormal conditions the appreciation of pain may be entirely lost, while touch remains unimpaired.

THE SENSE OF TASTE.

The Sense of Taste is localized mainly in the mucous membrane covering the superior surface of the tongue.

The Tongue is situated in the floor of the mouth; its base is directed backward, and connected with the hyoid bone, by numerous muscles, with the epiglottis and soft palate; its apex is directed forward against the posterior surface of the teeth.

The *substance* of the tongue is made up of intrinsic muscular fibres, the *linguales*; it is attached to surrounding parts, and its various movements performed by the extrinsic muscles, *e. g.*, stylo-glossus, genio-hyo-glossus, etc.

The *mucous membrane* covering the tongue is continuous with that lining the commencement of the alimentary canal, and is furnished with *vascular and nervous papillæ*.

The *papillæ* are analogous in their structure to those of the skin, and are distributed over the dorsum of the tongue, giving it its characteristic roughness.

There are three principal varieties:—

1. The *filiform papillæ* are most numerous, and cover the anterior two-thirds of the tongue; they are conical or filiform in shape, often prolonged into filamentous tufts, of a whitish color, and covered by horny epithelium.

2. The *fungiform papillæ* are found chiefly at the tip and sides of the tongue; they are larger than the preceding, and may be recognized by their deep red color.

3. The *circumvallate papillæ* are rounded eminences, from 8 to 10 in number, situated at the base of the tongue, where they form a V-shaped figure. They are quite large, and consist of a central projection of mucous membrane, surrounded by a wall, or circumvallation, from which they derive their name.

The Taste Beakers, supposed to be the true organs of taste, are flask-like bodies, ovoid in form, about the $\frac{1}{300}$ of an inch in length, situated in the epithelial covering of the mucous membrane, on the circumvallate papillæ. They consist of a number of fusiform, narrow cells, and curved so as to form the walls of this flask-like body; in the interior are elongated cells, with large, clear nuclei, the *taste cells*.

Nerves of Taste. The *chorda tympani* nerve, a branch of the facial, after leaving the cavity of the tympanum, joins the 3d division of the 5th nerve between the two pterygoid muscles, and then passes forward in the lingual branches, to be distributed to the mucous membrane of the anterior two-thirds of the tongue. Division or disease of this nerve is followed by a *loss of taste* in the part to which it is distributed.

The *glosso-pharyngeal* enters the tongue at the posterior border of the hyo-glossus muscle, and is distributed to the mucous membrane of the base and sides of the tongue, fauces, etc.

The *lingual branch of the trifacial* nerve endows the tongue with general sensibility; the *hypoglossal* endows it with motion.

The *nerves of taste* in the superficial layer of the mucous membrane form a fine plexus from which branches pass to the epithelium and penetrate it; others enter the taste beakers, and are directly connected with the taste cells.

The *seat* of the sense of taste has been shown by experiment to be the whole of the mucous membrane over the dorsum of the tongue, soft palate, fauces, and upper part of the pharynx.

The Sense of Taste enables us to distinguish the savor of substances introduced into the mouth, which is different from tactile sensibility. The sapid quality of substances appreciated by the tongue are designated as bitter, sweet, alkaline, sour, salt, etc.

The Essential Conditions for the production of the impressions of taste are (1) a state of solubility of the food; (2) a free secretion of the saliva, and (3) active movements on the part of the tongue, exerting pres-

sure against the roof of the mouth, gums, etc., thus aiding the solution of various articles and their osmosis into the lingual papillæ. Sapid substances, when in a state of solution, pass into the interior of the taste beakers and come into contact, through the medium of the taste cells, with the terminal filaments of the gustatory nerves.

THE SENSE OF SMELL.

The Sense of Smell is located in the mucous membrane lining the upper part of the nasal cavity, in which the olfactory nerves are distributed.

The Nasal Fossæ are two cavities, irregular in shape, separated by the vomer, the perpendicular plate of the ethmoid bone, and the triangular cartilage. They open anteriorly and posteriorly by the anterior and posterior nares, the latter communicating with the pharynx. They are lined by mucous membrane, of which the only portion capable of receiving odorous impressions is the part lining the upper one-third of the fossæ.

The Olfactory Nerves, arising by three roots from the posterior and inferior surface of the anterior lobes, pass forward to the cribriform plate of the ethmoid bone, where they each expand into an oblong body, the *olfactory bulb*. From its under surface from 15 to 20 filaments pass downward through the foramina, to be distributed to the olfactory mucous membrane, where they terminate in long, delicate, spindle-shaped cells, the *olfactory cells*, situated between the ordinary epithelial cells.

The olfactory bulbs are the centres in which odorous impressions are perceived as sensations; destruction of these bulbs being attended by an abolition of the sense of smell.

In animals which possess an acute sense of smell, there is a corresponding increase in the development of the olfactory bulbs.

The Essential Conditions for the sense of smell are, (1) a special nerve centre capable of receiving impressions and transforming them into odorous sensations. (2) Emanations from bodies which are in a gaseous or vaporous condition. (3) The odorous emanations must be drawn freely through the nasal fossæ; if the odor be very faint, a peculiar inspiratory movement is made, by which the air is forcibly brought into contact with the olfactory filaments. The secretions of the nasal fossæ probably dissolve the odorous particles.

Various substances, as ammonia, horseradish, etc., excite the sensibility of the mucous membrane, which must be distinguished from the perception of true odors.

THE SENSE OF SIGHT.

The Eyeball. The eyeball, or organ of vision, is situated within the orbital cavity, and loosely held in position by the fibrous capsule of Tenon. It rests upon a cushion of fat, which never disappears, except in cases of extreme starvation; it is protected from injury by the bony orbital walls, and is so situated as to permit an extensive range of vision.

Blood vessels and Nerves. The structures of the eyeball are supplied with blood by the ciliary arteries, which pierce the posterior surface around the optic nerve.

The *Ciliary* or *Ophthalmic ganglion*, about the size of a pin's head, situated in the posterior portion of the orbital cavity, receives filaments of communication from the trifacial or 5th nerve, the motor oculi or 3d nerve, and the sympathetic. From its anterior portion are given off the ciliary nerves, which enter the ball posteriorly and are distributed to the structures of which it is composed.

Structure. The form of the eyeball is that of a sphere; it is about one inch in the transverse diameter, and a little longer in the antero-posterior diameter, on account of its having the segment of a smaller sphere inserted into the anterior surface. It is composed of 3 coats; in its interior is contained the refracting apparatus.

The Sclerotic and Cornea together form the external coat of the eye; the former covering the posterior $\frac{5}{8}$, the latter covering the anterior $\frac{1}{8}$. The sclerotic is a dense, opaque, fibrous membrane, varying in thickness from the $\frac{1}{30}$ to the $\frac{1}{25}$ of an inch; it is composed of connective tissue and is slightly vascular. Posteriorly it is continuous with the sheath of the optic nerve, and is pierced by that nerve, as well as by the ciliary vessels and nerves; anteriorly its fibres become quite pale, and after passing into the cornea, transparent. It is a protective covering, and gives attachment to the tendons of the muscles by which the eyeball is moved.

The *Cornea* is a non-vascular, transparent membrane, composed for the most part of connective tissue in which are contained stellate corpuscles filled with a clear fluid. It is covered anteriorly by the basement membrane of the conjunctiva, upon which rests several layers of epithelial cells; posteriorly it is lined by the membrane of Descemet, which is reflected on to the anterior surface of the iris.

The Choroid, the Iris, the Ciliary Muscle and Ciliary Processes, together constitute the middle coat of the eye.

The *Choroid coat*, about the $\frac{1}{30}$ of an inch in thickness, is both a vas-


cular and pigmentary membrane ; it is of a dark brown color externally, and of a deep black internally.

The outer portion is made up of a rich network of vessels, the branches of the ciliary arteries and veins ; the inner portion, the pigmentary layer, is a delicate membrane formed of hexagonal cells, containing black pigment.

The *Function* of the choroid is mainly to absorb the rays of light which pass through the retina, and thus prevent them from interfering with the distinctness of vision by being again reflected upon the retina.

The *Iris* is a circular, muscular diaphragm, placed in the anterior por-

FIG. II.



SCLEROTIC COAT REMOVED TO SHOW THE CHOROID, CILIARY MUSCLE AND NERVES.

a. Sclerotic coat. *b.* Veins of the choroid. *c* Ciliary nerves. *d.* Veins of the choroid. *e.* Ciliary body. *f.* Iris.—From Holden's Anatomy.

tion of the eye, and perforated a little to the nasal side of the centre by a circular opening, the pupil ; it is attached by its periphery to the point of junction of the sclerotic and cornea. It is composed of a connective tissue stroma, blood vessels and non-striated muscular fibres, circular and radiating. The *circular* fibres surround the margin of the pupil like a sphincter, and are controlled by the 3d pair of nerves ; the *radiating* fibres (dilators of the pupil) radiate from the centre toward its circumference, and are controlled by the sympathetic system of nerves.

The *Ciliary muscle* is a grayish circular band, consisting of unstriated muscular fibres, about one-eighth of an inch long, running from before

backward; beneath the radiating fibres are small bands of circular fibres running around the eye. It arises from the line of junction of the sclerotic, cornea and iris; passing backward it is attached to the outer surface of the choroid; it is the principal agent in accommodation, and innervated by the 3d pair of nerves.

The **Retina** forms the internal coat of the eye; in the fresh state it is a delicate, transparent membrane, but soon becomes opaque and of a pinkish tint; it extends forward almost to the ciliary processes, where it terminates in the *ora serrata*. In the posterior portion of the retina, at a point corresponding to the axis of vision, is a rounded, elevated yellow spot, the *limbus luteus*, having a central depression, the *fovea centralis*; about $\frac{1}{10}$ of an inch to the inner side is the point of entrance of the optic nerve, where it spreads out to assist in the formation of the retina. The *arteria centralis retina* pierces the optic nerve near the sclerotic, runs forward in its substance and is distributed in the retina as far forward as the ciliary processes.

The Retina consists of nine distinct layers, from within outward, supported by connective tissue. 1. *Membrana limitans interna*. 2. Fibres of optic nerve. 3. Layers of ganglionic corpuscles. 4. Molecular layer. 5. Internal granular layer. 6. Molecular layer. 7. External granular layer. 8. *Membrana limitans externa*. 9. Layer of rods and cones.

In the *Fovea centralis*, at the point of most distinct vision, all of the layers disappear except the layer of rods and cones, which becomes somewhat longer and more slender.

The *Aqueous humor* is a clear fluid, alkaline in reaction, occupying the anterior chamber of the eye; this chamber is bounded in front by the cornea, posteriorly by the iris.

The *Vitreous humor* forms about four-fifths of the entire ball. It supports the retina, and is excavated anteriorly for the reception of the lens; it is transparent, of a jelly-like consistence, and surrounded by a structureless, transparent membrane, the *hyaloid* membrane.

The *Crystalline lens* is situated immediately behind the pupil, in the concavity of the vitreous humor. It is inclosed in a highly elastic, transparent membrane, the *capsule*. The lens is a transparent, doubly-convex body, $\frac{1}{3}$ of an inch transversely, $\frac{1}{4}$ of an inch antero-posteriorly; it is held in position by the *suspensory ligament*, formed by a splitting of the hyaloid tunic, the external layer of which passes in front of the lens, the internal layer behind it. Its function is to refract the rays of light and bring them to a focus upon the retina.

Vision. The eye may be regarded as a *camera obscura*, in which images of external objects are thrown upon a screen, the retina, by means of a double convex lens.

The Essential Conditions for proper vision are: 1. *Certain refracting media, e. g., cornea, aqueous humor, and crystalline lens, by which the rays of light are so disposed as to form an image.* 2. *A diaphragm, the iris, which, by alternately contracting and dilating, increases or diminishes the amount of light entering the eye.* 3. *A sensitive surface, to*

FIG. 12.




DIAGRAM OF A VERTICAL SECTION OF THE EYE.

1. Anterior chamber filled with aqueous humor. 2. Posterior chamber. 3. Canal of Petit.

a. Hyaloid membrane. b. Retina (dotted line). c. Choroid coat (black line). d. Sclerotic coat. e. Cornea. f. Iris. g. Ciliary processes. h. Canal of Schlemm or Fontana. i. Ciliary muscle. (From Holden's Anatomy.)

receive the image and transmit the luminous impressions through the optic nerve to the brain. 4. A contractile structure, the ciliary muscle, which can so manipulate the lens as to enable external objects to be seen at near or far distances.

The Refracting Apparatus, by which parallel rays of light are brought to a focus on the retina, consists mainly of the crystalline lens, though aided by the cornea and aqueous humor. A ray of light passing through the pupil is refracted and concentrated by the lens at a given point pos-

terior to it. For the correct perception of images of external objects, the rays of light must be accurately focused on the retina; in order that this may be accomplished, the lens must have a certain density and a proper curvature of its surfaces. When the lens is too convex, its refracting power is greatly increased, the rays of light are brought to a focus in front of the retina, and the visual perception becomes dim and confused. When it is too flat, the rays are not focused at all, and the resulting perception is the same.

The *Crystalline lens*, therefore, produces a distinct perception of the outline and form of external objects.

Action of the Iris. The iris, consisting of contracting and dilating fibres, transmits and regulates the quantity of light passing through its central aperture, the pupil, which is necessary for distinct vision.

If the light be too intense or excessive, the circular fibres contract under the stimulus of the 3d pair of nerves, and the aperture is diminished in size; if the quantity of light be insufficient, the dilating fibres contract under the stimulus of the sympathetic, and the pupillary aperture is increased in size.

The Retina, which is formed partly by the expansion of the optic nerve, and partly by new nervous structures, is the membrane which receives the impressions of light. Its posterior surface, which is in contact with the choroid, and especially the layer of rods and cones, is the sensitive portion, in which the rays of light produce their effects.

The *point of most distinct vision* is in the macula lutea, and especially in its central depression, the fovea, which corresponds to the central axis of the eye; it is situated about $\frac{1}{10}$ of an inch to the outside of the entrance of the optic nerve. It is at this point that images of external objects are seen most distinctly, while all around it the perceptions are more or less obscure; at the macula all the layers disappear except the layer of rods and cones.

Blind Spot. At the point of entrance of the optic nerve is a region in which the rays of light make no impression, owing to the absence of the proper retinal elements; the fibres of the optic nerve being insensible to the action of light.

The *course* which a ray of light takes is as follows: After passing through the cornea, lens, and vitreous humor and the layers of the retina, it is finally arrested by the pigmentary layer of the choroid; here it excites in the layer of rods and cones some physical or chemical change, which is then transmitted to the fibres of the optic nerve, and thence to the brain, where it is perceived as a sensation of light.

The Accommodation of the eye to vision for different distances is accomplished by a change in the convexities of the lens, caused by the action of the ciliary muscle. When the eye is accommodated for vision at far distances, the structures are in a passive condition and the lens is flattened; when it is adjusted for vision at short distances, the convexities of the lens are increased.

When the *Ciliary muscle* contracts and draws the choroid coat forward, the suspensory ligament is relaxed and the lens becomes more convex, in virtue of its own elasticity.

Optical Defects. *Astigmatism* is a condition of the eye which prevents vertical and horizontal lines from being focused at the same time, and is due to a greater curvature of the eye in one direction than another.

Spherical aberration is a condition in which there is an indistinctness of an image from the unequal refraction of the rays of light passing through the circumference and the centre of the lens; it is corrected mainly by the iris, which cuts off the marginal rays, and only transmits those passing through the centre.

Chromatic aberration, in which the image is surrounded by a colored margin, from the decomposition of the rays of light into their elementary parts, is corrected by the different refractive powers of the transparent media in front of the retina.

Myopia, or *short-sightedness*, is caused by an abnormal increase in the antero-posterior diameter of the eyeball; the lens being too far removed from the retina, forms the image in front of it, and the perception becomes dim and blurred. *Concave* glasses correct this defect, by preventing the rays from converging too soon.

Hypermetropia, or *long-sightedness*, is caused by a shortening of the antero-posterior diameter; the lens consequently focuses the rays of light behind the retina. *Convex* glasses correct this defect, by converging the rays of light more anteriorly.

Presbyopia is a loss of the power of accommodation of the eye to near objects, and usually occurs between the ages of 40 and 60; it is remedied by the use of a convex eye-glass.

Accessory Structures. The muscles which move the eyeball are six in number; the superior and inferior recti, the external and internal recti, the superior and inferior oblique muscles. The four recti muscles, arising from the apex of the orbit, pass forward and are inserted into the sides of the sclerotic coat; the *superior* and *inferior* muscles rotate the

eye around a *horizontal* axis; the *external* and *internal* rotate it around a *vertical* axis.

The *Superior oblique* muscle, having the same origin, passes forward to the inner and upper angle of the orbital cavity, where its tendon passes through a cartilaginous pulley; it is then reflected backward and inserted into the sclerotic just behind the transverse diameter. Its *function* is to rotate the eyeball in such a manner as to direct the pupil *downward* and *outward*.

The *inferior oblique* muscle arises at the inner angle of the orbit and then passes outward and backward to be inserted into the sclerotic. Its function is to rotate the eyeball and direct the pupil *upward* and *outward*.

By the associated action of all these muscles, the eyeball is capable of performing all the varied and complex movements necessary for distinct vision.

The *Eyelids*, bordered with short, stiff hairs, shade the eye and protect it from injury. On the posterior surface, just beneath the conjunctiva, are the Meibomian glands, which secrete an oily fluid; it covers the edge of the lids and prevents the tears from flowing over the cheek.

The *Lachrymal Glands* are ovoid in shape and situated at the upper and outer part of the orbital cavity; they open by from six to eight ducts at the outer portion of the upper lids.

The *Tears*, secreted by the lachrymal glands, are distributed over the cornea by the lids during the act of winking, and keep it moist and free from dust. The excess of tears passes into the lachrymal ducts, which begin by two minute orifices, one on each lid, at the inner canthus. They conduct the tears into the nasal duct, and so into the nose.

THE SENSE OF HEARING.

The **Organ of Hearing** is situated in the petrous portion of the temporal bone, and is divided into three portions, viz: the external ear, the middle ear and the internal ear.

The **External Ear** consists of two portions, the pinna or auricle, and the external auditory canal. The *former*, consisting of cartilage, which is irregularly folded and covered by integument, is united to the side of the head by ligaments and muscles; the *latter*, partly cartilaginous and partly bony, is about one and a quarter inches in length; it runs downward and forward from the concha to the middle ear, and is lined by a reflection of the general integument, in which is lodged a number of glands, which secrete the *cerumen*.

The *function* of the external ear is to collect the waves of sound coming from all directions and to transmit them to the *membrana tympani*.

The **Middle Ear or Tympanum** is an irregularly shaped cavity, narrow from side to side, but long in its vertical and antero-posterior diameters.

It is separated from the external ear by the *membrana tympani*, and from the internal ear by a second *membrana tympani*; it communicates posteriorly with the mastoid cells, anteriorly with the pharynx, through the Eustachian tube. It is lined by mucous membrane, and contains three small bones, forming a connected chain running across its cavity.

The *Membrana tympani* is a thin, delicate, translucent membrane, circular in shape and measuring about two-fifths of an inch in diameter; it is received into a delicate ring of bone, which in the adult becomes consolidated with the temporal bone; it is concave externally and situated obliquely, inclining at an angle of 45 degrees.

The membrane consists of three layers; the outer is formed by a reflection of the integument lining the external auditory canal; the middle is composed of fibrous tissue, and the internal of mucous membrane.

The *Function* of the *membrana tympani* is to receive and transmit the waves of sound to the chain of bones; it is capable of being made tense and lax by the action of the *tensor tympani* and *laxator tympani* muscles, so as to vibrate in unison with the waves of sound in the external auditory meatus. When the membrane is relaxed, its vibrations have a greater amplitude, and it appreciates sounds of a low pitch. When it is made tense it vibrates less forcibly and appreciates sounds of a high pitch.

The *Chain of bones* is formed by the malleus, incus and stapes, united together by ligaments. The *malleus* consists of a head, neck and handle, of which the latter is attached to the inner surface of the *membrana tympani*. The *incus*, or anvil bone, articulates with the head of the malleus by a capsular joint, and with the stapes by the end of its long process. The *stapes* resembles a stirrup in shape; it articulates externally with the long process of the incus, and internally its oval base is applied to the edges of the *foramen ovale*.

The *Function* of the chain of bones is to transmit the waves of sound across the tympanum to the internal ear; being surrounded by air, and acting as a solid rod, they prevent the vibrations from losing but little in intensity.

The *Tensor tympani muscle* arises mainly from the cartilaginous part of the Eustachian tube; it then passes backward into the tympanic cavity, where it bends at a right angle around a process of bone, and is inserted

into the root of the handle of the malleus. Its *function* is to draw the handle of the malleus internally, and thus increase the tension of the membrana tympani, so as to make it capable of vibrating with sounds of greater or less intensity; at the same time it tightens the joints of the chain of bones, so that they may the better conduct waves of sound to the internal ear, with but a slight loss of intensity.

The *Laxator tympani muscle*, arising from the spinous process of the sphenoid bone, passes backward through the Glasserian fissure, into the tympanic cavity, and is inserted into the neck of the malleus. Its *function* is to draw the handle of the malleus outward, and so relax the membrana tympani, and enable it to receive waves of sound of greater amplitude than when it is tense.

The *Stapedius muscle*, emerging from the cavity of the pyramid of bone projecting from the posterior wall of the tympanum, is inserted into the head of the stapes bone. Its *function* is to draw the stapes backward, preventing too great movement of the bone, and at the same time relaxing the membrana tympani.

The *Eustachian tube*, by means of which the middle ear communicates with the pharynx, is partly bony and partly cartilaginous in its structure. It is about one and a half inches in length; commencing at its opening in the pharynx, it passes upward and outward to the spine of the sphenoid bone, where it is slightly contracted; it then gradually dilates as it passes backward into the tympanic cavity. It is lined by mucous membrane, which is continued into the middle ear and into the mastoid cells.

The Eustachian tube permits the passage of air from the pharynx into the middle ear; in this way the pressure of the air within and without the membrana tympani is equalized, which is one of the essential conditions for the reception of sonorous vibrations.

By closing the mouth and nose, and blowing out the cheeks, air can be forced into the middle ear, producing undue pressure and bulging out of the membrana tympani; by making an effort at swallowing, with the mouth and nose closed, the air in the tympanum can be rarefied and the tympanic membrane will be pressed in. In both such cases the acuteness of hearing is very much diminished.

The pharyngeal orifice of the Eustachian tube is opened by the action of certain of the muscles of deglutition, viz. : the levator palati, tensor palati, and at times the palato-pharyngei muscles.

The Internal Ear, or Labyrinth, is located in the petrous portion of the temporal bone, and consists of an osseous and membranous portion.

The **Osseous Labyrinth** is divisible into three parts, viz.: the vestibule, the semi-circular canals and the cochlea.

The *Vestibule* is a small, triangular cavity, which communicates with the middle ear by the foramen ovale; in the natural condition it is closed by the base of the stapes bone. The filaments of the auditory nerve enter the vestibule through small foramina in the inner wall, at the fovea hemispherica.

The *Semi-circular canals* are three in number: the superior vertical, the inferior vertical and the horizontal, each of which opens into the cavity of the vestibule by two openings, with the exception of the two vertical, which at one extremity open by a common orifice.

The *Cochlea* forms the anterior part of the internal ear. It is a gradually tapering canal, about one and a half inches in length, which winds spirally around a central axis, the *modiolus*, two and a half times. The interior of the cochlea is partly divided into two passages by a thin plate of bone, the *lamina osseous spiralis*, which projects from the central axis two-thirds across the canal. These passages are termed the *scala vestibuli* and the *scala tympani*, from their communication with the vestibule and tympanum. The *scala tympani* communicates with the middle ear through the *foramen rotundum*, which, in the natural condition, is closed by the second membrana tympani; superiorly they are united by an opening, the *helicotrema*.

The whole interior of the labyrinth, the vestibule, the semi-circular canals, and the scala of the cochlea, contains a clear, limpid fluid, the *peri lymph*, secreted by the periosteum lining the osseous walls.

The **Membranous Labyrinth** corresponds to the osseous labyrinth with respect to form, though somewhat smaller in size.

The *Vestibular* portion consists of two small sacs, the *utricle* and *saccul*e.

The *Semi-circular canals* communicate with the *utricle* in the same manner as the bony canals communicate with the vestibule. The *saccul*e communicates with the membranous cochlea by the *canalis reuniens*. In the interior of the utricle and saccul, at the entrance of the auditory nerve, are small masses of carbonate of lime crystals, constituting the *otoliths*. Their function is unknown.

The *Membranous cochlea* is a closed tube, commencing by a blind extremity at the first turn of the cochlea, and terminating at its apex by a blind extremity also. It is situated between the edge of the *osseous lamina spiralis* and the outer wall of the bony cochlea, and follows it in its turns around the *modiolus*.

A transverse section of the cochlea shows that it is divided into two

portions by the osseous lamina and the basilar membrane: 1. The *scala vestibuli*, bounded by the periosteum and membrane of Reissner. 2. The *scala tympani*, occupying the inferior portion, and bounded above by the septum, composed of the osseous lamina and the membrana basilaris.

The *true membranous canal* is situated between the membrane of Reissner and the basilar membrane. It is triangular in shape, but is partly divided into a triangular portion and a quadrilateral portion by the *tectorial membrane*.

The *Organ of Corti* is situated in the quadrilateral portion of the canal, and consists of pillars or rods, of the consistence of cartilage. They are arranged in two rows; the one internal, the other external; these rods rest upon the basilar membrane; their bases are separated from each other, but their upper extremities are united, forming an arcade. In the internal row it is estimated there are about 3500, and in the external row about 5200 of these rods.

On the inner side of the internal row is a single layer of elongated hair cells; on the outer surface of the external row are three such layers of hair cells. Nothing definite is known as to their function.

The *Endolymph* occupies the interior of the utricle, saccule, membranous canals, and bathes the structures in the interior of the membranous cochlea, throughout its entire extent.

The Auditory Nerve at the bottom of the internal auditory meatus divides into (1) a vestibular branch, which is distributed to the utricle and semi-circular canals; (2) a cochlear branch, which passes into the central axis at its base, and ascends to its apex; as it ascends, fibres are given off, which pass between the plates of the osseous lamina, to be ultimately connected with the organ of Corti.

The *Function* of the semi-circular canals appears to be to assist in maintaining the equilibrium of the body; destruction of the vertical canal is followed by an oscillation of the head upward and downward; destruction of the horizontal canal is followed by oscillations from left to right. When the canals are injured on both sides, the animal loses the power of maintaining equilibrium upon making muscular movements.

Function of the Cochlea. It is regarded as possessing the power of appreciating the quality of pitch and the shades of different musical tones. The elements of the organ of Corti are analogous, in some respects, to a musical instrument, and are supposed, by Helmholtz, to be tuned so as to vibrate in unison with the different tones conveyed to the internal ear.

Summary. The waves of sound are gathered together by the pinna and external auditory meatus, and conveyed to the membrana tympani. This membrane, made tense or lax by the action of the tensor tympani and laxator tympani muscles, is enabled to receive sound waves of either a high or low pitch. The vibrations are conducted across the middle ear by the chain of bones to the foramen ovale, and by the column of air of the tympanum to the foramen rotundum, which is closed by the second membrana tympani; the pressure of the air in the tympanum being regulated by the Eustachian tube.

The internal ear finally receives the vibrations which excite vibrations successively in the perilymph, the walls of the membranous labyrinth, the endolymph, and, lastly, the terminal filaments of the auditory nerve, by which they are conveyed to the brain.

VOICE AND SPEECH.

The Larynx is the organ of voice. Speech is a modification of voice, and is produced by the teeth and the muscles of the lips and tongue, co-ordinated in their action by stimuli derived from the cerebrum.

The Structures entering into the formation of the larynx are mainly the *thyroid*, *cricoid* and *arytenoid* cartilages; they are so situated and united by means of ligaments and muscles as to form a firm cartilaginous box. The Larynx is covered externally by fibrous tissue and lined internally with mucous membrane.

The Vocal Cords are four ligamentous bands, running antero-posteriorly across the upper portion of the larynx, and are divided into the *two superior* or *false* vocal cords, and the *two inferior* or *true* vocal cords; they are attached anteriorly to the receding angle of the thyroid cartilages and posteriorly to the anterior part of the base of the arytenoid cartilages. The space between the true vocal cords is the *rima glottidis*.

The Muscles which have a direct action upon the movements of the vocal cords are nine in number, and take their names from their points of origin and insertion, viz: the two *crico-thyroid*, two *thyro-arytenoid*, two *posterior crico-arytenoid*, two *lateral crico arytenoid*, and one *arytenoid* muscles.

The *crico thyroid* muscles, by their contraction, render the vocal cords *more tense* by drawing down the anterior portion of the thyroid cartilage and approximating it to the cricoid, and at the same time tilting the posterior portion of the cricoid and arytenoid cartilages backward.

The *thyro-arytenoid*, by their contraction, *relax* the vocal cords by drawing the arytenoid cartilage forward and the thyroid backward.

The *posterior crico-arytenoid* muscles, by their contraction rotate the arytenoid cartilages outward and thus separate the vocal cords and enlarge the aperture of the glottis. They principally aid the respiratory movements during inspiration.

The *lateral crico-arytenoid* muscles are antagonistic to the former, and by their contraction rotate the arytenoid cartilages so as to approximate the vocal cords and constrict the glottis.

The *arytenoid* muscle assists in the closure of the aperture of the glottis.

The *inferior laryngeal* nerve animates all the muscles of the larynx, with the exception of the crico-thyroid.

Movements of the Vocal Cords. During respiration the movements of the vocal cords differ from those occurring during the production of voice.

At each *inspiration*, the true vocal cords are widely separated, and the aperture of the glottis is enlarged by the action of the crico-arytenoid muscles, which rotate outward the anterior angle of the base of the arytenoid cartilages; at each *expiration* the larynx becomes passive; the elasticity of the vocal cords returns them to their original position, and the air is forced out by the elasticity of the lungs and the walls of the thorax.

Phonation. As soon as phonation is about to be accomplished a marked change in the glottis is noticed with the aid of the laryngoscope. The true vocal cords suddenly become approximated and are made parallel, giving to the glottis the appearance of a narrow slit, the edges of which are capable of vibrating accurately and rapidly; at the same time their tension is much increased.

With the vocal cords thus prepared, the expiratory muscles force the column of air into the lungs and trachea through the glottis, throwing the edges of the cords into vibration.

The *pitch* of sounds depends upon the extent to which the vocal cords are made tense and the length of the aperture through which the air passes. In the production of sounds of a high pitch the tension of the vocal cords becomes very marked, and the glottis diminishes in length. When grave sounds, having a low pitch, are emitted from the larynx, the vocal cords are less tense and their vibrations are large and loose.

The *quality* of voice depends upon the length, size and thickness of the cords, and the size, form and construction of the trachea, larynx and the resonant cavities of the pharynx, nose and mouth.

The *compass* of the voice comprehends from two to three octaves. The range is different in the two sexes; the lowest note of the male being about one octave lower than the lowest note of the female; while the highest note of the male is an octave less than the highest of the female.

The *varieties* of voices, *e. g.*, bass, baritone, tenor, contralto, mezzo-soprano and soprano, are due to the length of the vocal cords; being longer when the voice has a low pitch, and shorter when it has a high pitch.

Speech is the faculty of expressing ideas by means of combination of sounds, in obedience to the dictates of the cerebrum.

Articulate sounds may be divided into *vowels* and *consonants*. The *vowel sounds*, *a, e, i, o, u*, are produced in the larynx by the vocal cords. The *consonantal sounds* are produced in the air passages above the larynx by an interruption of the current of air by the lips, tongue and teeth; the consonants may be divided into : (1) mutes, *b, d, k, p, t, c, g*; (2) dentals, *d, j, s, l, z*; (3) nasals, *m, n, ng*; (4) labials, *b, p, f, v, m*; (5) gutturals, *k, g, c*, and *g* hard; (6) liquids, *l, m, n, r*.

REPRODUCTION.

Reproduction is the function by which the species is preserved, and is accomplished by the organs of generation in the two sexes.

GENERATIVE ORGANS OF THE FEMALE.

The **Generative Organs of the Female** consist of the ovaries, Fallopian tubes, uterus and vagina.

The **Ovaries** are two small, ovoid, flattened bodies, measuring one inch and a half in length and three-quarters of an inch in width; they are situated in the cavity of the pelvis, and imbedded in the posterior layer of the broad ligament; attached to the uterus by a round ligament, and to the extremities of the Fallopian tubes by the fimbriae. The ovary consists of an external membrane of fibrous tissue, the *cortical* portion, in which are imbedded the *Graafian vesicles*, and an internal portion, the *stroma*, containing blood vessels.

The **Graafian Vesicles** are exceedingly numerous, but situated only in the cortical portion. Although the ovary contains the vesicles from the period of birth, it is only at the period of puberty that they attain their full development. From this time onward to the catamenial period, there is a constant growth and maturation of the Graafian vesicles. They consist of an external investment, composed of fibrous tissue and blood vessels, in the interior of which is a layer of cells forming the *membrana granulosa*; at its lower portion there is an accumulation of cells, the *proligerous disc*, in which the *ovum* is contained. The cavity of the vesicle contains a slightly yellowish, alkaline, albuminous fluid.

The **Ovum** is a globular body, measuring about the $\frac{1}{125}$ of an inch in diameter; it consists of an external investing membrane, the *vitelline membrane*, a central granular substance, the *vitellus*, or *yolk*, a nucleus, the *germinal vesicle*, in the interior of which is imbedded the nucleolus, or *germinal spot*.

The **Fallopian Tubes** are about four inches in length, and extend outward from the upper angles of the uterus, between the folds of the broad ligaments, and terminate in a fringed extremity, which is attached by one of the fringes to the ovary. They consist of three coats: (1) the external, or peritoneal, (2) middle, or muscular, the fibres of which are arranged in a circular or longitudinal direction, (3) internal, or mucous,

covered with ciliated epithelial cells, which are always waving from the ovary toward the uterus.

The **Uterus** is pyriform in shape, and may be divided into a body and neck; it measures about three inches in length and two inches in breadth in the unimpregnated state. At the lower extremity of the neck is the os externum; at the junction of the neck with the body is a constriction, the os internum. The cavity of the uterus is triangular in shape, the walls of which are almost in contact.

The walls of the uterus are made up of several layers of non-striated muscular fibres, covered externally by peritoneum, and lined internally by mucous membrane, containing numerous tubular glands, and covered by ciliated epithelial cells.

The **Vagina** is a membranous canal, from five to six inches in length, situated between the rectum and bladder. It extends obliquely upward from the surface, almost to the brim of the pelvis, and embraces at its upper extremity the neck of the uterus.

Discharge of the Ovum. As the Graafian vesicle matures, it increases in size, from an augmentation of its liquid contents, and approaches the surface of the ovary, where it forms a projection, measuring from one-fourth to one-half an inch in size. The maturation of the vesicle occurs periodically, about every twenty-eight days, and is attended by the phenomena of menstruation. During this period of active congestion of the reproductive organs the Graafian vesicle ruptures, the ovum and liquid contents escape, and are caught by the fimbriated extremity of the Fallopian tube, which has adapted itself to the posterior surface of the ovary. The passage of the ovum through the Fallopian tube into the uterus occupies from ten to fourteen days, and is accomplished by muscular contraction and the action of the ciliated epithelium.

Menstruation is a periodical discharge of blood from the mucous membrane of the uterus, due to a fatty degeneration of the small blood vessels. Under the pressure of an increased amount of blood in the reproductive organs, attending the process of ovulation, the blood vessels rupture, and a hemorrhage takes place into the uterine cavity; thence it passes into the vagina, where it is kept in a fluid condition from admixture with the vaginal mucus. Menstruation lasts from five to six days, and the amount of blood discharged averages about five ounces.

Corpus Luteum. For some time anterior to the rupture of a Graafian vesicle, it increases in size and becomes vascular; its walls become thickened, from the deposition of a reddish-yellow, glutinous substance, a

product of cell growth from the proper coat of the follicle and the *membrana granulosa*. After the ovum escapes, there is usually a small effusion of blood into the cavity of the follicle, which soon coagulates, loses its coloring matter, and acquires the characteristics of fibrin, but it takes no part in the formation of the corpus luteum. The walls of the follicle become convoluted, vascular, and undergo hypertrophy, until they occupy the whole of the follicular cavity. At its period of fullest development, the corpus luteum measures three-fourths of an inch in length and one-half inch in depth. In a few weeks the mass loses its red color, and becomes yellow, constituting the *corpus luteum* or *yellow body*. It then begins to retract, and becomes pale; and at the end of two months nothing remains but a small cicatrix upon the surface of the ovary. Such are the changes in the follicle, if the ovum has not been impregnated.

The corpus luteum, after impregnation has taken place, undergoes a much slower development, becomes larger, and continues during the entire period of gestation. The differences between the corpus luteum of the unimpregnated and pregnant condition is expressed in the following table by Dalton :—

	Corpus Luteum of Menstruation.	Corpus Luteum of Pregnancy.
At the end of three weeks.	Three-quarters of an inch in diameter; central clot reddish; convoluted wall pale.	
One month.	Smaller; convoluted wall bright yellow; clot still reddish.	Larger; convoluted wall bright yellow; clot still reddish.
Two months.	Reduced to the condition of an insignificant cicatrix.	Seven-eighths of an inch in diameter; convoluted wall bright yellow; clot perfectly decolorized.
Four months.	Absent or unnoticeable.	Seven-eighths of an inch in diameter; clot pale and fibrinous; convoluted wall dull yellow.
Six months.	Absent.	Still as large as at the end of second month; clot fibrinous; convoluted wall paler.
Nine months.	Absent.	Half an inch in diameter; central clot converted into a radiating cicatrix; external wall tolerably thick and convoluted, but without any bright yellow color.

GENERATIVE ORGANS OF THE MALE.

The Generative Organs of the Male consist of the testicles, vasa deferentia, vesiculæ seminales and penis.

The Testicles, the essential organs of reproduction in the male, are two oblong glands, about an inch and a half in length, compressed from side to side, and situated in the cavity of the scrotum.

The proper coat of the testicle, the *tunica albuginea*, is a white, fibrous structure, about the $\frac{1}{25}$ of an inch in thickness; after enveloping the testicle, it is reflected into its interior at the posterior border, and forms a vertical process, the *mediastinum testes*, from which septa are given off, dividing the testicle in lobules.

The substance of the testicle is made up of the *seminiferous tubules*, which exist to the number of 840; they are exceedingly convoluted, and when unraveled, are about 30 inches in length. As they pass toward the apices of the lobules they become less convoluted and terminate in from 20 to 30 straight ducts, the *vasa recta*, which pass upward through the *mediastinum* and constitute the *rete testis*. At the upper part of the mediastinum the tubules unite to form from 9 to 30 small ducts, the *vasa efferentia*, which become convoluted, and form the *globus major* of the *epididymis*; the continuation of the tubes downward behind the testicle and a second convolution constitutes the *body* and *globus minor*.

The seminal tubule consists of a basement membrane lined by granular nucleated epithelium.

The Vas Deferens, the excretory duct of the testicle, is about two feet in length, and may be traced upward from the epididymis to the under surface of the base of the bladder, where it unites with the duct of the vesiculæ seminales, to form the ejaculatory duct.

The Vesiculæ Seminales are two lobulated, pyriform bodies, about two inches in length, situated on the under surface of the bladder.

They have an external fibrous coat, a middle muscular coat, and an internal mucous coat, covered by epithelium, which secretes a mucous fluid. The vesiculæ seminales serve as reservoirs, in which the seminal fluid is temporarily stored up.

The Ejaculatory Duct, about $\frac{3}{4}$ of an inch in length, opens into the urethra, and is formed by the union of the vasa deferentia and the ducts of the vesiculæ seminales.

The Prostate Gland surrounds the posterior extremity of the urethra, and opens into it by from twenty to thirty openings, the orifices of the *pros-*

tatic tubules. The gland secretes a fluid which forms part of the semen, and assists in maintaining the vitality of the spermatozoa.

Semen is a complex fluid, made up of the secretions from the testicles, the vesiculæ seminales, the prostatic and urethral glands. It is grayish-white in color, mucilaginous in consistence, of a characteristic odor, and somewhat heavier than water. From half a drachm to a drachm is ejaculated at each orgasm.

The Spermatozoa are peculiar anatomical elements, developed within the seminal tubules, and possess the power of spontaneous movement. The spermatozoa consist of a conoidal head and a long filamentous tail, which is in continuous and active motion; as long as they remain in the vas deferens they are quiescent, but when free to move in the fluid of the vesiculæ seminales, become very active.

Origin. The spermatozoa appear at the age of puberty, and are then constantly formed until an advanced age. They are developed from the nuclei of large, round cells contained in the interior of the seminal tubules, as many as fifteen to twenty developing in a single cell.

When the spermatozoa are introduced into the vagina, they pass readily into the uterus and through the Fallopian tubes toward the ovaries, where they remain and retain their vitality for a period of from 8 to 10 days.

Fecundation is the union of the spermatozoa with the ovum during its passage toward the uterus, and usually takes place in the Fallopian tube, just outside of the womb. After floating around the ovum in an active manner, they penetrate the vitelline membrane, pass into the interior of the vitellus, where they lose their vitality, and along with the germinal vesicle entirely disappear.

DEVELOPMENT OF ACCESSORY STRUCTURES.

Segmentation of the Vitellus. After the disappearance of the spermatozoa and the germinal vesicle there remains a transparent, granular, albuminous substance, in the centre of which a new nucleus soon appears; this constitutes the *parent cell*, and is the first stage in the development of the new being.

Following this, the vitellus undergoes *segmentation*; a constriction appears on the opposite sides of the vitellus, which gradually deepens, until the yolk is divided into two segments, each of which has a distinct nucleus and nucleolus; these two segments undergo a further division into four, the four into eight, the eight into sixteen, and so on, until the entire

vitellus is divided into a great number of cells, each of which contains a nucleus and nucleolus.

The peripheral cells of this "mulberry mass" then arrange themselves so as to form a membrane, and as they are subjected to mutual pressure, assume a polyhedral shape, which gives to the membrane a mosaic appearance. The central part of the vitellus becomes filled with a clear fluid. A secondary membrane shortly appears within the first, and the two together constitute the external and internal blastodermic membranes.

Germinal Area. At about this period there is an accumulation of cells at a certain spot upon the surface of the blastodermic membranes, which marks the position of the future embryo. This spot, at first circular, soon becomes elongated, and forms the *primitive trace*, around which is a clear space, the *area pellucida*, which is itself surrounded by a darker region, the *area opaca*.

The primitive trace soon disappears, and the *area pellucida* becomes guitar-shaped; a new groove, the *medullary groove*, is now formed, which develops from before backward, and becomes the neural canal.

Blastodermic Membranes. The embryo, at this period, consists of three layers, viz.: the external and internal blastodermic membranes, and a middle membrane formed by a genesis of cells from their internal surfaces. These layers are known as the epiblast, mesoblast and hypoblast.

The *Epiblast* gives rise to the central nervous system, the epidermis of the skin and its appendages, and the primitive kidneys.

The *Mesoblast* gives rise to the dermis, muscles, bones, nerves, blood vessels, sympathetic nervous system, connective tissue, the urinary and reproductive apparatus and the walls of the alimentary canal.

The *Hypoblast* gives rise to the epithelial lining of the alimentary canal and its glandular appendages, the liver and pancreas, and the epithelium of the respiratory tract.

Dorsal Laminæ. As development advances, the true medullary groove deepens, and there arise two longitudinal elevations of the epiblast, the *dorsal laminæ*, one on either side of the groove, which grow up, arch over and unite so as to form a closed tube, the primitive central nervous system.

The *Chorda Dorsalis* is a cylindrical rod running almost throughout the entire length of the embryo. It is formed by an aggregation of mesoblastic cells, and situated immediately beneath the medullary groove.

Primitive Vertebrae. On either side of the neural canal the cells of the mesoblast undergo a longitudinal thickening, which develops and

extends around the neural canal and the chorda dorsalis, and forms the arches and bodies of the vertebrae. They become divided transversely into four-sided segments.

The *Mesoblast* now separates into two layers; the external, joining with the epiblast, forms the *somatopleure*; the internal, joining with the hypoblast, forms the *splanchnopleure*; the space between them constituting the *pleuro-peritoneal cavity*.

Visceral Laminae. The walls of the pleuro-peritoneal cavity are formed by a downward prolongation of the somatopleure (the *visceral laminae*), which, as they extend around in front, pinch off a portion of the yolk sac (formed by the splanchnopleure), which becomes the primitive alimentary canal; the lower portion, remaining outside of the body cavity, forms the *umbilical vesicle*, which after a time disappears.

Formation of Fœtal Membranes. The Amnion appears shortly after the embryo begins to develop, and is formed by folds of the epiblast and external layer of the mesoblast, rising up in front and behind, and on each side; these amniotic folds gradually extend over the back of the embryo to a certain point, where they coalesce, and enclose a cavity, the amniotic cavity. The membranous partition between the folds disappears, and the outer layer recedes and becomes blended with the vitelline membrane, constituting the *chorion*, the external covering of the embryo.

The Allantois. As the amnion develops, there grows out from the posterior portion of the alimentary canal a pouch, or diverticulum, the *allantois*, which carries blood vessels derived from the intestinal circulation. As it gradually enlarges, it becomes more vascular, and inserts itself between the two layers of the amnion, coming into intimate contact with the external layer. Finally, from increased growth, it completely surrounds the embryo, and its edges become fused together.

In the bird, the allantois is a respiratory organ, absorbing oxygen and exhaling carbonic acid; it also absorbs nutritious matter from the interior of the egg.

Amniotic fluid. The amnion, when first formed, is in close contact with the surface of the ovum; but it soon enlarges, and becomes filled with a clear, transparent fluid, containing albumen, glucose, fatty matters, urea and inorganic salts. It increases in amount up to the latter period of gestation, when it amounts to about two pints. In the space between the amnion and allantois is a gelatinous material, which is encroached upon, and finally disappears as the amnion and allantois come in contact, at about the fifth month.

The Chorion, the external investment of the embryo, is formed by a fusion of the original vitelline membrane, the external layer of the amnion, and the allantois. The external surface now becomes covered with villous processes, which increase in number and size by the continual budding and growth of club-shaped processes from the main stem, and give to the chorion a shaggy appearance. They consist of a homogeneous granular matter, and are penetrated by branches of the blood vessels derived from the aorta.

The presence of villous processes in the uterine cavity is proof positive of the previous existence of a foetus. They are characteristic of the chorion, and are found under no other circumstances.

At about the end of the second month the villositities begin to atrophy and disappear from the surface of the chorion, with the exception of those situated at the points of entrance of the foetal blood vessels, which occupy about one-third of its surface, where they continue to grow longer, become more vascular, and ultimately assist in the formation of the placenta; the remaining two thirds of the surface loses its villi and blood vessels, and becomes a simple membrane.

The Umbilical Cord connects the foetus with that portion of the chorion which forms the foetal side of the placenta. It is a process of the allantois, and contains two arteries and a vein, which have a more or less spiral direction. It appears at the end of the first month, and gradually increases in length, until at the end of gestation it measures about 20 inches. The cord is also surrounded by a process of the amnion.

Development of the Decidual Membrane. The interior of the uterus is lined by a thin, delicate mucous membrane, in which are imbedded immense numbers of tubules, terminating in blind extremities, the *uterine tubules*. At each period of menstruation the mucous membrane becomes thickened and vascular, which condition, however, disappears after the usual menstrual discharge. When the ovum becomes fecundated, the mucous membrane takes on an increased growth, becomes more hypertrophied and vascular, sends up little processes, or elevations from its surface, and constitutes the *decidua vera*.

As the ovum passes from the Fallopian tube into the interior of the uterus, the primitive vitelline membrane, covered with villositities, becomes entangled with the processes of the mucous membrane. A portion of the decidua vera then grows up on all sides, and encloses the ovum, forming the decidua reflexa, while the villous processes of the chorion insert themselves into the uterine tubules, and in the mucous membrane between them.

As development advances the decidua reflexa increases in size, and at about the end of the fourth month comes in contact with the decidua vera, with which it is ultimately fused.

The Placenta. Of all the embryonic structures, the placenta is the most important. It begins to be formed toward the end of the second month, and then increases in size until the end of gestation, when it assumes an oval or rounded shape, and measures from 7 to 9 inches in length, 6 to 8 inches in breadth, and weighs from 15 to 20 ounces. It is most frequently situated at the upper and posterior part of the inner surface of the uterus.

The placenta consists of two portions, a foetal and a maternal.

The *Foetal portion* is formed by the villi of the chorion, which, by developing, rapidly increase in size and number. They become branched and penetrate the uterine tubules, which enlarge and receive their many ramifications. The capillary blood vessels in the interior of the villi also enlarge and freely anastomose with each other.

The *Maternal portion* is formed from that part of the hypertrophied and vascular decidual membrane between the ovum and the uterus, the *decidua serotina*. As the placenta increases in size, the maternal blood vessels around the tubules become more and more numerous, and gradually fuse together, forming great lakes, which constitute sinuses in the walls of the uterus.

As the latter period of gestation approaches, the villi extend deeper into the decidua, while the sinuses in the maternal portion become larger and extend further into the chorion. Finally, from excessive development of the blood vessels, the structures between them disappear, and as their walls come in contact, they fuse together, so that, ultimately, the maternal and foetal blood are only separated by a thin layer of a homogeneous substance. When fully formed, the placenta consists principally of blood vessels interlacing in every direction. The blood of the mother passes from the uterine vessels into the lakes surrounding the villi; the blood from the child flows from the umbilical arteries into the interior of the villi; but there is not at any time an intermingling of blood, the two being separated by a delicate membrane formed by a fusion of the walls of the blood vessels and the walls of the villi and uterine sinuses.

The function of the placenta is that of a *respiratory organ*, permitting the oxygen of the maternal blood to pass by osmosis through the delicate placental membrane into the blood of the foetus; at the same time permitting the carbonic acid and other waste products, the result of nutritive changes in the foetus, to pass into the maternal blood, and so to be carried to the various eliminating organs.

Through the placenta also passes all the nutritious materials of the maternal blood which are essential for the development of the embryo.

At about the middle of gestation there develops beneath the decidua membrane a new mucous membrane, destined to perform the functions of the old when it is extruded from the womb, along with the other embryonic structures, during parturition.

DEVELOPMENT OF THE EMBRYO.

Nervous System. The cerebro-spinal axis is formed within the medullary canal by the development of cells from its inner surfaces, which as they increase fill up the canal, and there remains only the central canal of the cord. The external surface gives rise to the dura mater and pia mater. The neural canal thus formed is a tubular membrane; it terminates posteriorly in an oval dilatation, and anteriorly in a bulbous extremity, which soon becomes partially contracted, and forms the anterior, middle and posterior cerebral vesicles, from which are ultimately developed the cerebrum, the corpora quadrigemina, and medulla oblongata, respectively.

The *anterior vesicle* soon subdivides into two secondary vesicles, the larger of which becomes the hemispheres, the smaller, the optic thalami; the *posterior vesicle* also divides into two; the anterior becoming the cerebellum, the posterior, the pons Varolii and medulla oblongata.

About the seventh week the straight chain of cerebral vesicles becomes curved from behind forward and forms three prominent angles. As development advances, the relative size of the encephalic masses changes. The cerebrum developing more rapidly than the posterior portion of the brain, soon grows backward and arches over the optic thalami and the tubercula quadrigemina; the cerebellum overlaps the medulla oblongata.

The surface of the cerebral hemispheres is at first smooth, but at about the fourth month begins to be marked by the future fissures and convolutions.

The Eye is formed from a little bud projecting from the side of the anterior vesicle. It is at first hollow, but becomes lined with nervous matter, forming the *optic nerve* and *retina*; the remainder of the cavity is occupied by the *vitreous body*. The anterior portion of the pouch becomes invaginated and receives the *crystalline lens*, which is a product of the epiblast, as is also the cornea. The *iris* appears as a circular membrane without a central aperture, about the seventh week; the eyelids are formed between the second and third months.

The Internal Ear is developed from the *auditory vesicle*, budding from the third cerebral vesicle; the membranous vestibule appears first,

and from it diverticula are given off, which become the semi-circular canals and cochlea.

The cavity of the tympanum, the Eustachian tube, and the external auditory canal are the remains of the first branchial cleft; the cavity of this cleft being subdivided into the tympanum and external auditory meatus by the *membrana tympani*.

The Skeleton. The *chorda dorsalis*, the primitive part of the vertebral column, is a cartilaginous rod situated beneath the medullary groove. It is a temporary structure, and disappears as the true bony vertebræ develop. On either side are the quadrate masses of the *mesoblast*, the primitive vertebræ, which send processes upward and around the medullary groove, and downward and around the *chorda dorsalis*, forming in these situations the arches and bodies of the future vertebræ.

More externally the outer layer of the *mesoblast* and *epiblast* arch downward and forward, forming the *ventral laminae*, in which develop the *muscles* and *bones* of the abdominal walls.

The *true cranium* is an anterior development of the vertebral column, and consists of the occipital, parietal and frontal segments, which correspond to the three cerebral vesicles. The *base of the cranium* consists, at this period, of a cartilaginous rod on either side of the anterior extremity of the *chorda dorsalis*, in which three centres of ossification appear, the *basi-occipital*, the *basi-sphenoidal*, and the *pre-sphenoidal*. They ultimately develop into the basilar process of the occipital bone and the body of the sphenoid.

The *entire skeleton* is at first either membranous or cartilaginous. At the beginning of the second month centres of ossification appear in the jaws and clavicle; as development advances, the ossific points in all the future bones extend, until ossification is completed.

The *limbs* develop from four little buds projecting from the sides of the embryo, which, as they increase in length, separate into the thigh, leg and foot, and the arm, forearm and hand; the extremities of the limbs undergo subdivision, to form the fingers and toes.

Face and Visceral Arches. In the facial and cervical regions the visceral laminae send up three processes, the *visceral arches*, separated by clefts, the *visceral clefts*.

The *first*, or the *mandibular arches*, unite in the median line to form the lower jaw, and superiorly form the malleus. A process jutting from its base grows forward, unites with the fronto-nasal process growing from above, and forms the upper jaw. When the superior maxillary processes

fail to unite, there results the *cleft-palate* deformity; if the integument also fails to unite, there results the *hare-lip* deformity. The space above the mandibular arch becomes the mouth.

The *second arch* develops the incus and stapes bones, the styloid process and ligament, and the lesser cornu of the hyoid bone. The cleft between the first and second arches partially closes up, but there remains an opening at the side which becomes the Eustachian tube, tympanic cavity, and external auditory meatus.

The *third arch* develops the body and greater cornu of the hyoid bone.

Alimentary Canal and its Appendages. The alimentary canal is formed by a pinching off of the yolk sac by the visceral plates as they grow downward and forward. It consists of three distinct portions, the fore gut, the hind gut, and the central part, which communicates for some time with the yolk sac. It is at first a straight tube, closed at both extremities, lying just beneath the vertebral column. The canal gradually increases in length, and becomes more or less convoluted; at its anterior portion two pouches appear, which become the cardiac and pyloric extremities of the stomach. At about the seventh week the inferior extremity of the intestine is brought into communication with the exterior, by an opening, the anus. Anteriorly the mouth and pharynx are formed by an involution of epiblast, which deepens until it communicates with the fore gut.

The *Liver* appears as a slight protrusion from the sides of the alimentary canal, about the end of the first month; it grows very rapidly, attains a large size, and almost fills up the abdominal cavity. The hepatic cells are derived from the intestinal epithelium, the vessels and connective tissue from the mesoblast.

The *Pancreas* is formed by the hypoblastic membrane. It originates in two small ducts budding from the duodenum, which divide and subdivide, and develop the glandular structure.

The *Lungs* are developed from the anterior part of the œsophagus. At first a small bud appears, which, as it lengthens, divides into two branches; secondary and tertiary processes are given off these, which form the bronchial tubes and air cells. The lungs originally extended into the abdominal cavity, but became confined to the thorax by the development of the diaphragm.

The *Bladder* is formed by a dilatation of that portion of the allantois remaining within the abdominal cavity. It is at first pear-shaped, and communicates with the intestine, but later becomes separated, and opens

exteriorly by the urethra. It is attached to the abdominal walls by a rounded cord, the urachus, the remains of a portion of the allantois.

Genito-Urinary Apparatus. The *Wolffian bodies* appear about the thirtieth day, as long hollow tubes running along each side of the primitive vertebral column. They are temporary structures, and are sometimes called the primordial kidneys. The Wolffian bodies consist of tubules which run transversely and are lined with epithelium; internally they become invaginated to receive tufts of blood vessels; externally they open into a common excretory duct, the *duct* of the *Wolffian body*, which unites with the duct of the opposite body, and empties into the intestinal canal at a point opposite the allantois. On the outer side of the Wolffian body there appears another duct, the duct of Müller, which also opens into the intestine.

Behind the Wolffian bodies are developed the structures which become either the ovaries or testicles. In the development of the female, the Wolffian bodies and their ducts disappear; the extremities of the Müllerian ducts dilate and form the fimbriated extremity of the Fallopian tubes, while the lower portions coalesce to form the body of the uterus and vagina, which now separate themselves from the intestine.

In the development of the male, the Müllerian ducts atrophy, and the ducts of the Wolffian body ultimately form the epididymis and vas deferens. About the seventh month the testicles begin to descend, and by the ninth month have passed through the abdominal ring into the scrotum.

The *Kidneys* are developed out of the Wolffian bodies. They consist of little pyramidal lobules, composed of tubules which open at the apex into the pelvis. As they pass outward they become convoluted and cup-shaped at their extremities, receive a tuft of blood vessels, and form the Malpighian bodies.

The ureters are developed from the kidneys, and pass downward to be connected with the bladder.

The Circulatory Apparatus assumes three different forms at different periods of life, all having reference to the manner in which the embryo receives nutritious matter and is freed of waste products.

The *Vitelline circulation* appears first and absorbs nutritious material from the vitellus. It is formed by blood vessels which emerge from the body and ramify over a portion of the vitelline membrane, constituting the *area vasculosa*. The heart, lying in the median line, gives off two arches which unite to form the abdominal aorta, from which two large arteries are given off, passing into the vascular area; the venous blood is returned

by veins which enter the heart. These vessels are known as the *omphalo-mesenteric arteries and veins*. The vitelline circulation is of short duration in the mammals, as the supply of nutritious matter in the vitellus soon becomes exhausted.

The *Placental circulation* becomes established when the blood vessels in the allantois enter the villous processes of the chorion and come into close relationship with the maternal blood vessels. This circulation lasts during the whole of intra-uterine life, but gives way at birth to the adult circulation, the change being made possible by the development of the *circulatory apparatus*.

The *Heart* appears as a mass of cells coming off from the anterior portion of the intestine; its central part liquefies, and pulsations soon begin. The heart is at first tubular, receiving posteriorly the venous trunks and giving off anteriorly the arterial trunks. It soon becomes twisted upon itself, so that the two extremities lie upon the same plane.

The heart now consists of a single auricle and a single ventricle. A septum growing from the apex of the ventricle divides it into two cavities, a right and a left. The auricles also become partly separated by a septum which is perforated by the foramen ovale. The arterial trunk becomes separated by a partition, into two canals, which become, ultimately, the aorta and pulmonary artery. The auricles are separated from the ventricles by incomplete septa, through which the blood passes into the ventricles.

Arteries. The aorta arises from the cephalic extremity of the heart and divides into two branches which ascend, one on each side of the intestine, and unite posteriorly to form the main aorta; posteriorly to these first aortic arches four others are developed, so that there are five altogether running along the visceral arches. The two anterior soon disappear. The *third arch* becomes the internal carotid and the external carotid; a part of the *fourth arch*, on the right side, becomes the subclavian artery, and the remainder atrophies and disappears, but on the left side it enlarges and becomes the permanent aorta; the *fifth arch* becomes the pulmonary artery on the left side. The communication between the pulmonary artery and the aorta, the *ductus arteriosus*, disappears at an early period.

Veins. The venous system appears first as two short, transverse veins, the canals of Cuvier, formed by the union of the vertebral veins and the cardinal veins, which empty into the auricle. The inferior vena cava is formed as the kidneys develop, by the union of the renal veins, which, in a short time, receive branches from the lower extremities. The subclavian veins join the jugular as the upper extremities develop. The heart descends in the thorax, and the canals of Cuvier become oblique; they

shortly communicate by a transverse duct, which ultimately becomes the left innominate vein. The left canal of Cuvier atrophies and becomes a fibrous cord. A transverse branch now appears, which carries the blood from the left cardinal vein into the right, and becomes the vena azygos minor; the right cardinal vein becomes the vena azygos major.

Circulation of Blood in the Fœtus. The blood returning from the placenta, after having received oxygen, and being freed from carbonic acid, is carried by the umbilical vein to the under surface of the liver; here a portion of it passes through the *ductus venosus* into the ascending vena cava, while the remainder flows through the liver, and passes into the vena cava by the hepatic veins. When the blood is emptied into the right auricle, it is directed by the Eustachian valve, through the foramen ovale, into the left auricle, thence into the left ventricle, and so into the aorta to all parts of the system. The venous blood returning from the head and upper extremities is emptied, by the superior vena cava, into the right auricle, from which it passes into the right ventricle, and thence into the pulmonary artery. Owing to the condition of the lung, only a small portion flows through the pulmonary capillaries, the greater part passing through the ductus arteriosus, which opens into the aorta at a point below the origin of the carotid and subclavian arteries. The mixed blood now passes down the aorta to supply the lower extremities, but a portion of it is directed, by the hypogastric arteries, to the placenta, to be again oxygenated.

At birth, the placental circulation gives way to the circulation of the adult. As soon as the child begins to breathe, the lungs expand, blood flows freely through the pulmonary capillaries, and the ductus arteriosus begins to contract. The foramen ovale closes about the tenth day. The umbilical vein, the ductus venosus, and the hypogastric arteries become impervious in several days, and ultimately form rounded cords.

TABLE OF PHYSIOLOGICAL CONSTANTS.

Mean height of male, 5 feet 6½ inches; of female, 5 feet 2 inches.

Mean weight of male, 145 pounds; of female, 121 pounds.

Number of chemical elements in the human body; from 16 to 18.

Number of proximate principles in the human body; about 100.

Amount of water in the body weighing 145 pounds; 108 pounds.

Amount of solids in the body weighing 145 pounds; 36 pounds.

Amount of food required daily; 16 ounces meat, 19 ounces of bread, 3½ ounces of fat, 52 ounces of water.

Amount of saliva secreted in 24 hours; about 3½ pounds.

Function of saliva; converts starch into glucose.

Active principle of saliva; **ptyalin**.

Amount of gastric juice secreted in 24 hours; from 8 to 14 pounds.

Functions of gastric juice; converts albumen into albuminose.

Active principles of gastric juice; **pepsin** and **hydrochloric acid**.

Duration of digestion; from 3 to 5 hours.

Amount of intestinal juice secreted in 24 hours; about 1 pound.

Function of intestinal juice; converts starch into glucose.

Amount of pancreatic juice secreted in 24 hours; about 1½ pounds.

Active principles of pancreatic juice; **pancreatin** and **trypsin**.

Functions: { 1. Emulsifies fats.
2. Converts albumen into albuminose.
3. Converts starch into glucose.

Amount of bile poured into the intestines daily; about 2½ pounds.

Functions: { 1. Assists in the emulsification of fats.
2. Stimulates the peristaltic movements.
3. Prevents putrefactive changes in the food.
4. Promotes the absorption of the fat.

Amount of blood in the body; from 16 to 18 pounds.

Size of red corpuscles; $\frac{1}{3200}$ of an inch.

Size of white corpuscles; $\frac{1}{2500}$ of an inch.

Shape of red corpuscles; circular biconcave disks.

Shape of white corpuscles; globular.

Number of red corpuscles in a cubic millimetre of blood (the cubic $\frac{1}{25}$ of an inch); 5,000,000.

Function of **red corpuscles** ; to carry oxygen from the lungs to the tissues.

Frequency of the **heart's pulsations** per minute ; 72, on the average.

Velocity of the **blood movement** in the arteries ; about 16 inches per second.

Length of time required for the **blood** to make an entire circuit of the vascular system ; about 20 seconds.

Amount of **air** passing in and out of the lungs at each respiratory act ; from 20 to 30 cubic inches.

Amount of **air** that can be taken into the lungs on a forced inspiration ; 110 cubic inches.

Amount of **reserve air** in the lungs after an ordinary expiration ; 100 cubic inches.

Amount of **residual air** always remaining in the lungs ; about 100 cubic inches.

Vital capacity of the **lungs** ; 230 cubic inches.

Entire volume of **air** passing in and out of the lungs in 24 hours ; about 400 cubic feet.

Composition of the **air** ; nitrogen, 79.19, oxygen, 20.81, per 100 parts.

Amount of **oxygen** absorbed in 24 hours ; 18 cubic feet.

Amount of **carbonic acid** exhaled in 24 hours ; 14 cubic feet.

Temperature of the **human body** at the surface ; $98\frac{5}{10}^{\circ}$ F.

Amount of **urine** excreted daily ; from 40 to 50 ounces.

Amount of **urea** excreted daily ; 512 grains.

Specific gravity of **urine** ; from 1.010 to 1.015.

Number of **spinal nerves** ; 31 pairs.

Number of **roots of origin** ; two ; 1st, anterior, motor ; 2d, posterior, sensory.

Rate of transmission of **nerve force** ; about 100 feet per second.

Number of **cranial nerves** ; 12 pairs.

Nerves of special sense :	{	1. Olfactory, or 1st pair.
		2. Optic, or 2d pair.
		3. Auditory, or 8th pair.
		4. Chorda tympani for anterior $\frac{2}{3}$ of tongue.
		5. Branches of glosso-pharyngeal, or 8th pair, for posterior $\frac{1}{3}$ of tongue.

Motor nerves to **eyeball** and **accessory structures** ; motor oculi, or 3d pair ; pathetic, or 4th pair ; abducens, or 6th pair.

Motor nerves to **facial muscles** ; portio dura, facial, or 7th pair.

Motor nerve to **tongue** ; hypoglossal or 12th pair.

Motor nerve to **laryngeal muscles** ; spinal accessory or 11th pair.

- Sensory nerve of the **face** ; trifacial or 5th pair.
- Sensory nerve of the **pharynx** ; glosso-pharyngeal or 9th pair.
- Sensory nerve of the **lungs, stomach, etc.** ; pneumogastric or 10th pair.
- Length of **spinal cord** ; 16 to 18 inches, weight $1\frac{1}{2}$ ounces.
- Point of decussation of **motor fibres** ; at the medulla oblongata.
- Point of decussation of **sensory fibres** ; throughout the spinal cord.
- Function of **antero-lateral columns of spinal cord** ; transmit motor impulses from the brain to the muscles.
- Functions of the **posterior columns** ; assist in the coördination of muscular movements.
- Functions of the **medulla oblongata** ; controls the functions of insalivation, mastication, deglutition, respiration, circulation, etc.
- Functions of the **corpora quadrigemina** ; physical centres for sight.
- Functions of the **corpora striata** ; centres for motion.
- Functions of the **optic thalami** ; centres for sensation.
- Function of the **cerebellum** ; centre for the coördination of muscular movements.
- Function of the **cerebrum** ; centre for intelligence, reason and will.
- Centre for **articulate language** ; 3d frontal convolution on left side of cerebrum.
- Number of coats to the **eye** ; three ; 1st, cornea and sclerotic ; 2d, choroid ; 3d, retina.
- Function of **iris** ; regulates the amount of light entering the eye.
- Function of **crystalline lens** ; refracts the rays of light so as to form an image on the retina.
- Function of **retina** ; receives the impressions of light.
- Function of **membrana tympani** ; receives and transmits waves of sound to internal ear.
- Function of **Eustachian tube** ; regulates the passage of air into and from the middle ear.
- Function of **semi-circular canals** ; assist in maintaining the equipoise of the body.
- Function of the **cochlea** ; appreciates the shades and combinations of musical tones.
- Size of **human ovum** ; $\frac{1}{125}$ of an inch in diameter.
- Size of **spermatozoa** ; $\frac{1}{4000}$ of an inch in length.
- Function of the **placenta** ; acts as a respiratory and digestive organ for the foetus.
- Duration of **pregnancy** ; 280 days.

TABLE SHOWING RELATION OF WEIGHTS AND MEASURES OF THE METRIC SYSTEM TO APPROXIMATE WEIGHTS AND MEASURES OF THE U. S.

MEASURES OF LENGTH.

One Myriametre	= 10,000 metres	= 32800. feet.
One Kilometre	= 1,000 "	= 3280. "
One Hectometre	= 100 "	= 328.0 "
One Decametre	= 10 "	= 32.80 "
One METRE	= { the ten millionth part of a quarter of the Meridian of the Earth }	= 39.368 inches.
One Decimetre	= the tenth part of one metre	= 3.936 "
One Centimetre	= { the one hundredth part of one metre }	= .393 ($\frac{2}{5}$) "
One Millimetre	= { the one thousandth part of one metre }	= .039 ($\frac{1}{25}$) "

WEIGHTS.

One Myriagramme	= 10,000 grammes	= 26 $\frac{3}{4}$ pounds Troy.
One Kilogramme	= 1,000 "	= 2 $\frac{2}{3}$ " "
One Hectogramme	= 100 "	= 3 $\frac{1}{4}$ ounces "
One Decagramme	= 10 "	= 2 $\frac{1}{2}$ drachms "
One GRAMME	= { the weight of a cubic centi- metre of water at 4° C. }	= 15.434 grains.
One Decigramme	= the tenth part of one gramme	= 1.543 ($1\frac{1}{2}$) "
One Centigramme	= { the hundredth part of one gramme }	= .154 ($\frac{1}{6}$) "
One Milligramme	= { the thousandth part of one gramme }	= .015 ($\frac{1}{64}$) "

MEASURES OF CAPACITY.

One Myrialitre	= { 10 cubic Metres or the measure of 10 Milliers of water }	= 2600. gallons.
One Kilolitre	= { 1 cubic Metre or the meas- ure of 1 Millier of water }	= 260. "
One Hectolitre	= { 100 cubic Decimetres or the measure of 1 Quintal of water }	= 26. "
One Decalitre	= { 10 cubic Decimetres or the measure of 1 Myriagramme of water }	= 2.6 "
One LITRE	= { 1 cubic Decimetre or the measure of 1 Kilogramme of water }	= 2.1 pints.
One Decilitre	= { 100 cubic Centimetres or the measure of 1 Hecto- gramme of water }	= 3.3 ounces.
One Centilitre	= { 10 cubic Centimetres or the measure of one Deca- gramme of water }	= 2.7 drachms.
One Millilitre	= { 1 cubic Centimetre or the measure of 1 gramme of water }	= 16.2 minims.

INDEX.

	PAGE
ABDUCENS NERVE	90
— Aberration, chromatic.....	135
— spherical.....	135
Absorption.....	33
— by the lacteals.....	38
— by the blood vessels.....	34
— of oxygen in respiration.....	56
Accommodation of the eye.....	135
Adipose tissue, uses of in the body....	12
Adult circulation, establishment of at birth.....	157
Air, atmospheric, composition of.....	56
— amount exchanged in respiration.....	56
— changes in during respiration....	57
Albumen, uses of in the body.....	13
Albuminoid substances.....	13
Alcohol, action of.....	21
Alimentary principles, classification of.....	19
— albuminous principles.....	19
— saccharine principles.....	19
— oleaginous principles.....	19
— inorganic principles.....	20
Alimentary canal, development of.....	155
Allantois, development and function of.....	150
Amnion, formation of.....	150
Animal heat.....	53
Anterior columns of spinal cord.....	105
Area, germinal.....	149
Arteries, properties of.....	49
Asphyxia.....	58
Astigmatism.....	135
Axis, cerebro-spinal.....	100
— cylinder of nerves.....	81
BILE	75
— Bladder, urinary.....	69
Blastodermic membranes.....	148
Blood.....	39
— composition of plasma.....	40
— coagulation of.....	43
— coloring matter of.....	34
— changes in, during respiration.....	56
— circulation of.....	40
— rapidity of flow in arteries.....	55
— rapidity of flow in capillaries....	51
— pathological conditions of.....	45
— corpuscles.....	41
— origin of.....	42
— pressure.....	49
Burdach, column of.....	102

	PAGE
CANALS OF CUVIER	157
— Capillary blood vessels.....	50
Capsule, internal.....	115
— external.....	115
Caudate nucleus.....	115
Cells, structure of.....	16
— manifestations of life by.....	16
— of anterior horns of gray matter.....	104
Centre for articulate language.....	121
Cerebrum.....	113
— fissures and convolutions.....	183
— functions of.....	120
— localization of functions.....	121
— motor area of.....	122
— special centres of.....	122
Cerebellum.....	116
— forced movements of.....	117
Cerebral vesicles of embryo.....	153
Chemical composition of human body.....	10
— elements, proximate quantity of in body.....	15
Chorda dorsalis.....	149
— tympani nerve, course and function of.....	93
Chorion.....	151
Chyle.....	38
Ciliary muscle.....	131
Circulation of blood.....	45
Clastrum.....	115
Cochlea.....	139
Columns of spinal cord.....	101
Corium.....	85
Corpora Wolffiana.....	156
— quadrigemina.....	114
Corpus luteum.....	145
— striatum.....	114
Corti, organ of.....	140
Cranial nerves.....	87
Crura cerebri.....	113
Crystalline lens.....	132
DECIDUAL MEMBRANE	151
— Decussation of motor and sensory fibres.....	104
Deglutition.....	26
— nervous circle of.....	111
Development of accessory structures of embryo.....	148
Digestion.....	21
Ductus arteriosus.....	153
— venosus.....	158

	PAGE		PAGE
EAR	136	Heart, work done by.....	47
— Electrotonus	86	— course of blood through.....	46
Embryo, development of.....	153	— influence of nervous system	
Endolymph.....	139	upon.....	48
Epidermis.....	85	Hyaloid membrane.....	132
Epididymis.....	147	Hypermetropia.....	135
Epiglottis.....	27	Hypoglossal nerve.....	99
Eustachian tube.....	138		
Excretion	66	INCUS BONE	137
Eye.....	130	— Ingesta and egesta, comparison of	
— refracting apparatus of.....	133	in 24 hours.....	22
— blind spot of.....	134	Insalivation.....	24
		— nervous circle of.....	25
FACIAL NERVE	92	Inspiration, movements of thorax in.....	54
— paralysis, symptoms of ..	93	Internal capsule.....	115
Fallopian tubes.....	144	— results of injury to.....	115
Facies.....	33	Intestinal juice.....	30
Fat, uses of in the body.....	13	Iris.....	130
Female organs of generation.....	144	— action of.....	134
Fissures and convolutions of brain.....	118	Island of Reil.....	119
Food.....	18		
— percentage composition of.....	22	KIDNEYS	66
— daily amount required.....	22	— excretion of urine by.....	70
— albuminous principles of.....	20		
— saccharine principles of.....	20	LABYRINTH OF INTERNAL	
— oleaginous principles of.....	20	ear	138
— inorganic principles of.....	19	— function of cochlea.....	141
Fovea centralis.....	134	— function of semicircular canals.....	140
		Language, articulate, centre for.....	121
GALVANIC CURRENTS, EF-		Larynx.....	52
fect on nerves.....	86	Lateral columns of spinal cord.....	101
Ganglia.....	80	Laws of muscular contraction.....	87
— ophthalmic.....	123	Latator tympani muscle.....	138
— Gasserian.....	91	Lens, crys-talline.....	132
— sphenopalatine.....	122	Lime phosphate.....	12
— otic.....	123	Liver.....	73
— sub-maxillary.....	123	— secretion of bile by.....	75
— semi-lunar.....	124	— glyco-genic function of.....	75
Gases of the intestine.....	35	— elaboration of blood.....	75
— condition of, in blood.....	57	— cells.....	74
Gastric juice.....	28	Localization of functions in cerebrum.....	121
— action of.....	29	Lungs.....	53
Generation, male organs of.....	147	— changes in blood while passing	
— female organs of.....	144	through.....	57
Globules of the blood.....	41	Lymph.....	36
— of the lymph.....	36	Lymphatic glands.....	36
Glomeruli of the kidneys.....	69	— vessels, origin and course of... ..	35
Glossopharyngeal nerve.....	95		
Glottis, respiratory movements of	54	MAMMARY GLANDS	63
Glycogen.....	76	— Malleus bone.....	137
Glyco-genic function of liver.....	75	Mastication.....	25
Golf, column of.....	102	— nervous circle of.....	25
Graafian follicles.....	144	— muscles of.....	24
Gray matter of nervous system.....	80	Medulla oblongata.....	109
		— properties and functions of.....	110
HAIR	77	Membrana basilaris.....	140
— Hæmoglobin.....	42	— tympani.....	137
Hearing, sense of.....	136	Menstruation.....	145
Heart.....	45	Middle ear.....	137
— valves of.....	45	Milk.....	63
— sounds of.....	46	Motor centres of cerebrum.....	122
— influence of pneumogastric		Muscles, properties of.....	35
nerve upon.....	48	Myopia.....	135
— ganglia of.....	48		
— force exerted by left ventricle..	47		

	PAGE		PAGE
NERVE, OLFACTORY	88	Proximate quantity of chemical ele-	
— optic.....	88	ments in body.....	15
— motor oculi.....	89	Ptyadin.....	25
— pathetic.....	90	Pulse.....	49
— trigeminal.....	90	Pyramidal tracts.....	101
— abducens.....	90	RED CORPUSCLES OF	
— facial.....	92	blood.....	41
— auditory.....	94	Reflex movements of spinal cord.....	106
— glosso-pharyngeal.....	95	— action, laws of.....	107
— pneumogastric.....	95	Reproduction.....	144
— spinal accessory.....	96	Respiration.....	52
— hypoglossal.....	99	— movements of.....	54
— cells, structure of.....	80	— nervous mechanism of.....	55
— fibres, terminations of.....	83	— types of.....	55
— force, rate of transmission of...	87	— nervous circle of.....	112
— roots, function of anterior and		Retina.....	134
posterior.....	82	SALIVA	25
Nerves, centrifugal and centripetal....	84	Sebaceous glands.....	78
— cranial.....	87	Secretion.....	60
— decussation of motor and sensory		Semi-circular canals.....	139
vaso-motor.....	104	Semen.....	148
— properties and functions of.....	84	Sight, sense of.....	130
— spinal.....	84	Skin.....	76
Nervous system.....	80	— relative sensibility of.....	126
— white and gray matter of.....	80	— appendages of.....	77
— cerebro-spinal.....	80	Smell, sense of.....	129
— sympathetic.....	123	Sounds of heart.....	46
Nucleus caudatus.....	115	Spermatozoa.....	148
— lenticularis.....	115	Spheno-palatine ganglion.....	122
OLFACTORY NERVES	88	Spinal accessory nerve.....	98
Ophthalmic ganglion.....	123	Spinal cord.....	101
Optic nerves.....	88	— membranes of.....	100
— thalamus.....	115	— structure of white matter.....	101
— function of.....	115	— structure of gray matter.....	102
Organs of Corti.....	140	— properties of.....	104
Otic ganglion.....	123	— function of as a conductor.....	105
Ovaries.....	144	— as an independent centre.....	106
Ovum.....	144	— decussation of motor and sensory	
— discharge of from the ovary.....	145	fibres.....	104
Oxygen, absorption of by hæmoglobin		— reflex action of.....	106
42		— special centres of.....	106
PACINIAN CORPUSCLES	83	— paralysis, from disease of.....	108
Pancreatic juice.....	30	— nerves, origin of.....	102
Patheticus nerve.....	90	— course of anterior and posterior	
Peptones.....	29	roots of.....	102
Perilymph.....	139	Spleen.....	65
Perspiration.....	69	Stapes bone.....	137
Petrosal nerves, large and small.....	93	Starvation, phenomena of.....	18
Phonation.....	142	Stomach.....	26
Physiology, definition of.....	9	Structural composition of the body...	15
Placenta, formation and function of...	152	Submaxillary ganglion.....	123
Pneumogastric nerve.....	95	Sugar, uses of in the body.....	12
Pons varolii.....	112	Supra-renal capsules.....	66
Portal vein.....	34	Sudoriparous glands.....	78
Posterior columns of spinal cord.....	105	Sympathetic nervous system.....	123
— functions of.....	105	— properties and functions of.....	124
Prehension.....	23	TASTE, SENSE OF	127
Presbyopia.....	135	— nerve of.....	128
Pressure of blood in arteries.....	49	Teeth.....	24
Proximate principles.....	11	Tensor tympani muscle.....	137
— inorganic.....	11	Testicles.....	147
— organic, non-nitrogenized.....	12	Thoracic duct.....	36
— organic, nitrogenized.....	13	Thorax, enlargement of in inspiration	54
— of waste.....	15		

	PAGE		PAGE
Tissues, classification of.....	17	V APOR, WATERY, OF	
Tongue.....	127	breath.....	57
—— motor nerve of.....	128	Vascular glands.....	65
—— sensory nerve of.....	128	—— system, development of.....	156
Touch, sense of.....	126	Vaso-motor nerves, origin of.....	111
Türck column.....	101	Veins.....	51
U MBILICAL CORD.....	151	Vesiculæ seminales.....	147
Urea.....	72	Vision, psychical centre for.....	122
Uric acid.....	72	—— physical centre for.....	114
Urine.....	70	Vital capacity of lungs.....	56
—— composition of.....	71	Vocal cords.....	141
—— average quantity of constitu-		Voice.....	141
ents secreted daily.....	71	W ATER, AMOUNT OF IN	
Urination, nervous mechanism of.....	70	body.....	11
Uterus.....	145	Wolffian bodies.....	156

CATALOGUE No. 7. *

FEBRUARY, 1886.



A CATALOGUE
OF
BOOKS FOR STUDENTS;

INCLUDING A FULL LIST OF

The ?Quiz-Compend?,
MANUALS,
Text-Books and Students' Aids,

PUBLISHED BY

P. BLAKISTON, SON & CO.

Medical Booksellers, Importers and Publishers.

LARGE STOCK OF ALL STUDENTS' BOOKS, AT
THE LOWEST PRICES.

No. 1012 WALNUT STREET,

PHILADELPHIA.

*** For sale by all Booksellers, or any book will be sent by mail, postpaid, upon receipt of price. Catalogues of books on all branches of Medicine, Dentistry, Pharmacy, etc., supplied upon application.

? QUIZ-COMPENDS?

A NEW SERIES OF COMPENDS FOR STUDENTS

For Use in the Quiz Class and when
Preparing for Examinations.

Price of Each, Bound in Cloth, \$1.00 Interleaved, \$1.25.

Based on the most popular text-books, and on the lectures of prominent professors, they form a most complete set of manuals, containing information nowhere else collected in such a condensed, practical shape. The authors have had large experience as quiz masters and attachés of colleges, with exceptional opportunities for noting the most recent advances and methods. The arrangement of the subjects, illustrations, types, etc., are all of the most improved form, and the size of the books is such that they may be easily carried in the pocket. They are constantly revised, so as to include the latest and best teachings, and can be used by students of any college.

Nos. 1. ANATOMY. (Illustrated.)

THIRD REVISED EDITION.

A Compend of Human Anatomy. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army. With 63 Illustrations.

"The work is reliable and complete, and just what the student needs in reviewing the subject for his examinations."—*The Physician and Surgeon's Investigator*, Buffalo, N. Y.

"The arrangement is well calculated to facilitate accurate memorizing, and the illustrations are clear and good."—*North Carolina Medical Journal*.

Nos. 2 and 3. PRACTICE.

NEW REVISED EDITIONS.

A Compend of the Practice of Medicine, especially adapted to the use of Students. By DAN'L E. HUGHES, M.D., Demonstrator of Clinical Medicine in Jefferson Medical College, Philadelphia. Second Edition. Enlarged and thoroughly Revised. In two parts.

PART I.—Continued, Eruptive, and Periodical Fevers, Diseases of the Mouth, Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, Intestinal Parasites, etc., and General Diseases.

PART II.—Diseases of the Respiratory System, Circulatory System and Blood, Nervous System, etc.

Price of each Book, Cloth, \$1.00. Interleaved for Notes, \$1.25.

* * These little books can be regarded as a full set of notes upon the Practice of Medicine, containing the Synonyms, Definitions, Causes, Symptoms, Prognosis, Diagnosis, Treatment, etc., of each disease, and including a number of new prescriptions. They have been compiled from the lectures of prominent Professors, and reference has been made to the latest writings of Professors FLINT, DA COSTA, BARTHOLOW, ROBERTS, etc.

"It is brief and concise, and at the same time possesses an accuracy not generally found in compends."—*Jas. M. French, M.D., Ass't to the Prof. of Practice, Medical College of Ohio, Cincinnati.*

"The book seems very concise, yet very comprehensive. . . . An unusually superior book."—*Dr. E. T. Bruen, Demonstrator of Clinical Medicine, University of Pennsylvania.*

"I have used it considerably in connection with my branches in the Quiz-class of the University of La."—*J. H. Bemiss.*

"Dr. Hughes has prepared a very useful little book, and I shall take pleasure in advising my class to use it."—*Dr. George W. Hall, Prof. of Practice, St. Louis College of Phys. and Surgeons.*

No. 4. PHYSIOLOGY. Illustrated.

THIRD REVISED EDITION.

Compend of Human Physiology, adapted to the use of Students. By ALBERT P. BRUBAKER, M.D., Demonstrator of Physiology in Jefferson Medical College, Philadelphia. Third Ed. Enlarged and Revised.

"Dr. Brubaker deserves the hearty thanks of medical students for his *Compend of Physiology*. He has arranged the fundamental and practical principles of the science in a peculiarly inviting and accessible manner. I have already introduced the work to my class."—*Maurice N. Miller, M.D., Instructor in Histology, formerly Demonstrator of Physiology, University City of New York.*

"'Quiz-Compend' No. 4 is fully up to the high standard established by its predecessors of the same series."—*Medical Bulletin, Philadelphia.*

"I can recommend it as a valuable aid to the student."—*C. N. Ellinwood, M.D., Professor of Physiology, Cooper Medical College, San Francisco.*

"This is a well written little book."—*London Lancet.*

No. 5. OBSTETRICS. Second Ed.

A Compend of Obstetrics. For Physicians and Students.

By HENRY G. LANDIS, M.D., Professor of Obstetrics and Diseases of Women, in Starling Medical College, Columbus. New Revised Ed. New Illustrations.

"We have no doubt that many students will find in it a most valuable aid in preparing for examination."—*The American Journal of Obstetrics.*

"It is complete, accurate and scientific. The very best book of its kind I have seen."—*J. S. Knox, M.D., Lecturer on Obstetrics, Rush Medical College, Chicago.*

Price of each Book, Cloth, \$1.00. Interleaved for Notes, \$1.25.

"I have been teaching in this department for many years, and am free to say that this will be the best assistant I ever had. It is accurate and comprehensive, but brief and pointed."—*Prof. P. D. Yost, St. Louis.*

No. 6. MATERIA MEDICA. Revised Ed.

A Compend on Materia Medica and Therapeutics, with especial reference to the Physiological Actions of Drugs. For the use of Medical, Dental, and Pharmaceutical Students and Practitioners. Based on the New Revision (Sixth) of the U. S. Pharmacopœia, and including many unofficial remedies. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army.

"I have examined the little volume carefully, and find it just such a book as I require in my private Quiz, and shall certainly recommend it to my classes. Your Compends are all popular here in Washington."—*John E. Brackett, M.D., Professor of Materia Medica and Therapeutics, Howard Medical College, Washington.*

"Part of a series of small but valuable text-books. . . . While the work is, owing to its therapeutic contents, more useful to the medical student, the pharmaceutical student may derive much useful information from it."—*N. Y. Pharmaceutical Record.*

No. 7. CHEMISTRY. Revised Ed.

A Compend of Chemistry. By G. MASON WARD, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. Including Table of Elements and various Analytical Tables.

"Brief, but excellent. . . . It will doubtless prove an admirable aid to the student, by fixing these facts in his memory. It is worthy the study of both medical and pharmaceutical students in this branch."—*Pharmaceutical Record, New York.*

No. 8. VISCERAL ANATOMY.

SECOND EDITION. ILLUSTRATED.

A Compend of Visceral Anatomy. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army. With 40 Illustrations.

. This is the only Compend that contains full descriptions of the viscera, and will, together with No. 1 of this series, form the only complete Compend of Anatomy published.

"This work is very happily arranged, very thorough in practical details, and will no doubt prove universally popular with medical students."—*Medical Herald.*

"I believe it will prove of great usefulness to the busy teacher or student; short, concise helps are always welcome."—*Dr. R. N. Hall, Demonstrator of Anatomy, College of Physicians and Surgeons, Chicago.*

"It is a very concise and convenient help to the memory, and quite accurate."—*Prof. L. B. How, Medical Department, Dartmouth College.*

Price of Each Book, Cloth, \$1.00. Interleaved for Notes, \$1.25.

No. 9. SURGERY. Second Edition.**ILLUSTRATED.**

A Compend of Surgery; including Fractures, Wounds, Dislocations, Sprains, Amputations and other operations, Inflammation, Suppuration, Ulcers, Syphilis, Tumors, Shock, etc. Diseases of the Spine, Ear, Eye, Bladder, Testicles, Anus, and other Surgical Diseases. By ORVILLE HORWITZ, A.M., M.D., with 62 Illustrations. Second Edition. Enlarged and Revised.

. This compend has been prepared with great care, from the standard authorities on Surgery and from notes taken by the author during attendance on lectures by prominent professors. The rapid sale of the first edition allowed the addition of much valuable matter, besides a thorough revision of the whole book.

No. 10. ORGANIC CHEMISTRY.

A Compend of Organic Chemistry, including Medical Chemistry, Urine Analysis, and the Analysis of Water and Food, etc. By HENRY LEFFMANN, M.D., Professor of Clinical Chemistry and Hygiene in the Philadelphia Polyclinic; Professor of Chemistry, Pennsylvania College of Dental Surgery; Member of the N. Y. Medico-Legal Society. Cloth. \$1.00.

Interleaved, for the addition of Notes, \$1.25.

"Compact, substantial and exact; well suited as a remembrancer to students."—*Pacific Medical and Surgical Journal*.

"This neat, handy and exceedingly useful volume is a valuable aid to the student."—*Pharmaceutical Record*.

"It contains, in compact form, the most of modern organic and medical chemistry essential to the student of medicine, and will be of great value in bringing this subject within his grasp."—C. C. Howard, Prof. of Chemistry, Starling Medical College, Columbus, Ohio.

"It has the decided merit of being written in a clear and understandable language."—Dr. J. Sickels, Instructor in Chemistry, University Medical College, New York.

No. 11. PHARMACY.

A Compend of Pharmacy. By F. E. STEWART, M.D., Ph. G., Quiz Master in Chemistry and Theoretical Pharmacy, Philadelphia College of Pharmacy; Demonstrator and Lecturer in Pharmacology, Medico-Chirurgical College; Member of the American Pharmaceutical Association.

✚ The ? Quiz Compend ? are adapted to students of any college, because they are based upon the text-books in use throughout the country. They contain the latest and best information, in such a shape that it can be easily memorized.

Price of Each Book, Cloth, \$1.00; Interleaved for Notes \$1.25.

ANATOMY.

Holden's Anatomy. A manual of Dissection of the Human Body. Fifth Edition. Enlarged, with Marginal References and over 200 Illustrations. Octavo. Cloth, 5.00; Leather, 6.00

Bound in Oilcloth, for the Dissecting Room, \$4.50.

"No student of Anatomy can take up this book without being pleased and instructed. Its Diagrams are original, striking and suggestive, giving more at a glance than pages of text description.

* * * The text matches the illustrations in directness of practical application and clearness of detail."—*New York Medical Record*.

Holden's Human Osteology. Comprising a Description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. With Lithographic Plates and Numerous Illustrations. Sixth Edition. 8vo. Cloth, 6.00

Heath's Practical Anatomy. Sixth London Edition. 24 Colored Plates, and nearly 300 other Illustrations. Cloth, 5.00

CHEMISTRY.

Bartley's Medical Chemistry. A text-book prepared specially for Medical, Pharmaceutical and Dental Students. With 40 Illustrations, Plate of Absorption Spectra and Glossary of Chemical Terms. Cloth, 2.50

* * * This book has been written especially for students and physicians. It is practical and concise, dealing only with those parts of chemistry pertaining to medicine; no time being wasted in long descriptions of substances and theories of interest only to the advanced chemical student.

Bloxam's Chemistry, Inorganic and Organic, with Experiments. Fifth Edition, nearly 300 Illustrations. Cloth, 3.75; Leather, 4.75

Bowman's Practical Chemistry. Including Analysis. About 100 Illustrations. Eighth Edition. Cloth, 2.00

Muter's Practical and Analytical Chemistry. 8vo. Cloth, 2.50

Richter's Inorganic Chemistry. A text-book for Students. Second American, from Fourth German Edition. Translated by Prof. Edgar F. Smith, PH.D. 89 Wood Engravings and Colored Plate of Spectra. *Just Ready.* Cloth, 2.00

Richter's Organic Chemistry, or Chemistry of the Carbon Compounds. Translated by Prof. Edgar F. Smith, PH.D. Illustrated. *Just Ready.* Cloth, 3.00

Trimble. Practical and Analytical Chemistry. A Complete Course in Chemical Analysis, by Henry Trimble, Professor of Analytical Chemistry in the Philadelphia College of Pharmacy. Illustrated. 8vo. Cloth, 1.50

See pages 2 to 5 for list of ? Quiz-Compend ?

Chemistry:—Continued.

Wolff's Applied Medical Chemistry. By Lawrence Wolff, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. *Just Ready.* Cloth, 1.50

CHILDREN.

Goodhart and Starr. The Diseases of Children. A Manual for Students and Physicians. By J. F. Goodhart, M.D., Physician to the Evelina Hospital for Children; Assistant Physician to Guy's Hospital, London. American Edition, Revised and Edited by Louis Starr, M.D., Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania; Physician to the Children's Hospital, Philadelphia. Containing many new Prescriptions, a List of over 50 Formulæ, conforming to the U. S. Pharmacopœia, and Directions for making Artificial Human Milk, for the Artificial Digestion of Milk, etc. *Just Ready.* Demi-Octavo, 738 Pages. Cloth, 3.00; Leather, 4.00

The NEW YORK MEDICAL RECORD says:—"As it is said of some men, so it might be said of some books, that they are 'born to greatness.' This new volume has, we believe, a mission, particularly in the hands of the younger members of the profession. In these days of prolixity in medical literature, it is refreshing to meet with an author who knows both what to say, and when he has said it. The work of Dr. Goodhart (admirably conformed, by Dr. Starr, to meet American requirements) is the nearest approach to clinical teaching, without the actual presence of clinical material, that we have yet seen. The details of management so gratefully read by the young practitioner are fully elucidated. Altogether, the book is one of as great practical working value as we have seen for many months."

Day. On Children. A Practical and Systematic Treatise. Second Edition. 8vo. 752 pages. Cloth, 3.00; Leather, 4.00
Meigs and Pepper. The Diseases of Children. Seventh Edition. 8vo. Cloth, 6.00; Leather, 7.00

DENTISTRY.

Flagg's Plastics and Plastic Filling. 2d Ed. Cloth, 4.00
Gorgas. Dental Medicine. A Manual of Materia Medica and Therapeutics, by Professor F. J. S. Gorgas, M.D., D.D.S., Professor of the Principles and Practice of Dental Science, in Dental Department, University of Maryland. 8vo. Second Edition. Revised and Enlarged. Cloth, 3.25

Harris' Principles and Practice of Dentistry. Including Anatomy, Physiology, Pathology, Therapeutics, Dental Surgery and Mechanism. Eleventh Edition. Revised and enlarged by Professor Gorgas. 744 Illustrations. Cloth, 6.50; Leather, 7.50

Richardson's Mechanical Dentistry. Third Edition. 185 Illustrations. 8vo. Cloth, 4.00; Leather, 4.75

Stocken's Dental Materia Medica. Third Edition. Cloth, 2.50

See pages 2 to 5 for list of Quiz-Compends

Dentistry:—Continued.

Tomes' Dental Anatomy, Human and Comparative. Second Edition. 191 Illustrations. Cloth, 4.25

Tomes' Dental Surgery. New Revised Edition. *Preparing.*

Taft's Operative Dentistry. A Text-book for Dental Students and Practitioners. Fourth Edition. Over 100 Illustrations. Cloth, 4.25; Leather, 5.00

DICTIONARIES.

Cleaveland's Pocket Medical Lexicon. Thirty-first Edition. Giving correct Pronunciation and Definition of Terms used in Medicine and the Collateral Sciences. Very small pocket size, red edges. Cloth, .75; pocket-book style, 1.00

Longley's Pocket Dictionary. The Student's Medical Lexicon, giving Definition and Pronunciation of all Terms used in Medicine, with an Appendix giving Poisons and Their Antidotes, Abbreviations used in Prescriptions, Metric Scale of Doses, etc. 24mo. Cloth, 1.00; pocket-book style, 1.25

Harris' Dictionary of Medical Terminology and Dental Surgery By Prof. Gorgas. Fourth Edition. Cloth, 6.50; Leather, 7.50

EYE.

Arlt. Diseases of the Eye. Including those of the Conjunctiva, Cornea, Sclerotic, Iris and Ciliary Body. By Professor Fred. Ritter von Arlt. Translated by Dr. Lyman Ware. Illustrated. 8vo. Cloth, 2.50

Higgins. Ophthalmic Practice. A Handbook for Students and Practitioners. 16mo. Cloth, .50

Macnamara. On Diseases of the Eye. Fourth Edition, revised, with Marginal References, numerous Colored Plates and Diagrams, Wood Cuts and Test Types. Cloth, 4.00


HYGIENE.

Parke's Practical Hygiene. Sixth Edition, enlarged. Illustrated. 8vo. Cloth, 3.00

Wilson's Handbook of Hygiene and Sanitary Science. Fifth Edition. Revised and Illustrated. Cloth, 2.75

MATERIA MEDICA AND THERAPEUTICS.

Biddle's Materia Medica. Tenth Edition. For the use of Students and Physicians. By the late Prof. John B. Biddle, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Tenth Edition, thoroughly revised, and in many parts rewritten, by his son, Clement Biddle, M.D., Past Assistant Surgeon, U. S. Navy, assisted by Henry Morris, M.D., Demonstrator of Obstetrics in Jefferson Medical College. 8vo., illustrated. *Just Ready.* Cloth, 4.00; Leather, 4.75

 See pages 2 to 5 for list of ? Quia-Compendis?

Biddle's Materia Medica and Therapeutics:—Continued.

"The larger works usually recommended as text-books in our medical schools are too voluminous for convenient use. This work will be found to contain in a condensed form all that is most valuable, and will supply students with a reliable guide."—*Chicago Med. J.*

Merrell's Digest of Materia Medica. 8vo. Half Calf, 4.00

Roberts' Compend of Materia Medica and Pharmacy. By the author of "Roberts' Practice." Cloth, 2.00

"It contains an immense amount of matter."—*The National Druggist.*

Headland's Action of Medicines. 9th Ed. 8vo. Cloth, 3.00

Waring. Therapeutics. A Practical Manual. Fourth Edition, revised and enlarged. *In Press.*

MEDICAL JURISPRUDENCE.

Reese. A Text-book of Medical Jurisprudence and Toxicology. By John J. Reese, M.D., Professor of Medical Jurisprudence and Toxicology in the Medical and Law Departments of the University of Pennsylvania; Vice-President of the Medical Jurisprudence Society of Philadelphia; Physician to St. Joseph's Hospital; Corresponding Member of The New York Medico-legal Society. Demi-Octavo. Cloth, 4.00; Leather, 5.00

"Professor Reese is so well known as a skilled medical jurist, that his authorship of any work virtually guarantees the thoroughness and practical character of the latter. And such is the case in the book before us. * * * * We might call these the essentials for the study of medical jurisprudence. The subject is skeletonized, condensed, and made thoroughly up to the wants of the general medical practitioner, and the requirements of prosecuting and defending attorneys. If any section deserves more distinction than any other, as to intrinsic excellence, it is that on toxicology. This part of the book comprises the best outline of the subject in a given space that can be found anywhere. As a whole, the work is everything it promises, and more, and considering its size, condensation, and practical character, it is by far the most useful one for ready reference, that we have met with. It is well printed and neatly bound."—*New York Medical Record.*

Abercrombie's Students' Guide to Medical Jurisprudence. 12mo. Cloth, 2.50

Mann's Manual of Psychological Medicine, and Allied Nervous Diseases, Their Diagnosis, Pathology and Treatment, and their Medico-Legal Aspects. Illustrated. 8vo.

Cloth, 5.00; Leather, 6.00

Woodman and Tidy's Medical Jurisprudence and Toxicology. Chromo-Lithographic Plates and 116 Wood engravings.

Cloth, 7.50; Leather, 8.50

 See pages 2 to 5 for list of ? Quiz-Compend?

MISCELLANEOUS.

- Beale. Slight Ailments.** Their Nature and Treatment. Illustrated. 8vo. Paper cover, .75; Cloth, 1.25
- Dulles. Surgical and other Emergencies.** Illustrated. Second Edition. 12mo. Cloth, .75
- Fothergill. Diseases of the Heart and Their Treatment.** Second Edition. 8vo. Cloth, 3.50
- Tanner. Memoranda of Poisons.** Their Antidotes and Tests. Fifth Edition. 12mo. Cloth, .75
- Allingham. Diseases of the Rectum.** Fourth Edition. Illustrated. 8vo. Paper covers, .75; Cloth, 1.25

OBSTETRICS AND GYNÆCOLOGY.

- Byford. The Diseases of Women.** By W. H. Byford, A.M., M.D., Professor of Gynæcology in Rush Medical College; of Obstetrics in the Woman's Medical College; and Surgeon to the Woman's Hospital, Chicago. Third Edition. Over 160 Illustrations. Octavo. Cloth, 5.00; Leather, 6.00

"The treatise is as complete a one as the present state of our science will admit of being written. We commend it to the diligent study of every practitioner and student, as a work calculated to inculcate sound principles and lead to enlightened practice."—*New York Medical Record*.

- Cazeaux and Tarnier. Obstetrics; the Theory and Practice of; including the Diseases of Pregnancy and Parturition, Obstetrical Operations, etc.** By P. Cazeaux, Member of the Imperial Academy of Medicine, etc. Revised, with additions, by S. Tarnier, Prof. of Obstetrics and Diseases of Women and Children in the Faculty of Medicine, of Paris. A New American, from the Eighth French and First Italian Editions. Edited and enlarged by Robert J. Hess, M.D., Physician to the Northern Dispensary, Philadelphia; Member of the College of Physicians of Philadelphia, etc. 1100 pages, 4to, with 12 Full-page Lithographic plates, 5 of which are Colored, and over 175 Wood Engravings.

Sold by subscription only. Full information and Four-page Circular upon application to the publishers.

"I have examined this edition of Cazeaux and Tarnier's Theory and Practice of Obstetrics, just from the publishing house of P. Blakiston, Son & Co., Philadelphia, and pronounce it practical and just what is needed by every practitioner. I highly recommend the work. It should be prominent in every library."—*T. Gaillard Thomas, M. D., Professor of Gynæcology in College of Physicians and Surgeons, New York*.

- Gallabin's Midwifery.** A New Manual for Students. Illustrated. *In Press.*

See pages 2 to 5 for list of ? Quiz-Compends ?

Obstetrics and Gynecology:—Continued.

- Meadows' Manual of Midwifery.** Including the Signs and Symptoms of Pregnancy, Obstetric Operations, Diseases of the Puerperal State, etc. 145 Illustrations. 494 pages. Cloth, 2.00
- Rigby's Obstetric Memoranda.** 4th Ed. 32mo. Cloth, .50
- Swayne's Obstetric Aphorisms.** For the use of Students commencing Midwifery Practice. 8th Ed. 12mo. Cloth, 1.25

PATHOLOGY AND HISTOLOGY.

- Rindfleisch's General Pathology.** For Students and Physicians. By Prof. Edward Rindfleisch, of Würzburg. Translated by Wm. H. Mercur, M.D., of Pittsburg, Pa., Edited by James Tyson, M.D., Professor of Pathology and Morbid Anatomy in the University of Pennsylvania. 12mo. Cloth, 2.00
- Gilliam's Essentials of Pathology.** A Handbook for Students. 47 Illustrations. 12mo. Cloth, 2.00

. The object of this book is to unfold to the beginner the fundamentals of pathology in a plain, practical way, and by bringing them within easy comprehension to increase his interest in the study of the subject. Though it will not altogether supplant larger works, it will be found to impart clear-cut conceptions of the generally accepted doctrines of the day, and to prevent confusion in the mind of the student.

- Gibbes' Practical Histology and Pathology.** Third Edition. Enlarged. 12mo. Cloth, 1.75

PHYSICAL DIAGNOSIS.

- Bruen's Physical Diagnosis of the Heart and Lungs.** By Dr. Edward T. Bruen, Assistant Professor of Clinical Medicine in the University of Pennsylvania. Second Edition, revised. With new Illustrations. 12mo. Cloth, 1.50

. The subject is treated in a plain, practical manner, avoiding questions of historical or theoretical interest, and without laying special claim to originality of matter, the author has made a book that presents to the student the somewhat difficult points of Physical Diagnosis clearly and distinctly.

PHYSIOLOGY.

- Yeo's Physiology.** The most Popular Students' Book. By Gerald F. Yeo, M.D., F.R.C.S., Professor of Physiology in King's College, London. Small Octavo. 750 pages. Over 300 carefully printed Illustrations. With a Full Glossary and Index.

Cloth, 4.00; Leather, 5.00

"The work will take a high rank among the smaller text-books of Physiology."—*Prof. H. P. Bowditch, Harvard Med. School, Boston.*

"The brief examination I have given it was so favorable that I placed it in the list of text-books recommended in the circular of the University Medical College."—*Prof. Lewis A. Stimpson, M. D., 37 East 33d Street, New York.*

See pages 2 to 5 for list of Quiz-Compends

Physiology:—Continued.

Kirke's Physiology. 11th Ed. Illus. Cloth, 4.00; Leather, 5.00

Landois' Human Physiology. Including Histology and Microscopical Anatomy. 2 volumes. Cloth, 10.00

"So great are the advantages offered by Prof. Landois' Text-book, from the exhaustive and eminently practical manner in which the subject is treated, that, notwithstanding it is one of the largest works on Physiology, it has yet passed through four large editions in the same number of years. Dr. Stirling's annotations have materially added to the value of the work. . . . Admirably adapted for the practitioner. . . . With this Text-book at his command, no student could fail in his examination."—*Lancet*.

Sanderson's Physiological Laboratory. Being Practical Exercises for the Student. 350 Illustrations. 8vo. Cloth, 5.00

Tyson's Cell Doctrine. Its History and Present State. Illustrated. Second Edition. Cloth, 2.00

PRACTICE.

Roberts' Practice. Fifth American Edition. A Handbook of the Theory and Practice of Medicine. By Frederick T. Roberts, M.D.; M.B.C.P., Professor of Clinical Medicine and Therapeutics in University College Hospital, London. Fifth Edition. Octavo. Cloth, 5.00; Leather, 6.00

* This new edition has been subjected to a careful revision. Many chapters have been rewritten. Important additions have been made throughout, and new illustrations introduced. Recommended as a Text-book at University of Pennsylvania, Long Island College Hospital, Yale and Harvard Colleges, Bishop's College, Montreal, University of Michigan, and over twenty other Medical Schools.

"I have become thoroughly convinced of its great value, and have cordially recommended it to my class in *Yale College*."—*Prof. David P. Smith*.

"I have examined it with some care, and think it a good book, and shall take pleasure in mentioning it among the works which may properly be put in the hands of students."—*A. B. Palmer, Prof. of the Practice of Medicine, University of Michigan*.

"A clear, yet concise, scientific and practical work. It is a capital compendium of the classified knowledge of the subject."—*Prof. J. Adams Allen, Rush Medical College, Chicago*.

"It is unsurpassed by any work that has fallen into our hands, as a compendium for students preparing for examination. It is thoroughly practical, and fully up to the times."—*The Clinic*.

Aitken's Practice of Medicine. Seventh Edition. 196 Illustrations. 2 vols. Cloth, 12.00; Leather, 14.00

Fagge's Principles and Practice of Medicine. A Complete Text-book. 2 vols. *Now Ready.*

Tanner's Index of Diseases, and Their Treatment. Cloth, 3.00

"This work has won for itself a reputation. . . . It is, in truth, what its Title indicates."—*N. Y. Medical Record*.

See pages 2 to 5 for list of ? Quiz-Compend ?

PRESCRIPTION BOOKS.

Wythe's Dose and Symptom Book. Containing the Doses and Uses of all the principal Articles of the *Materia Medica*, etc. Sixteenth edition. 32mo. Cloth, 1.00; Pocket-book style, 1.25

Pereira's Physician's Prescription Book. Containing Lists of Terms, Phrases, Contractions and Abbreviations used in Prescriptions, Explanatory Notes, Grammatical Construction of Prescriptions, etc., etc. By Professor Jonathan Pereira, M.D. Sixteenth Edition. 32mo. Cloth, 1.00; Pocket-book style, 1.25

SKIN DISEASES.

Van Harlingen on Skin Diseases. A Handbook of the Diseases of the Skin, their Diagnosis and Treatment. By Arthur Van Harlingen, M.D., Prof. of Diseases of the Skin in the Philadelphia Polyclinic; Consulting Physician to the Dispensary for Skin Diseases, etc. With colored plates. 12mo. Cloth, 1.75

*** This is a complete epitome of skin diseases, arranged in alphabetical order, giving the diagnosis and treatment in a concise, practical way. Many prescriptions are given that have never been published in any text-book, and an article incorporated on Diet. The plates do not represent one or two cases, but are composed of a number of figures, accurately colored, showing the appearance of various lesions, and will be found to give great aid in diagnosing.

"This new handbook is essentially a small encyclopædia. *** We heartily commend it for its brevity, clearness and evidently careful preparation."—*Philadelphia Medical Times*.

"This is an excellent little book, in which, for ease of reference, the more common diseases of the skin are arranged in alphabetical order, while many good prescriptions are given, together with clear and sensible directions as to their proper application."—*Boston Medical and Surgical Journal*.

Bulkley. The Skin in Health and Disease. By L. Duncan Bulkley, Physician to the N. Y. Hospital. Illus. Cloth, .50


SURGERY.

Heath's Minor Surgery, and Bandaging. New Edition. With many Illustrations. *In Press.*

Mears' Practical Surgery. Second Edition. Enlarged. 490 Illustrations. Cloth, \$3.75; Leather, \$4.75

Pye's Surgical Handicraft. A Manual of Surgical Manipulations, Minor Surgery, Bandaging, Dressing, etc., etc. With special chapters on Aural Surgery, Extraction of Teeth, Anæsthetics, etc. 208 Illustrations. 8vo. Cloth, 5.00

Watson on Amputation of the Extremities, and their Complications. 2 colored plates and 250 wood cuts. 8vo. Cloth, 5.50

 See pages 2 to 5 for list of ?Quiz-Compends?

THROAT.

Mackenzie on the Throat and Nose. By Morell Mackenzie, M.D., Senior Physician to the Hospital for Diseases of the Chest and Throat; Lecturer on Diseases of the Throat at the London Hospital, etc.

Vol. I. Including the Pharynx, Larynx, Trachea, etc., with Formulæ and 112 Illustrations.

Vol. II. Diseases of the Œsophagus, Nose and Naso-Pharynx, with Formulæ and 93 Illustrations.

The two volumes, Cloth, 6.00; Leather, 7.50
Vol. II, separately, Cloth, 3.00; Leather, 4.00

"It is both practical and learned; abundantly and well illustrated; its descriptions of disease are graphic and the diagnosis the best we have anywhere seen."—*Philadelphia Medical Times*.

Cohen. The Throat and Voice. Illustrated. Cloth, .50

James. Sore Throat. Its Nature, Varieties and Treatment. 12mo. Illustrated. Paper cover, .75; Cloth, 1.25

URINE AND URINARY ORGANS.

Acton. The Reproductive Organs. In Childhood, Youth Adult Life and Old Age. Sixth Edition. Cloth, 2.00

Beale. Urinary and Renal Diseases and Calculous Disorders. Hints on Diagnosis and Treatment. 12mo. *Just Ready.* Cloth, 1.75

Ralfe. Kidney Diseases and Urinary Derangements. 42 Illustrations. 12mo. 572 pages. *Just Ready.* Cloth, 2.75

Legg. On the Urine. A Practical Guide. Sixth Edition. 12mo. Cloth, .75

Marshall and Smith. On the Urine. The Chemical Analysis of the Urine. By John Marshall, M.D., Chemical Laboratory, University of Pennsylvania, and Prof. E. F. Smith, Ph.D. With Colored Plates. Cloth, 1.00

Thompson. Diseases of the Urinary Organs. Seventh Edition. Illustrated. Cloth, 1.25

Tyson. On the Urine. A Practical Guide to the Examination of Urine. For Physicians and Students. By James Tyson, M.D., Professor of Pathology and Morbid Anatomy, University of Pennsylvania. With Colored Plates and Wood Engravings. Fifth Edition. Revised and Enlarged. 12mo. Cloth, 1.50

Durkee. On Gonorrhœa and Syphilis. Illus. Cloth, 3.50

☞ See pages 2 to 5 for list of ? Quiz-Compend? ?

Richter's Chemistries.

AUTHORIZED TRANSLATIONS.

By EDGAR F. SMITH, M.A., Ph.D.,

*Prof. of Chemistry in Wittenberg College, Springfield, Ohio;
formerly in the Laboratories of the University of Pennsylvania
and Muhlenburg College; Member of the Chemical
Societies of Berlin and Paris; of the Academy of
Natural Sciences of Philadelphia, etc., etc.*

EACH VOLUME SOLD SEPARATELY.

INORGANIC CHEMISTRY. Second American,
from the Fourth German Edition; thoroughly revised
and in many parts rewritten. With 89 Illustrations
and Colored Plate of Spectra. Cloth, \$2.00

**THE CHEMISTRY OF THE CARBON COM-
POUNDS,** or Organic Chemistry. First Ameri-
can, from Fourth German Edition. Illustrated.

Cloth, \$3.00

The success attending the publication of the first edition of Richter's Inorganic Chemistry encourages the translator and publishers to believe that the companion volume will have an equally warm reception. Professor Richter's methods of arrangement and teaching have proved their superiority, abroad, by the very large sale of his books all over the Continent, translations having been made in Germany, Russia, Holland and Italy.

From PROF. B. SILLIMAN, Yale College, New Haven, Conn.

"It is decidedly a good book, and in some respects the best manual we have."

From JOHN MARSHALL, M.D., NAT. SC. D. (Tübingen), Demonstrator of Chemistry in the University of Pennsylvania, Medical Department.

"The work is of undoubted value. The theory of chemistry, which is generally the bugbear of students, is, in this book, very clearly explained, and the explanations are so well distributed through the book that students are brought easily from the simplest to the most difficult problems.

"That part descriptive of the elements and their compounds is full, and all that could be desired in a text-book, while the cuts, with which the work is profusely illustrated, are an excellent aid to the student. Altogether, it is one of our best modern works on chemistry."

Yeo's Physiology.

A MANUAL FOR STUDENTS.

By **GERALD F. YEO, M.D., F.R.C.S.,**
Professor of Physiology in King's College, London.

WITH OVER 300 CAREFULLY PRINTED ILLUSTRATIONS AND A GLOSSARY.

Small Octavo. Cloth, \$4.00; Leather, \$5.00.

RECOMMENDATIONS.

"After a careful examination of this manual of Physiology, I can truthfully say that it is a most valuable addition to the list of text-books upon this subject. That it should and will receive a welcome from both students and teachers there can be no doubt; for, in addition to the familiar but well presented facts of most text-books, it contains all the more important facts of physiological science which have been established in the last few years. The author presents his subject in a manner that is clear, concise and logical. Each section has had a careful revision, and reveals the author's familiarity with the scope and tendencies of modern physiology. It will prove an interesting and instructive book to those commencing the study of this subject."—*A. P. Brubaker, M.D., Demonstrator of Physiology at Jefferson Medical College, Philadelphia.*

"The writer has endeavored to 'avoid theories which have not borne the test of time, and such details of methods as are unnecessary for junior students.' His experience as a teacher and as an examiner has led him to lay stress on the points which are hard to grasp and are commonly misunderstood, and he has directed attention more to those subjects which have a practical medical or surgical application than to those relating to abstract physiological subjects. * * * We have pleasure in recommending this book, as a most excellent manual, being what it pretends to be—elementary, and yet containing all that is really of importance to the student of medicine. It is difficult to devise an original method of treating such a well worn subject as physiology, but the present volume undoubtedly has originality in its method. We think the author treats his subject in the best manner possible when he combines microscopical anatomy intimately with the discussion of strict physiology."—*The Medical Times and Gazette.*

VAN HARLINGEN

On Skin Diseases.

WITH COLORED ILLUSTRATIONS.

A HANDBOOK OF THE DISEASES OF THE SKIN, their Diagnosis and Treatment. By ARTHUR VAN HARLINGEN, M.D., Professor of Diseases of the Skin in the Philadelphia Polyclinic, Consulting Physician to the Dispensary for Skin Diseases, etc. Illustrated by colored lithographic plates.

12mo. 284 pages. Cloth. Price, \$1.75.

RECOMMENDATIONS.

"It is a most useful compendium of the knowledge to be had at the present time upon the important subjects to which it is devoted; and is in all respects a credit to the well recognized abilities of its author."—*James Nevins Hyde, M.D., Professor of Skin and Venereal Diseases, Rush Medical College, Chicago.*

"This new handbook is essentially a small encyclopedia of pathology and treatment of Skin Diseases, in which the subjects are arranged alphabetically. This arrangement was that followed by the late Tilbury Fox, of London, in his handbook, which we believe was remarkably successful; and we have no doubt it will be equally appreciated in the present work, which (compendious in form) contains a very complete summary of the present state of dermatology. Dr. Van Harlingen's position in the profession, being at present vice-president of the American Dermatological Association, which he served as secretary for several years, and the high standard of his communications to his department, are sufficient to warrant the confidence in his teachings, which is fully sustained by an examination of this handbook, which we heartily commend for its brevity, clearness and evident careful preparation."—*Philadelphia Medical Times, October 18th, 1884.*

RINDFLEISCH.

The Elements of Pathology.

A Text-Book in the University of Pennsylvania.

THE ELEMENTS OF PATHOLOGY. For Students and Physicians.

By EDWARD RINDFLEISCH, M.D., Professor of Pathological Anatomy in the University of Wurzburg. Authorized translation by WILLIAM H. MERCUR, M.D., of Pittsburgh, Pa. Revised by JAMES TYSON, M.D., Professor of General Pathology and Morbid Anatomy in the University of Pennsylvania.

12mo. 263 pages. Cloth, \$2.00.

RECOMMENDATIONS.

"The practical views of one of the best of the modern histologists is placed before the profession in this admirable work, in a most careful and systematic manner. The author, who is one of the leading pathologists, sets forth not only the ground-work in his department, but treats and makes clear many of the more difficult points of the study of pathological processes. The work is divided into a consideration, *first*, of the local outbreak of diseases; *second*, into the anatomical extension of disease; *thirdly*, into the physiological extension of disease, and lastly, into an examination of special parts."—*The Medical Bulletin.*

P. BLAKISTON, SON & CO., Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

BIDDLE'S Materia Medica.

TENTH REVISED EDITION.

Contains all Changes in the New Pharmacopœia.

Recommended as a Text-book at Yale College, University of Michigan, College of Physicians and Surgeons, Baltimore, Baltimore Medical College, Louisville Medical College, and a number of other Colleges throughout the United States.

BIDDLE'S MATERIA MEDICA, For the Use of Students and Physicians. By the late PROF. JOHN B. BIDDLE, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Tenth Edition, thoroughly revised, and in many parts rewritten, by his son, CLEMENT BIDDLE, M.D., Assistant Surgeon, U. S. Navy, assisted by HENRY MORRIS, M.D., Demonstrator of Obstetrics, Jefferson Medical College, Philadelphia. Containing all the additions and changes made in the last revision of the United States Pharmacopœia. 8vo.

Bound in Cloth. Price \$4.00; Leather, \$4.75.

RECOMMENDATIONS.

"It will be found a useful handbook by students, especially, who may be under the instruction of its able and accomplished author."—*American Medical Journal*.

"In short, it is just the work for a student, embracing as it does what will be discussed in a course of lectures on materia medica."—*Cincinnati Medical News*.

"In truth, the work is well adapted to the wants of students."—*The Clinic*.

"Nothing has escaped the writer's scan. All the new remedies against disease are duly and judiciously noted. Students will certainly appreciate its shapely form, grace of manner, and general *multum in parvo* style."—*American Practitioner*.

"Biddle's 'Materia Medica' is well known to the profession, being a standard text-book in several leading colleges."—*New York Medical Journal*.

"It contains, in a condensed form, all that is valuable in materia medica, and furnishes the medical student with a complete manual on this subject."—*Canada Lancet*.

"The necessity for a new edition of this work in so short a time is the best proof of the value in which it is held by the profession."—*Medical and Surgical Reporter*.

"The standard 'Materia Medica' with a large number of medical students is Biddle's."—*Buffalo Medical and Surgical Journal*.

"The larger works usually recommended as text-books in our medical schools are too voluminous for convenient use. This work will be found to contain in a condensed form all that is most valuable, and will supply students with a reliable guide."—*Chicago Medical Journal*.

* * This Ninth Edition contains all the additions and changes in the U. S. Pharmacopœia, Sixth Revision.

P. BLAKISTON, SON & CO., Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

REESE'S Medical Jurisprudence AND TOXICOLOGY.

A TEXT-BOOK FOR STUDENTS AND PHYSICIANS. By JOHN J. REESE, M.D., Professor of Medical Jurisprudence and Toxicology in the University of Pennsylvania; Vice-President of the Medical Jurisprudence Society of Philadelphia; Member of the College of Physicians of Philadelphia; Corresponding Member of the New York Medico-Legal Society 606 pages. Demi-octavo.

Bound in Cloth, \$4.00; Leather, \$5.00.

RECOMMENDATIONS.

"I have just concluded a careful review of Dr. John J. Reese's 'Text-Book of Medical Jurisprudence and Toxicology,' and I take great pleasure in saying that it is by far the best book of its size on this subject which I have ever seen. * * * It is, for the medical student and the general practitioner, a much more convenient, readable, and in most respects better book than the voluminous treatises, to say nothing of the fact that it is a much less costly book than are these."—*Willis G. Tucker, M. D., Professor of Inorganic and Analytical Chemistry and Medical Jurisprudence, Albany Medical College.*

"There has long been needed a work like this one. The student of the subject has had to choose between very prolix and expensive treatises by American writers, or those prepared by foreigners, whose customs and laws differ so much from ours that it materially detracts from their usefulness.

"In this single volume, printed in clear type and on excellent paper, Dr. Reese presents all that the student or the general practitioner will have occasion to learn about the subject."—*The Medical and Surgical Reporter, Philadelphia, October 11th, 1884.*

"I have examined the work, and find it to be a well compiled manual of the various doctrines obtaining, both in law and in medicine, upon the subject of which it treats. In a field so wide as that of medical jurisprudence, it is a matter largely of individual choice as to the scope which shall be given by an author to the treatment of its several topics. Professor Reese seems to me to have struck the golden mean in this respect, and to have combined lucidity of style with brevity of statement, so as to give to each topic its due proportion of development. It cannot fail to become a most useful handbook for students who desire to have in outline the whole body of medical jurisprudence, including even the special department of toxicology, with all which that now embraces."—*John Ordronaux, Professor of Medical Jurisprudence in Columbia College, New York, and in Dartmouth College, Hanover, N. H.*

"It may be called a practical encyclopædia, giving just those things which are wanted in medico-legal inquiries. * * * The thoroughness and skill of the author is apparent on every page, and the work may be especially commended as the best single volume on this subject in print."—*Quarterly Journal of Inebriety, October, 1884.*

"It not only treats of the medical side of the subject, but gives much valuable legal information as to the rights and duties of physicians in such cases."—*Popular Science News, October, 1884.*

"It comes to the profession at an opportune time, as a valuable addition to the subject of State medicine, and the volume should be in the hands of every medical man."—*Prof. James F. Harrison, University of Virginia.*

"We might call these the essentials for the study of medical jurisprudence. * * * * If any section deserves more distinction than any other, as to intrinsic excellence, it is that on toxicology. This part of the book comprises the best outline of the subject in a given space that can be found anywhere. As a whole, the work is everything it promises and more, and considering its size, condensation and practical character, it is by far the most useful one for ready reference that we have met with. It is well printed and neatly bound."—*N. Y. Medical Record, Sept. 13th, 1884.*

P. BLAKISTON, SON & CO., Publishers and Booksellers,
1012 WALNUT STREET, PHILADELPHIA.

NATIONAL LIBRARY OF MEDICINE



NLM 00052909 3